

**NUCLEIC ACID MOLECULES AND OTHER MOLECULES ASSOCIATED WITH
THE CYTOKININ PATHWAY**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C §119(e) and/or 35 U.S.C §120 of applications No. 60/067000 filed November 24, 1997; No. 60/069472 filed December 9, 1997; No. 60/071,064 filed January 9, 1998; No. 60/074,201 filed February 10, 1998; No. 60/074281 filed February 10, 1998; No. 60/074567 filed February 12, 1998; No. 60/074565 filed February 12, 1998; No. 60/075462 filed February 19, 1998; No. 60/075461 filed February 19, 1998; No. 60/075464 filed February 19, 1998; No. 60/075460 filed February 19, 1998; No. 60/075463 filed February 19, 1998; No. 60/077231 filed March 9, 1998; No. 60/077229 filed March 9, 1998; No. 60/077230 filed March 9, 1998; No. 60/078368 filed March 18, 1998; No. 60/080844 filed April 7, 1998; No. 60/083067 filed April 27, 1998; No. 60/083387 filed April 29, 1998; No. 60/083388 filed April 29, 1998; No. 60/085224 filed May 13, 1998; No. 60/085223 filed May 13, 1998; No. 60/085222 filed May 13, 1998; No. 60/086186 filed May 21, 1998; No. 60/086187 filed May 21, 1998; No. 60/086185 filed May 21, 1998; No. 60/086184 filed May 21, 1998; No. 60/086188 filed May 21, 1998; No. 60/089,524 filed June 16, 1998; No. 60/089,810 filed June 18, 1998; No. 60/089,814 filed June 18, 1998; No. 60/090,170 filed June 22, 1998; No. 60/092,036 filed July 8, 1998; No. 60/099670 filed September 9, 1998; No. 60/099697 filed September 9, 1998; No. 60/100674 filed September 16, 1998; No. 60/101132 filed September 21, 1998; No. 60/101130 filed September 21, 1998; No. 60/101,508 filed September 22, 1998; No. 60/101344 filed September 22, 1998; No. 60/101347 filed September 22, 1998; No. 60/101343 filed September 22, 1998; No. 60/104,126 filed October 13, 1998; No. 60/104,127 filed October 13, 1998; No. 60/104,124 filed October 13, 1998; No. 60/104,121 filed October 13, 1998; "Nucleic

Acid Molecules and Other Molecules Associated With Plants" docket No. 38-21(15075)B filed November 24, 1998; "Nucleic Acid Molecules and Other Molecules Associated With Plants" docket No. 38-21(15076)B filed December 8, 1998; and "Nucleic acid Molecules and other Molecules associated with Plants" docket No. 38-21(15668)A filed December 11, 1998, all of which are herein incorporated by reference in their entirety.

FIELD OF THE INVENTION

The present invention is in the field of plant biochemistry. More specifically the invention relates to nucleic acid sequences from plant cells, in particular, nucleic acid sequences from maize and soybean plants associated with the cytokinin pathway. The invention encompasses nucleic acid molecules that encode proteins and fragments of proteins. In addition, the invention also encompasses proteins and fragments of proteins so encoded and antibodies capable of binding these proteins or fragments. The invention also relates to methods of using the nucleic acid molecules, proteins and fragments of proteins and antibodies, for example for genome mapping, gene identification and analysis, plant breeding, preparation of constructs for use in plant gene expression and transgenic plants.

BACKGROUND OF THE INVENTION

Plant hormones, produced in response to genetic, environmental or chemical stimuli (Goldberg, *Science* 240: 1460-1467 (1988); Letham, In: *Phytohormones and Related Compounds—A Comprehensive Treatise*, eds. Letham et al., Amsterdam, Elsevier North Holland. 1: 205-263 (1978); von Sachs, *Arb. Bot. Inst. Wurzburg* 2:452-488 (1880), all of which are herein incorporated by reference in their entirety), play a role in controlling the growth, development and environmental responses of plants.

Cytokinins are a class of plant hormones with a structure resembling adenine.

Cytokinins, in combination with auxin, promote cell division. Cytokinins are associated with many aspects of plant growth and development (Horgan, *Advanced Plant Physiology*, ed. Wilkins, Pitman, London: 90-116 (1984); Skoog *et al.*, *Biochemical Actions of Hormones*, ed. Litwack, Academic Press, London, vol. VI: 335-413 (1979), all of which are herein incorporated by reference in their entirety). Cytokinins have been found in almost all higher plants as well as mosses, fungi, and bacteria. In addition to occurring in higher plants as free compounds, cytokinins may also occur as component nucleosides in tRNA of plants, animals, and microorganisms.

Kinetin, the first cytokinin to be discovered, was so named because of its ability to promote cytokinesis (cell division). Although kinetin is a natural compound, it is not made in plants, and is therefore usually considered a "synthetic" cytokinin. Two common forms of cytokinin in plants are zeatin and zeatin riboside (maize)(Letham, *Life Sci.* 2: 569-573 (1963), the entirety of which is herein incorporated by reference). More than 200 known natural and synthetic cytokinins have been reported.

Several cytokinin related mutations have also been reported. For example, the *ckr1* mutant of *Arabidopsis* is resistant to the cytokinin bezyladenine (Su and Howell, *Plant Physiol.* 99:1569-1574 (1992), the entirety of which is herein incorporated by reference). The *Arabidopsis* mutant *amp1* has been reported to be a negative regulator of cytokinin biosynthesis (Chadbury *et al.*, *Plant J.* 4:907-916 (1993), the entirety of which is herein incorporated by reference).

Cytokinin concentrations are highest in meristematic regions and areas of continuous growth potential such as roots, young leaves, developing fruits, and seeds (Arteca, *Plant Growth Substances: Principles and Applications*, eds. Chapman & Hall, New York (1996); Mauseth, *Botany: An Introduction to Plant Biology*, ed. Saunders, Philadelphia: 348-415 (1991); Raven *et al.*, *Biology of Plants*, ed. Worth, New York: 545-572 (1992); Salisbury and Ross, *Plant*

Physiology, ed. Wadsworth, Belmont, CA: 357-407, 531-548 (1992), all of which are herein incorporated by reference in their entirety).

It has been reported that the induced cytokinin response varies depending on the type of cytokinin and plant species (Davies, *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, Kluwer, Dordrecht (1995); Mauseth, *Botany: An Introduction to Plant Biology*, Saunders, Philadelphia: 348-415 (1991); Raven *et al.*, *Biology of Plants*, ed. Worth, New York: 545-572 (1992); Salisbury and Ross, *Plant Physiology*, ed. Wadsworth, Belmont, CA: 357-407, 531-548 (1992), all of which are herein incorporated by reference in their entirety). Elevated cytokinin levels are associated with the development of seeds in higher plants, and have been demonstrated to coincide with maximal mitotic activity in the endosperm of developing maize kernels, cereal grains, and fruits. Exogenous cytokinin application (via stem injection) has been shown to directly correlate with increased kernel yield in maize. In addition, plant cells transformed with the *ipt* gene from *Agrobacterium tumefaciens* showed increased growth corresponding to an increase in endogenous cytokinin levels upon induction of the enzyme. Cytokinins have been reported to confer thermotolerance in certain physiological processes such as plastid biogenesis and endosperm cell division (Cheikh and Jones, *Plant Physiol.* 106: 45-51 (1994); Parthier, *Biochem. Physiol Pflanz* 174:173-214 (1979); Jones *et al.*, *Crop Science* 25: 830-834 (1985), all of which are herein incorporated by reference in their entirety).

Reviews of cytokinin metabolism, compartmentalization, conjugation and cytokinin metabolic enzymes have been presented by Jameson, *Cytokinins*, eds. Mok and Mok, Boca Raton, Fla, 113-128 (1994); Letham and Palni, *Ann. Rev. Plant Physiol.* 34: 163-197 (1983); McGaw *et al.* In: *Biosynthesis and metabolism of plant hormones*, Soc. Exp. Biol. Seminar Series, eds. Crozier and Hillman, Cambridge University Press, Cambridge, Vol. 23, chapter 5 (1984); McGaw and Horgan, *Biol. Plant* 27: 180 (1985); McGaw *et al.*, In: *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, ed. Davies, Kluwer, Dordrecht, 98-117 (1995); Mok and Martin, *Cytokinins*, eds. Mok and Mok, Boca Raton, Fla, 129-137 (1994); Salisbury

and Ross, *Plant Physiology*, Belmont, CA: ed. Wadsworth, 357-407, 531-548 (1992), all of which are hereby incorporated by reference in their entirety.

I. BIOSYNTHESIS OF CYTOKININS

Cytokinins are generally found in higher concentrations in meristematic regions and growing tissues. It has been reported that cytokinins are synthesized in the roots and translocated via the xylem to the meristematic regions and growing shoots of the plant. Although cytokinin biosynthesis in developed plants takes place mainly in roots (Engelbrecht, *Biochem. Physiol. Pflanzen* 163: 335-343 (1972); Henson *et al.*, *J. Exp. Bot* 27: 1268-1278 (1976); Sossountzov *et al.*, *Planta* 175: 291-304 (1988); Van Staden *et al.*, *Ann. Bot.* 42: 751-753 (1978), all of which are herein incorporated by reference in their entirety), smaller amounts can be synthesized by the shoot apex and some other plant tissues.

The level of active cytokinin at a particular site of action has been reported to be influenced by a large number of factors: de novo synthesis; oxidative degradation; reduction; formation and hydrolysis of inactive conjugates; transport into and out of particular cells; subcellular compartmentalization to or away from sites of action. It has also been reported that physiological responses may be modulated by variations in the ability of cells to respond to a particular concentration of free cytokinin.

Cytokinin biosynthesis happens through the biochemical modification of adenine (McGaw *et al.*, In: *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, ed. Davies, Kluwer, Dordrecht: 98-117 (1995), the entirety of which is herein incorporated by reference; Salisbury and Ross, *Plant Physiology*, Belmont, CA: ed. Wadsworth, 357-407, 531-548 (1992), the entirety of which is herein incorporated by reference). Plants appear to synthesize cytokinins either directly by addition of isopentenylpyrophosphate to AMP by an adenylate:isopentenyltransferase (cytokinin synthase) producing isopentenyladenosine 5' phosphate ("[9R-5'P]iP"), which in turn serves as an intermediate for further modifications, or indirectly via isopentenylation of adenosine residues of tRNA by tRNA:isopentenyltransferase

(McGaw *et al.*, *In: Plant Hormones: Physiology, Biochemistry and Molecular Biology*, ed. Davies, Kluwer, Dordrecht: 98-117 (1995)). [9R-5'P]iP may be modified by dephosphorylation, deribosylation, hydroxylation and reduction to produce a variety of derivatives with potential activity (Binns, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 45: 173-196 (1994), the entirety of which is herein incorporated by reference). Further, conjugation may modulate levels of active cytokinins (Letham and Palni, *Ann. Rev. Plant Physiol.* 34: 163-197 (1983), the entirety of which is herein incorporated by reference).

In the biosynthesis of tRNA cytokinins, mevalonic acid pyrophosphate undergoes decarboxylation, dehydration and isomerization to yield 2-isopentyl pyrophosphate ("iPP"). iPP then condenses with the relevant adenosine residue in the tRNA to give the N6(Δ 2-isopentenyl)adenosine ("[9R]iP") moiety. With the exception of [9R]iP and to a lesser extent cis- and trans-[9R]Z, the free and tRNA cytokinins are structurally distinct (e.g., free Zeatin ("Z") is mainly the trans isomer (trans-Zeatin while Z present in tRNA is mainly the cis isomer (McGaw *et al.*, *In: Plant Hormones: Physiology, Biochemistry and Molecular Biology*, ed. Davies, Kluwer, Dordrecht, 98-117 (1995)).

The de novo biosynthesis pathway of cytokinins in plants includes the following enzymes: isopentyltransferase, 5'-nucleosidase, adenine nucleotidase, adenine phosphorylase, adenine kinase, adenine phosphoribosyl transferase, microsomal mixed function oxidases, Zeatin reductase, O-glucosyltransferase, O-xylosyltransferase, β -(9-cytokinin-alanino)synthase, cytokinin oxidase, β -glucosidase, and Zeatin cis-trans isomerase.

Isopentyltransferase catalyzes the first reaction of the pathway in which N6(Δ 2-isopentenyl) adenosine-5'-monophosphate ("[9R-5'P]iP") is generated from iPP and AMP.

5'-nucleotidase catalyzes the conversion of [9R-5'P]iP to [9R]iP. The reaction catalyzed by the enzyme 5'-nucleotidase has been found in wheat germ extract (Chen *et al.*, *Plant Physiol.* 67:494-498 (1981); Chen *et al.*, *Plant Physiol.* 68:1020-1023 (1981), both of which are herein incorporated by reference in their entirety) and in tomato leaf and root extracts (Burch and Stuchbury, *Phytochemistry* 25:2445-2449 (1986); Burch and Stuchbury, *J. Plant Physiol.*

125:267-273 (1986), both of which are herein incorporated by reference in their entirety).

Adenine kinase catalyzes the reversion of [9R]iP to [9R-5'P]iP. Alternatively, [9R-5'P]iP can be converted to t-Zeatin riboside-5'-monophosphate ("[9R-5'P]Z") by a microsomal mixed function oxidase.

Adenosine nucleotidase catalyzes the conversion of [9R]iP to iP. This reaction can be reversed by the enzyme adenine phosphorylase. Alternatively, [9R]iP can be converted to t-Zeatin riboside ("[9R]Z") by a microsomal mixed function oxidase. Under another reaction mechanism, adenosine can be cleaved from [9R]iP by cytokinin oxidase. The enzyme adenine phosphoribosyl transferase can catalyze the conversion of iP to [9R-5'P]iP. Adenine phosphoribosyl transferase which is one of the salvage routes in plants for converting adenosine to AMP has also been shown to catalyze the phosphoribolyzation of cytokinin bases from a number of plant sources, including wheat germ (Chen *et al.*, *Arch. Biochem. Biophys.* 214:634-641 (1982), the entirety of which is herein incorporated by reference), tomato (Burch *et al.*, *Physiol. Plant* 69:283-288 (1987), the entirety of which is herein incorporated by reference), *A. thaliana* (Moffatt *et al.*, *Plant Physiol.* 95:900-908 (1991), the entirety of which is herein incorporated by reference) and *Acer pseudoplatanus* (Doree and Guern, *Biochem. Biophys. Acta* 304:611-622 (1973); Sadorge *et al.*, *Physiol. Veg.* 8:499-514 (1970), both of which are herein incorporated by reference in their entirety).

The cytokinins N6(Δ^2 -isopentenyl) adenosine-7-glucoside ("[7G]iP") and N6(Δ^2 -isopentenyl) adenosine-9-glucoside ("[9G]iP") are generated from iP from the enzymes Zeatin reductase and O-glucosyltransferase (such as cytokinin-9-glucosyl transferase), respectively. Under another reaction mechanism, adenine can be cleaved from iP by cytokinin oxidase.

In addition to converting [9R-5'P]iP to [9R]iP, 5'-nucleotidase can also catalyze the conversion of [9R-5'P]Z to [9R]Z. Adenine kinase can catalyze the conversion of [9R]Z to [9R-5'P]Z.

O-glucosyltransferase catalyzes the conversion of [9R]Z to t-Zeatin riboside-O-glucoside ("(OG)[9R]Z"). O-glucosyltransferase can also remove the glucoside group from (OG)[9R]Z to

regenerate [9R]Z. Adenosine can be cleaved from [9R]Z by cytokinin oxidase. Alternatively, adenine nucleotidase can convert [9R]Z to Z. Adenine phosphorylase can catalyze the conversion of Z back into [9R]Z.

The cytokinins dihydroZeatin (“(diH)Z”), Zeatin-7-glucoside ([7G]Z), Zeatin-9-glucoside (“[9G]Z”), and lupinic acid (“[9Ala]Z”) are generated from Z by the enzymes Zeatin reductase, O-glucosyltransferase, Zeatin reductase and β -(9-cytokinin alanino) synthase, respectively. Zeatin cis-trans isomerase catalyzes the isomerization of Zeatin between its cis and trans isomers. O-glucosyltransferase catalyzes the addition of a glucoside residue to Z to form t-Zeatin-O-glucoside (“(OG)Z”) or removal of a glucoside residue from (OG)Z to form Z.

The cytokinins dihydroZeatin-9-glucoside (“(diH)[9G]Z”), dihydroZeatin-7-glucoside (“(diH)[7G]Z”), and dihydrolupinic acid (“(diH)[9Ala]Z”) are generated from (diH)Z by the enzymes β -(9-cytokinin alanino)synthase, Zeatin reductase, and O-glucosyltransferase, respectively. O-glucosyltransferase catalyzes the addition of a glucoside residue to (diH)Z to form t-Zeatin-O-glucoside (“(diHOG)Z”) or removal of a glucoside residue from (diHOG)Z to form (diH)Z. Alternatively, (diH)Z can be converted into dihydroZeatin riboside ((diH)[9R]Z) by adenine phosphorylase. The enzyme adenine nucleotidase can catalyze the conversion of (diH)[9R]Z to (diH)Z.

O-glucosyltransferase catalyzes the addition of a glucoside residue to (diH)[9R]Z to form t-dihydroZeatin riboside-O-glucoside (“(diHOG)[9R]Z”) or the removal of a glucoside residue from (diHOG)[9R]Z to form (diH)[9R]Z. The cytokinin dihydroZeatin riboside-5'-monophosphate (“(diH)[9R-5'P]Z”) is generated from (diH)[9R]Z by the enzyme adenine kinase. This reaction can be reversed by the enzyme 5'-nucleotidase.

It is understood that the above description of the de novo biosynthesis of cytokinins only describes the core of the biosynthesis pathway. Other enzymes have been reported to be involved in this pathway.

Active cytokinins can be inactivated by degradation or conjugation to different low-molecular-weight metabolites, such as sugars and amino acids. The enzyme cytokinin oxidase

plays a role in the degradation of cytokinins. This enzyme removes the side chain and releases adenine, the backbone of all cytokinins. Cytokinin oxidases are reported to remove cytokinins from plant cells after cell division. Cytokinin derivatives are also made.

β -glucosidase (EC 3.2.1.21) has been reported to cleave the biologically inactive hormone conjugates of cytokinin-O-glucoside to release the active cytokinin (Brzobohaty et al., *Science* 262:1051-1054 (1993); Campos *et al.*, *Plant J.* 2:675-684 (1992), both of which are herein incorporated by reference in their entirety). β -glucosidase catalyzes the hydrolysis of aryl and alkyl β -D-glucosides and/or cellobiose with the release of β -D-glucose (Reese, *Recent Adv. Phytochem.* 11:311 (1977), the entirety of which is herein incorporated by reference). The enzyme has been purified from maize and has a molecular weight of 60 kD (Esen, *Plant Physiol.* 98:174-182 (1992); Esen *et al.*, *Biochem. Genet.* 28:319-336 (1990), both of which are herein incorporated by reference). Esen et al. have identified the rolC gene of *Agrobacterium rhizogenes* which encodes for a cytokinin β -glucosidase and which effects the growth and development of transgenic plants (Esen *et al.*, *EMBO J.* 10:2889-2895 (1991), the entirety of which is herein incorporated by reference).

Conjugation is often reported as a way of removing free and active hormones from a tissue. The conjugation process is often reversible, and, as conjugates can frequently accumulate in excess of free forms of phytohormone. The conjugate pools are also considered as sources of free hormone and may represent storage or inactive transportable forms of the hormone.

II. EXPRESSED SEQUENCE TAG NUCLEIC ACID MOLECULES

Expressed sequence tags, or ESTs are randomly sequenced members of a cDNA library (or complementary DNA)(McCombie *et al.*, *Nature Genetics* 1:124-130 (1992); Kurata *et al.*, *Nature Genetics* 8:365-372 (1994); Okubo *et al.*, *Nature Genetics* 2:173-179 (1992), all of which references are incorporated herein in their entirety). The randomly selected clones comprise insets that can represent a copy of up to the full length of a mRNA transcript.

Using conventional methodologies, cDNA libraries can be constructed from the mRNA (messenger RNA) of a given tissue or organism using poly dT primers and reverse transcriptase (Efstratiadis *et al.*, *Cell* 7:279-3680 (1976), the entirety of which is herein incorporated by reference; Higuchi *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 73:3146-3150 (1976), the entirety of which is herein incorporated by reference; Maniatis *et al.*, *Cell* 8:163-182 (1976) the entirety of which is herein incorporated by reference; Land *et al.*, *Nucleic Acids Res.* 9:2251-2266 (1981), the entirety of which is herein incorporated by reference; Okayama *et al.*, *Mol. Cell. Biol.* 2:161-170 (1982), the entirety of which is herein incorporated by reference; Gubler *et al.*, *Gene* 25:263-269 (1983), the entirety of which is herein incorporated by reference).

Several methods may be employed to obtain full-length cDNA constructs. For example, terminal transferase can be used to add homopolymeric tails of dC residues to the free 3' hydroxyl groups (Land *et al.*, *Nucleic Acids Res.* 9:2251-2266 (1981), the entirety of which is herein incorporated by reference). This tail can then be hybridized by a poly dG oligo which can act as a primer for the synthesis of full length second strand cDNA. Okayama and Berg, *Mol. Cell. Biol.* 2:161-170 (1982), the entirety of which is herein incorporated by reference, report a method for obtaining full length cDNA constructs. This method has been simplified by using synthetic primer-adapters that have both homopolymeric tails for priming the synthesis of the first and second strands and restriction sites for cloning into plasmids (Coleclough *et al.*, *Gene* 34:305-314 (1985), the entirety of which is herein incorporated by reference) and bacteriophage vectors (Krawinkel *et al.*, *Nucleic Acids Res.* 14:1913 (1986), the entirety of which is herein incorporated by reference; Han *et al.*, *Nucleic Acids Res.* 15:6304 (1987), the entirety of which is herein incorporated by reference).

These strategies have been coupled with additional strategies for isolating rare mRNA populations. For example, a typical mammalian cell contains between 10,000 and 30,000 different mRNA sequences (Davidson, *Gene Activity in Early Development*, 2nd ed., Academic Press, New York (1976), the entirety of which is herein incorporated by reference). The number of clones required to achieve a given probability that a low-abundance mRNA will be present in a cDNA library is $N = (\ln(1-P))/(\ln(1-1/n))$ where N is the number of clones required, P is the probability desired and 1/n is the fractional proportion of the total mRNA that is represented by a single rare mRNA (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press (1989), the entirety of which is herein incorporated by reference).

A method to enrich preparations of mRNA for sequences of interest is to fractionate by size. One such method is to fractionate by electrophoresis through an agarose gel (Pennica *et al.*, *Nature* 301:214-221 (1983), the entirety of which is herein incorporated by reference). Another such method employs sucrose gradient centrifugation in the presence of an agent, such as methylmercuric hydroxide, that denatures secondary structure in RNA (Schweinfest *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 79:4997-5000 (1982), the entirety of which is herein incorporated by reference).

A frequently adopted method is to construct equalized or normalized cDNA libraries (Ko, *Nucleic Acids Res.* 18:5705-5711 (1990), the entirety of which is herein incorporated by reference; Patanjali *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:1943-1947 (1991), the entirety of which is herein incorporated by reference). Typically, the cDNA population is normalized by subtractive hybridization (Schmid *et al.*, *J. Neurochem.* 48:307-312 (1987), the entirety of which is herein incorporated by reference; Fargnoli *et al.*, *Anal. Biochem.* 187:364-373 (1990), the

entirety of which is herein incorporated by reference; Travis *et al.*, *Proc. Natl. Acad. Sci (U.S.A.)* 85:1696-1700 (1988), the entirety of which is herein incorporated by reference; Kato, *Eur. J. Neurosci.* 2:704-711 (1990); and Schweinfest *et al.*, *Genet. Anal. Tech. Appl.* 7:64-70 (1990), the entirety of which is herein incorporated by reference). Subtraction represents another method for reducing the population of certain sequences in the cDNA library (Swaroop *et al.*, *Nucleic Acids Res.* 19:1954 (1991), the entirety of which is herein incorporated by reference).

ESTs can be sequenced by a number of methods. Two basic methods may be used for DNA sequencing, the chain termination method of Sanger *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 74:5463-5467 (1977), the entirety of which is herein incorporated by reference and the chemical degradation method of Maxam and Gilbert, *Proc. Nat. Acad. Sci. (U.S.A.)* 74:560-564 (1977), the entirety of which is herein incorporated by reference. Automation and advances in technology such as the replacement of radioisotopes with fluorescence-based sequencing have reduced the effort required to sequence DNA (Craxton, *Methods* 2:20-26 (1991), the entirety of which is herein incorporated by reference; Ju *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 92:4347-4351 (1995), the entirety of which is herein incorporated by reference; Tabor and Richardson, *Proc. Natl. Acad. Sci. (U.S.A.)* 92:6339-6343 (1995), the entirety of which is herein incorporated by reference). Automated sequencers are available from, for example, Pharmacia Biotech, Inc., Piscataway, New Jersey (Pharmacia ALF), LI-COR, Inc., Lincoln, Nebraska (LI-COR 4,000) and Millipore, Bedford, Massachusetts (Millipore BaseStation).

In addition, advances in capillary gel electrophoresis have also reduced the effort required to sequence DNA and such advances provide a rapid high resolution approach for sequencing DNA samples (Swerdlow and Gesteland, *Nucleic Acids Res.* 18:1415-1419 (1990); Smith, *Nature* 349:812-813 (1991); Luckey *et al.*, *Methods Enzymol.* 218:154-172 (1993); Lu *et al.*, *J.*

Chromatog. A. 680:497-501 (1994); Carson *et al.*, *Anal. Chem.* 65:3219-3226 (1993); Huang *et al.*, *Anal. Chem.* 64:2149-2154 (1992); Kheterpal *et al.*, *Electrophoresis* 17:1852-1859 (1996); Quesada and Zhang, *Electrophoresis* 17:1841-1851 (1996); Baba, *Yakugaku Zasshi* 117:265-281 (1997), all of which are herein incorporated by reference in their entirety).

ESTs longer than 150 nucleotides have been found to be useful for similarity searches and mapping (Adams *et al.*, *Science* 252:1651-1656 (1991), herein incorporated by reference). ESTs, which can represent copies of up to the full length transcript, may be partially or completely sequenced. Between 150-450 nucleotides of sequence information is usually generated as this is the length of sequence information that is routinely and reliably produced using single run sequence data. Typically, only single run sequence data is obtained from the cDNA library (Adams *et al.*, *Science* 252:1651-1656 (1991). Automated single run sequencing typically results in an approximately 2-3% error or base ambiguity rate (Boguski *et al.*, *Nature Genetics* 4:332-333 (1993), the entirety of which is herein incorporated by reference).

EST databases have been constructed or partially constructed from, for example, *C. elegans* (McCombie *et al.*, *Nature Genetics* 1:124-131 (1992)), human liver cell line HepG2 (Okubo *et al.*, *Nature Genetics* 2:173-179 (1992)), human brain RNA (Adams *et al.*, *Science* 252:1651-1656 (1991); Adams *et al.*, *Nature* 355:632-635 (1992)), *Arabidopsis*, (Newman *et al.*, *Plant Physiol.* 106:1241-1255 (1994)); and rice (Kurata *et al.*, *Nature Genetics* 8:365-372 (1994)).

III. SEQUENCE COMPARISONS

A characteristic feature of a DNA sequence is that it can be compared with other DNA sequences. Sequence comparisons can be undertaken by determining the similarity of the test or query sequence with sequences in publicly available or proprietary databases ("similarity

analysis”) or by searching for certain motifs (“intrinsic sequence analysis”)(e.g. *cis* elements)(Coulson, *Trends in Biotechnology* 12:76-80 (1994), the entirety of which is herein incorporated by reference); Birren *et al.*, *Genome Analysis 1*: Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 543-559 (1997), the entirety of which is herein incorporated by reference).

Similarity analysis includes database search and alignment. Examples of public databases include the DNA Database of Japan (DDBJ)(<http://www.ddbj.nig.ac.jp/>); Genebank (<http://www.ncbi.nlm.nih.gov/Web/Search/Index.html>); and the European Molecular Biology Laboratory Nucleic Acid Sequence Database (EMBL) (http://www.ebi.ac.uk/ebi_docs/embl_db/embl-db.html). Other appropriate databases include dbEST (<http://www.ncbi.nlm.nih.gov/dbEST/index.html>), SwissProt (http://www.ebi.ac.uk/ebi_docs/swisprot_db/swisshome.html), PIR (<http://www-nbrt.georgetown.edu/pir/>) and The Institute for Genome Research (<http://www.tigr.org/tdb/tdb.html>)

A number of different search algorithms have been developed, one example of which are the suite of programs referred to as BLAST programs. There are five implementations of BLAST, three designed for nucleotide sequences queries (BLASTN, BLASTX and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology* 12:76-80 (1994); Birren *et al.*, *Genome Analysis 1*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 543-559 (1997)).

BLASTN takes a nucleotide sequence (the query sequence) and its reverse complement and searches them against a nucleotide sequence database. BLASTN was designed for speed, not maximum sensitivity and may not find distantly related coding sequences. BLASTX takes a

nucleotide sequence, translates it in three forward reading frames and three reverse complement reading frames and then compares the six translations against a protein sequence database.

BLASTX is useful for sensitive analysis of preliminary (single-pass) sequence data and is tolerant of sequencing errors (Gish and States, *Nature Genetics* 3:266-272 (1993), the entirety of which is herein incorporated by reference). BLASTN and BLASTX may be used in concert for analyzing EST data (Coulson, *Trends in Biotechnology* 12:76-80 (1994); Birren *et al.*, *Genome Analysis* 1:543-559 (1997)).

Given a coding nucleotide sequence and the protein it encodes, it is often preferable to use the protein as the query sequence to search a database because of the greatly increased sensitivity to detect more subtle relationships. This is due to the larger alphabet of proteins (20 amino acids) compared with the alphabet of nucleic acid sequences (4 bases), where it is far easier to obtain a match by chance. In addition, with nucleotide alignments, only a match (positive score) or a mismatch (negative score) is obtained, but with proteins, the presence of conservative amino acid substitutions can be taken into account. Here, a mismatch may yield a positive score if the non-identical residue has physical/chemical properties similar to the one it replaced. Various scoring matrices are used to supply the substitution scores of all possible amino acid pairs. A general purpose scoring system is the BLOSUM62 matrix (Henikoff and Henikoff, *Proteins* 17:49-61 (1993), the entirety of which is herein incorporated by reference), which is currently the default choice for BLAST programs. BLOSUM62 is tailored for alignments of moderately diverged sequences and thus may not yield the best results under all conditions. Altschul, *J. Mol. Biol.* 36:290-300 (1993), the entirety of which is herein incorporated by reference, describes a combination of three matrices to cover all contingencies. This may improve sensitivity, but at the expense of slower searches. In practice, a single

BLOSUM62 matrix is often used but others (PAM40 and PAM250) may be attempted when additional analysis is necessary. Low PAM matrices are directed at detecting very strong but localized sequence similarities, whereas high PAM matrices are directed at detecting long but weak alignments between very distantly related sequences.

Homologues in other organisms are available that can be used for comparative sequence analysis. Multiple alignments are performed to study similarities and differences in a group of related sequences. CLUSTAL W is a multiple sequence alignment package that performs progressive multiple sequence alignments based on the method of Feng and Doolittle, *J. Mol. Evol.* 25:351-360 (1987), the entirety of which is herein incorporated by reference. Each pair of sequences is aligned and the distance between each pair is calculated; from this distance matrix, a guide tree is calculated and all of the sequences are progressively aligned based on this tree. A feature of the program is its sensitivity to the effect of gaps on the alignment; gap penalties are varied to encourage the insertion of gaps in probable loop regions instead of in the middle of structured regions. Users can specify gap penalties, choose between a number of scoring matrices, or supply their own scoring matrix for both pairwise alignments and multiple alignments. CLUSTAL W for UNIX and VMS systems is available at: <ftp.ebi.ac.uk>. Another program is MACAW (Schuler *et al.*, *Proteins Struct. Func. Genet.* 9:180-190 (1991), the entirety of which is herein incorporated by reference, for which both Macintosh and Microsoft Windows versions are available. MACAW uses a graphical interface, provides a choice of several alignment algorithms and is available by anonymous ftp at: <ncbi.nlm.nih.gov> (directory/pub/macaw).

Sequence motifs are derived from multiple alignments and can be used to examine individual sequences or an entire database for subtle patterns. With motifs, it is sometimes

possible to detect distant relationships that may not be demonstrable based on comparisons of primary sequences alone. Currently, the largest collection of sequence motifs in the world is PROSITE (Bairoch and Bucher, *Nucleic Acid Research* 22:3583-3589 (1994), the entirety of which is herein incorporated by reference). PROSITE may be accessed via either the ExPASy server on the World Wide Web or anonymous ftp site. Many commercial sequence analysis packages also provide search programs that use PROSITE data.

A resource for searching protein motifs is the BLOCKS E-mail server developed by Henikoff, *Trends Biochem Sci.* 18:267-268 (1993), the entirety of which is herein incorporated by reference; Henikoff and Henikoff, *Nucleic Acid Research* 19:6565-6572 (1991), the entirety of which is herein incorporated by reference; Henikoff and Henikoff, *Proteins* 17:49-61 (1993). BLOCKS searches a protein or nucleotide sequence against a database of protein motifs or "blocks." Blocks are defined as short, ungapped multiple alignments that represent highly conserved protein patterns. The blocks themselves are derived from entries in PROSITE as well as other sources. Either a protein query or a nucleotide query can be submitted to the BLOCKS server; if a nucleotide sequence is submitted, the sequence is translated in all six reading frames and motifs are sought for these conceptual translations. Once the search is completed, the server will return a ranked list of significant matches, along with an alignment of the query sequence to the matched BLOCKS entries.

Conserved protein domains can be represented by two-dimensional matrices, which measure either the frequency or probability of the occurrences of each amino acid residue and deletions or insertions in each position of the domain. This type of model, when used to search against protein databases, is sensitive and usually yields more accurate results than simple motif searches. Two popular implementations of this approach are profile searches such as GCG

program ProfileSearch and Hidden Markov Models (HMMs)(Krough *et al.*, *J. Mol. Biol.* 235:1501-1531, (1994); Eddy, *Current Opinion in Structural Biology* 6:361-365, (1996), both of which are herein incorporated by reference in their entirety). In both cases, a large number of common protein domains have been converted into profiles, as present in the PROSITE library, or HMM models, as in the Pfam protein domain library (Sonnhammer *et al.*, *Proteins* 28:405-420 (1997), the entirety of which is herein incorporated by reference). Pfam contains more than 500 HMM models for enzymes, transcription factors, signal transduction molecules and structural proteins. Protein databases can be queried with these profiles or HMM models, which will identify proteins containing the domain of interest. For example, HMMSW or HMMFS, two programs in a public domain package called HMMER (Sonnhammer *et al.*, *Proteins* 28:405-420 (1997)) can be used.

PROSITE and BLOCKS represent collected families of protein motifs. Thus, searching these databases entails submitting a single sequence to determine whether or not that sequence is similar to the members of an established family. Programs working in the opposite direction compare a collection of sequences with individual entries in the protein databases. An example of such a program is the Motif Search Tool, or MoST (Tatusov *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:12091-12095 (1994), the entirety of which is herein incorporated by reference). On the basis of an aligned set of input sequences, a weight matrix is calculated by using one of four methods (selected by the user). A weight matrix is simply a representation, position by position of how likely a particular amino acid will appear. The calculated weight matrix is then used to search the databases. To increase sensitivity, newly found sequences are added to the original data set, the weight matrix is recalculated and the search is performed again. This procedure continues until no new sequences are found.

SUMMARY OF THE INVENTION

The present invention provides a substantially purified nucleic acid molecule that encodes a maize or a soybean enzyme or fragment thereof, wherein the maize or the soybean enzyme is selected from the group consisting of: (a) adenine phosphoribosyl transferase (b) β glucosidase and (c) isopentyltransferase.

The present invention also provides a substantially purified nucleic acid molecule that encodes a plant cytokinin pathway enzyme or fragment thereof, wherein the nucleic acid molecule is selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof.

The present invention also provides a substantially purified maize or soybean enzyme or fragment thereof, wherein the maize or soybean enzyme is selected from the group consisting of (a) adenine phosphoribosyl transferase or fragment thereof; (b) β glucosidase or fragment thereof; and (c) isopentyltransferase or fragment thereof.

The present invention also provides a substantially purified maize or soybean cytokinin pathway protein or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 711.

The present invention also provides a substantially purified maize or soybean adenine phosphoribosyl transferase enzyme or fragment thereof encoded by a first nucleic acid molecule

which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515.

The present invention also provides a substantially purified maize or soybean adenine phosphoribosyl transferase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515.

The present invention also provides a substantially purified maize or soybean β glucosidase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710.

The present invention also provides a substantially purified maize or soybean β glucosidase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710.

The present invention also provides a substantially purified soybean isopentyltransferase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence consisting of a complement of SEQ ID NO: 711.

The present invention also provides a substantially purified soybean isopentyltransferase enzyme or fragment thereof encoded by a nucleic acid sequence comprising SEQ ID NO: 711.

The present invention also provides a purified antibody or fragment thereof which is capable of specifically binding to a maize or soybean enzyme or fragment thereof, wherein the maize or soybean enzyme or fragment thereof is encoded by a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515 and a maize or soybean adenine phosphoribosyl transferase enzyme or fragment thereof encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a maize or a soybean β glucosidase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a complement of SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710 and a maize or soybean β glucosidase enzyme or fragment thereof encoded by a nucleic acid sequence selected

from the group consisting of SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710.

The present invention also provides a substantially purified antibody or fragment thereof, the antibody or fragment thereof capable of specifically binding to a soybean isopentyltransferase enzyme or fragment thereof encoded by a first nucleic acid molecule which specifically hybridizes to a second nucleic acid molecule, the second nucleic acid molecule consisting of a complement of a nucleic acid sequence having SEQ ID NO: 711 or a soybean isopentyltransferase enzyme or fragment thereof encoded by a nucleic acid sequence comprising SEQ ID NO: 711.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; (B) a structural nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of (a) a nucleic acid sequence which encodes for adenine phosphoribosyl transferase or fragment thereof; (b) a nucleic acid sequence which encodes for β glucosidase or fragment thereof; and (c) a nucleic acid sequence which encodes for isopentyltransferase or fragment thereof; and (d) a nucleic acid sequence which is complementary to any of the nucleic acid sequences of (a) through (c); and (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; which is linked to (B) a structural nucleic acid molecule,

wherein the structural nucleic acid molecule encodes a plant cytokinin pathway enzyme or fragment thereof, the structural nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; which is linked to (B) a structural nucleic acid molecule, wherein the structural nucleic acid molecule is selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; which is linked to (B) a transcribed nucleic acid molecule with a transcribed strand and a non-transcribed strand, wherein the transcribed strand is complementary to a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof; which is

linked to (C) a 3' non-translated sequence that functions in plant cells to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a transformed plant having a nucleic acid molecule which comprises: (A) an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule; which is linked to: (B) a transcribed nucleic acid molecule with a transcribed strand and a non-transcribed strand, wherein a transcribed mRNA of the transcribed strand is complementary to an endogenous mRNA molecule having a nucleic acid sequence selected from the group consisting of an endogenous mRNA molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, an endogenous mRNA molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and an endogenous mRNA molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof; which is linked to (C) a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of the mRNA molecule.

The present invention also provides a method for determining a level or pattern in a plant cell of an enzyme in a plant metabolic pathway comprising: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule selected from the group of marker nucleic acid molecules which specifically hybridize to a nucleic acid molecule having the nucleic acid sequence of SEQ ID NO: 1 through SEQ ID NO: 711 or compliments thereof, with a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant

0997644-10601

tissue permits the detection of an mRNA for the enzyme; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the enzyme in the plant metabolic pathway.

The present invention also provides a method for determining a level or pattern of a plant cytokinin pathway enzyme in a plant cell or plant tissue comprising: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragment of either, with a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue permits the detection of the plant cytokinin pathway enzyme; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the plant cytokinin pathway enzyme.

The present invention also provides a method for determining a level or pattern of a plant cytokinin pathway enzyme in a plant cell or plant tissue comprising: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or complement thereof or fragment of either, a

09976364-104600

nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or complement thereof or fragment of either and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or complement thereof or fragment of either, with a complementary nucleic acid molecule obtained from the plant cell or plant tissue, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue permits the detection of the plant cytokinin pathway enzyme; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) detecting the level or pattern of the complementary nucleic acid, wherein the detection of the complementary nucleic acid is predictive of the level or pattern of the plant cytokinin pathway enzyme.

The present invention also provides a method for determining a level or pattern of a plant cytokinin pathway enzyme in a plant cell or plant tissue under evaluation which comprises assaying the concentration of a molecule, whose concentration is dependent upon the expression of a gene, the gene specifically hybridizes to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof, in comparison to the concentration of that molecule present in a reference plant cell or a reference plant tissue with a known level or pattern of the plant cytokinin pathway enzyme, wherein the assayed concentration of the molecule is compared to the assayed concentration of the molecule in the reference plant cell or reference plant tissue with the known level or pattern of the plant cytokinin pathway enzyme.

09976034-104504

The present invention also provides a method for determining a level or pattern of a plant cytokinin pathway enzyme in a plant cell or plant tissue under evaluation which comprises assaying the concentration of a molecule, whose concentration is dependent upon the expression of a gene, the gene specifically hybridizes to a nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or complement thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or complement thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or complement thereof, in comparison to the concentration of that molecule present in a reference plant cell or a reference plant tissue with a known level or pattern of the plant cytokinin pathway enzyme, wherein the assayed concentration of the molecule is compared to the assayed concentration of the molecule in the reference plant cell or the reference plant tissue with the known level or pattern of the plant cytokinin pathway enzyme.

The present invention provides a method of determining a mutation in a plant whose presence is predictive of a mutation affecting a level or pattern of a protein comprising the steps: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid, the marker nucleic acid selected from the group of marker nucleic acid molecules which specifically hybridize to a nucleic acid molecule having a nucleic acid sequence selected from the group of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the plant cytokinin pathway enzyme in the plant; (B) permitting hybridization

between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method for determining a mutation in a plant whose presence is predictive of a mutation affecting the level or pattern of a plant cytokinin pathway enzyme comprising the steps: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a nucleic acid molecule that is linked to a gene, the gene specifically hybridizes to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the plant cytokinin pathway enzyme in the plant; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method for determining a mutation in a plant whose presence is predictive of a mutation affecting the level or pattern of a plant cytokinin pathway enzyme comprising the steps: (A) incubating, under conditions permitting nucleic acid hybridization, a marker nucleic acid molecule, the marker nucleic acid molecule comprising a nucleic acid molecule that is linked to a gene, the gene specifically hybridizes to a nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or a

soybean adenine phosphoribosyl transferase enzyme or complement thereof, a nucleic acid molecule that encodes a soybean β glucosidase enzyme or complement thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or complement thereof and a complementary nucleic acid molecule obtained from the plant, wherein nucleic acid hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant permits the detection of a polymorphism whose presence is predictive of a mutation affecting the level or pattern of the plant cytokinin pathway enzyme in the plant; (B) permitting hybridization between the marker nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant; and (C) detecting the presence of the polymorphism, wherein the detection of the polymorphism is predictive of the mutation.

The present invention also provides a method of producing a plant containing an overexpressed protein comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region has a nucleic acid sequence selected from group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the protein; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing an overexpressed plant cytokinin enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter

0997604-101901

region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof; wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the plant cytokinin pathway enzyme; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing an overexpressed plant cytokinin pathway enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, a nucleic acid molecule that encodes a soybean β glucosidase enzyme or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof, wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in overexpression of the plant cytokinin pathway enzyme protein; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing reduced levels of a plant cytokinin pathway enzyme comprising: (A) transforming the plant with a

functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711; wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in co-suppression of the plant cytokinin pathway enzyme protein; and (B) growing the transformed plant.

The present invention also provides a method of producing a plant containing reduced levels of a plant cytokinin pathway enzyme comprising: (A) transforming the plant with a functional nucleic acid molecule, wherein the functional nucleic acid molecule comprises a promoter region, wherein the promoter region is linked to a structural region, wherein the structural region comprises a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof, wherein the structural region is linked to a 3' non-translated sequence that functions in the plant to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and wherein the functional nucleic acid molecule results in co-suppression of the plant cytokinin pathway enzyme; and (B) growing the transformed plant.

0924034-101803

The present invention also provides a method for reducing expression of a plant cytokinin pathway enzyme in a plant comprising: (A) transforming the plant with a nucleic acid molecule, the nucleic acid molecule having an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule, wherein the exogenous promoter region is linked to a transcribed nucleic acid molecule having a transcribed strand and a non-transcribed strand, wherein the transcribed strand is complementary to a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either and the transcribed strand is complementary to an endogenous mRNA molecule; and wherein the transcribed nucleic acid molecule is linked to a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and (B) growing the transformed plant.

The present invention also provides a method for reducing expression of a plant cytokinin pathway enzyme in a plant comprising: (A) transforming the plant with a nucleic acid molecule, the nucleic acid molecule having an exogenous promoter region which functions in a plant cell to cause the production of a mRNA molecule, wherein the exogenous promoter region is linked to a transcribed nucleic acid molecule having a transcribed strand and a non-transcribed strand, wherein a transcribed mRNA of the transcribed strand is complementary to a nucleic acid molecule selected from the group consisting of an endogenous mRNA molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or fragment thereof, an endogenous mRNA molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and an endogenous mRNA molecule that encodes a soybean

isopentyltransferase enzyme or fragment thereof, and wherein the transcribed nucleic acid molecule is linked to a 3' non-translated sequence that functions in the plant cell to cause termination of transcription and addition of polyadenylated ribonucleotides to a 3' end of a mRNA molecule; and (B) growing the transformed plant.

The present invention also provides a method of determining an association between a polymorphism and a plant trait comprising: (A) hybridizing a nucleic acid molecule specific for the polymorphism to genetic material of a plant, wherein the nucleic acid molecule has a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragment of either; and (B) calculating the degree of association between the polymorphism and the plant trait.

The present invention also provides a method of determining an association between a polymorphism and a plant trait comprising: (A) hybridizing a nucleic acid molecule specific for the polymorphism to genetic material of a plant, wherein the nucleic acid molecule is selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme complement thereof or fragment of either and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme complement thereof or fragment of either and (B) calculating the degree of association between the polymorphism and the plant trait.

The present invention also provides a method of isolating a nucleic acid that encodes a plant cytokinin pathway enzyme or fragment thereof comprising: (A) incubating under conditions permitting nucleic acid hybridization, a first nucleic acid molecule comprising a

097404-1034-0000

nucleic acid sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragment of either with a complementary second nucleic acid molecule obtained from a plant cell or plant tissue; (B) permitting hybridization between the first nucleic acid molecule and the second nucleic acid molecule obtained from the plant cell or plant tissue; and (C) isolating the second nucleic acid molecule.

The present invention also provides a method of isolating a nucleic acid molecule that encodes a plant cytokinin pathway enzyme or fragment thereof comprising: (A) incubating under conditions permitting nucleic acid hybridization, a first nucleic acid molecule selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or complement thereof or fragment of either and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme complement thereof or fragment of either, with a complementary second nucleic acid molecule obtained from a plant cell or plant tissue; (B) permitting hybridization between the plant cytokinin pathway nucleic acid molecule and the complementary nucleic acid molecule obtained from the plant cell or plant tissue; and (C) isolating the second nucleic acid molecule.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and Agents of the Present Invention

Definitions:

As used herein, a cytokinin pathway enzyme is any enzyme that is associated with the synthesis or degradation of cytokinin.

As used herein, a cytokinin synthesis enzyme is any enzyme that is associated with the synthesis of cytokinin.

As used herein, a cytokinin degradation enzyme is any enzyme that is associated with the degradation of cytokinin.

As used herein, adenine phosphoribosyl transferase is any enzyme that catalyzes the conversion of iP to [9R-5'P]iP.

As used herein, β glucosidase is any enzyme that catalyzes the hydrolysis of aryl and alkyl β -D-glucosides and/or cellobiose with release of β -D-glucose.

As used herein, isopentyltransferase is any enzyme that catalyzes the first reaction of the pathway in which N6(Δ^2 -isopentenyl) adenosine-5'-monophosphate ("[9R-5'P]iP") is generated from iPP and AMP.

Agents

(a) Nucleic Acid Molecules

Agents of the present invention include plant nucleic acid molecules and more preferably include maize and soybean nucleic acid molecules and more preferably include nucleic acid molecules of the maize genotypes B73 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), B73 x Mo17 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), DK604 (Dekalb Genetics, Dekalb, Illinois U.S.A.), H99 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), RX601 (Asgrow Seed Company, Des Moines, Iowa), Mo17 (Illinois Foundation Seeds, Champaign, Illinois U.S.A.), and soybean types Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa), C1944 (United States Department of Agriculture (USDA) Soybean Germplasm Collection, Urbana, Illinois U.S.A.), Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois

U.S.A.), FT108 (Monsoy, Brazil), Hartwig (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), BW211S Null (Tohoku University, Morioka, Japan), PI507354 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), Asgrow A4922 (Asgrow Seed Company, Des Moines, Iowa U.S.A.), PI227687 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.), PI229358 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and Asgrow A3237 (Asgrow Seed Company, Des Moines, Iowa U.S.A.).

A subset of the nucleic acid molecules of the present invention includes nucleic acid molecules that are marker molecules. Another subset of the nucleic acid molecules of the present invention include nucleic acid molecules that encode a protein or fragment thereof. Another subset of the nucleic acid molecules of the present invention are EST molecules.

Fragment nucleic acid molecules may encode significant portion(s) of, or indeed most of, these nucleic acid molecules. Alternatively, the fragments may comprise smaller oligonucleotides (having from about 15 to about 250 nucleotide residues and more preferably, about 15 to about 30 nucleotide residues).

As used herein, an agent, be it a naturally occurring molecule or otherwise may be "substantially purified," if desired, such that one or more molecules that is or may be present in a naturally occurring preparation containing that molecule will have been removed or will be present at a lower concentration than that at which it would normally be found.

The agents of the present invention will preferably be "biologically active" with respect to either a structural attribute, such as the capacity of a nucleic acid to hybridize to another nucleic acid molecule, or the ability of a protein to be bound by an antibody (or to compete with another molecule for such binding). Alternatively, such an attribute may be catalytic and thus involve the capacity of the agent to mediate a chemical reaction or response.

The agents of the present invention may also be recombinant. As used herein, the term recombinant means any agent (e.g. DNA, peptide etc.), that is, or results, however indirect, from human manipulation of a nucleic acid molecule.

It is understood that the agents of the present invention may be labeled with reagents that facilitate detection of the agent (e.g. fluorescent labels, Prober *et al.*, *Science* 238:336-340 (1987); Albarella *et al.*, EP 144914; chemical labels, Sheldon *et al.*, U.S. Patent 4,582,789; Albarella *et al.*, U.S. Patent 4,563,417; modified bases, Miyoshi *et al.*, EP 119448, all of which are hereby incorporated by reference in their entirety).

It is further understood, that the present invention provides recombinant bacterial, mammalian, microbial, insect, fungal and plant cells and viral constructs comprising the agents of the present invention. (See, for example, Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants; Section (b) Fungal Constructs and Fungal Transformants; Section (c) Mammalian Constructs and Transformed Mammalian Cells; Section (d) Insect Constructs and Transformed Insect Cells; and Section (e) Bacterial Constructs and Transformed Bacterial Cells)

Nucleic acid molecules or fragments thereof of the present invention are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally

complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional "low-stringency" conditions.

Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional "high-stringency" conditions. Conventional stringency conditions are described by Sambrook *et al.*, *Molecular Cloning*, A Laboratory Manual, 2nd Ed., Cold Spring Harbor Press, Cold Spring Harbor, New York (1989) and by Haymes *et al.*, *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, DC (1985), the entirety of which is herein incorporated by reference. Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. Thus, in order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

Appropriate stringency conditions which promote DNA hybridization, for example, 6.0 X sodium chloride/sodium citrate (SSC) at about 45°C, followed by a wash of 2.0 X SSC at 50°C, are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0 X SSC at 50°C to a high stringency of about 0.2 X SSC at 50°C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22°C, to high stringency conditions at about 65°C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed.

In a preferred embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof under moderately stringent conditions, for example at about 2.0 X SSC and about 65°C.

In a particularly preferred embodiment, a nucleic acid of the present invention will include those nucleic acid molecules that specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof under high stringency conditions such as 0.2 X SSC and about 65°C.

In one aspect of the present invention, the nucleic acid molecules of the present invention have one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof. In another aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 90% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof. In a further aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 95% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof. In a more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 98% sequence identity with one or more of the nucleic acid sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof. In an even more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention share between 100% and 99% sequence identity with one or more of the sequences set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof.

093607-400000

In a further more preferred aspect of the present invention, one or more of the nucleic acid molecules of the present invention exhibit 100% sequence identity with a nucleic acid molecule present within MONN01, SATMON001 through SATMON031, SATMON033, SATMON034, SATMON~001, SATMONNN01, SATMONNN04 through SATMONNN006, CMz029 through CMz031, CMz033, CMz035 through CMz037, CMz039 through CMz042, CMz044 through CMz045, CMz047 through CMz050, SOYMON001 through SOYMON038, Soy51 through Soy56, Soy58 through Soy62, Soy65 through Soy66, Soy 68 through Soy73 and Soy76 through Soy77, Lib9, Lib22 through Lib25, Lib35, Lib80 through Lib81, Lib 144, Lib146, Lib147, Lib190, Lib3032 through Lib3036 and Lib3099 (Monsanto Company, St. Louis, Missouri U.S.A.).

(i) Nucleic Acid Molecules Encoding Proteins or Fragments Thereof

Nucleic acid molecules of the present invention can comprise sequences that encode a cytokinin pathway protein or fragment thereof. Such proteins or fragments thereof include homologues of known proteins in other organisms.

In a preferred embodiment of the present invention, a maize or a soybean protein or fragment thereof of the present invention is a homologue of another plant protein. In another preferred embodiment of the present invention, a maize or a soybean protein or fragment thereof of the present invention is a homologue of a fungal protein. In another preferred embodiment of the present invention, a maize or a soybean protein of the present invention is a homologue of mammalian protein. In another preferred embodiment of the present invention, a maize or a soybean protein or fragment thereof of the present invention is a homologue of a bacterial protein. In another preferred embodiment of the present invention, a soybean protein or fragment thereof of the present invention is a homologue of a maize protein. In another preferred

embodiment of the present invention, a maize protein homologue or fragment thereof of the present invention is a homologue of a soybean protein.

In a preferred embodiment of the present invention, the nucleic molecule of the present invention encodes a maize or a soybean protein or fragment thereof where a maize or a soybean protein exhibits a BLAST probability score of greater than $1E-12$, preferably a BLAST probability score of between about $1E-30$ and about $1E-12$, even more preferably a BLAST probability score of greater than $1E-30$ with its homologue.

In another preferred embodiment of the present invention, the nucleic acid molecule encoding a maize or a soybean protein or fragment thereof exhibits a % identity with its homologue of between about 25% and about 40%, more preferably of between about 40 and about 70%, even more preferably of between about 70% and about 90% and even more preferably between about 90% and 99%. In another preferred embodiment, of the present invention, a maize or a soybean protein or fragments thereof exhibits a % identity with its homologue of 100%.

In a preferred embodiment of the present invention, the nucleic molecule of the present invention encodes a maize or a soybean protein or fragment thereof where a maize or a soybean protein exhibits a BLAST score of greater than 120, preferably a BLAST score of between about 1450 and about 120, even more preferably a BLAST score of greater than 1450 with its homologue.

Nucleic acid molecules of the present invention also include non-maize, non-soybean homologues. Preferred non- homologues are selected from the group consisting of alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut, pepper, potato, rice, rye, sorghum, strawberry,

sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm and *Phaseolus*.

In a preferred embodiment, nucleic acid molecules having SEQ ID NO: 1 through SEQ ID NO: 711 or complements and fragments of either can be utilized to obtain such homologues.

The degeneracy of the genetic code, which allows different nucleic acid sequences to code for the same protein or peptide, is known in the literature. (U.S. Patent No. 4,757,006, the entirety of which is herein incorporated by reference).

In an aspect of the present invention, one or more of the nucleic acid molecules of the present invention differ in nucleic acid sequence from those encoding a maize or a soybean protein or fragment thereof in SEQ ID NO: 1 through SEQ ID NO: 711 due to the degeneracy in the genetic code in that they encode the same protein but differ in nucleic acid sequence.

In another further aspect of the present invention, one or more of the nucleic acid molecules of the present invention differ in nucleic acid sequence from those encoding a maize or a soybean protein or fragment thereof in SEQ ID NO: 1 through SEQ ID NO: 711 due to fact that the different nucleic acid sequence encodes a protein having one or more conservative amino acid residue. Examples of conservative substitutions are set forth in Table 1. It is understood that codons capable of coding for such conservative substitutions are known in the art.

Table 1

<u>Original Residue</u>	<u>Conservative Substitutions</u>
Ala	Ser
Arg	Lys
Asn	Gln; His

Asp	Glu
Cys	Ser; Ala
Gln	Asn
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu; Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

In a further aspect of the present invention, one or more of the nucleic acid molecules of the present invention differ in nucleic acid sequence from those encoding a maize or a soybean protein or fragment thereof set forth in SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof due to the fact that one or more codons encoding an amino acid has been substituted for a codon that encodes a nonessential substitution of the amino acid originally encoded.

Agents of the present invention include nucleic acid molecules that encode a maize or a soybean cytokinin pathway protein or fragment thereof and particularly substantially purified nucleic acid molecules selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase protein or fragment thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase protein or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase protein or fragment thereof.

Non-limiting examples of such nucleic acid molecules of the present invention are nucleic acid molecules comprising: SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof that encode for a cytokinin pathway protein or fragment thereof, SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515 or fragment thereof that encode for an adenine phosphoribosyl transferase protein or fragment thereof, SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710 or fragment thereof that encode for a β glucosidase protein or fragment thereof and SEQ ID NO: 711 or fragment thereof that encodes for an isopentyltransferase protein or fragment thereof.

A nucleic acid molecule of the present invention can also encode an homologue of a maize or a soybean adenine phosphoribosyl transferase or fragment thereof, a maize or a soybean β glucosidase or fragment thereof or a soybean isopentyltransferase or fragment thereof. As used herein a homologue protein molecule or fragment thereof is a counterpart protein molecule or fragment thereof in a second species (*e.g.*, maize adenine phosphoribosyl transferase protein is a homologue of *Arabidopsis*' adenine phosphoribosyl transferase protein).

(ii) Nucleic Acid Molecule Markers and Probes

One aspect of the present invention concerns markers that include nucleic acid molecules SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either that can act as markers or other nucleic acid molecules of the present invention that can act as markers.. Genetic markers of the present invention include “dominant” or “codominant” markers “Codominant markers” reveal the presence of two or more alleles (two per diploid individual) at a locus. “Dominant markers” reveal the presence of only a single allele per locus. The presence of the dominant marker phenotype (e.g., a band of DNA) is an indication that one allele is present in either the homozygous or heterozygous condition. The absence of the dominant marker phenotype (e.g. absence of a DNA band) is merely evidence that “some other” undefined allele is present. In the case of populations where individuals are predominantly homozygous and loci are predominately dimorphic, dominant and codominant markers can be equally valuable. As populations become more heterozygous and multi-allelic, codominant markers often become more informative of the genotype than dominant markers. Marker molecules can be, for example, capable of detecting polymorphisms such as single nucleotide polymorphisms (SNPs).

SNPs are single base changes in genomic DNA sequence. They occur at greater frequency and are spaced with a greater uniformity throughout a genome than other reported forms of polymorphism. The greater frequency and uniformity of SNPs means that there is greater probability that such a polymorphism will be found near or in a genetic locus of interest than would be the case for other polymorphisms. SNPs are located in protein-coding regions and noncoding regions of a genome. Some of these SNPs may result in defective or variant protein expression (e.g., as a results of mutations or defective splicing). Analysis (genotyping) of

characterized SNPs can require only a plus/minus assay rather than a lengthy measurement, permitting easier automation.

SNPs can be characterized using any of a variety of methods. Such methods include the direct or indirect sequencing of the site, the use of restriction enzymes (Botstein *et al.*, *Am. J. Hum. Genet.* 32:314-331 (1980), the entirety of which is herein incorporated reference; Konieczny and Ausubel, *Plant J.* 4:403-410 (1993), the entirety of which is herein incorporated by reference), enzymatic and chemical mismatch assays (Myers *et al.*, *Nature* 313:495-498 (1985), the entirety of which is herein incorporated by reference), allele-specific PCR (Newton *et al.*, *Nucl. Acids Res.* 17:2503-2516 (1989), the entirety of which is herein incorporated by reference; Wu *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:2757-2760 (1989), the entirety of which is herein incorporated by reference), ligase chain reaction (Barany, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:189-193 (1991), the entirety of which is herein incorporated by reference), single-strand conformation polymorphism analysis (Labrune *et al.*, *Am. J. Hum. Genet.* 48: 1115-1120 (1991), the entirety of which is herein incorporated by reference), primer-directed nucleotide incorporation assays (Kuppuswami *et al.*, *Proc. Natl. Acad. Sci. USA* 88:1143-1147 (1991), the entirety of which is herein incorporated by reference), dideoxy fingerprinting (Sarkar *et al.*, *Genomics* 13:441-443 (1992), the entirety of which is herein incorporated by reference), solid-phase ELISA-based oligonucleotide ligation assays (Nikiforov *et al.*, *Nucl. Acids Res.* 22:4167-4175 (1994), the entirety of which is herein incorporated by reference), oligonucleotide fluorescence-quenching assays (Livak *et al.*, *PCR Methods Appl.* 4:357-362 (1995), the entirety of which is herein incorporated by reference), 5'-nuclease allele-specific hybridization TaqMan assay (Livak *et al.*, *Nature Genet.* 9:341-342 (1995), the entirety of which is herein incorporated by reference), template-directed dye-terminator incorporation (TDI) assay (Chen and Kwok,

Nucl. Acids Res. 25:347-353 (1997), the entirety of which is herein incorporated by reference), allele-specific molecular beacon assay (Tyagi *et al.*, *Nature Biotech.* 16: 49-53 (1998), the entirety of which is herein incorporated by reference), PinPoint assay (Haff and Smirnov, *Genome Res.* 7: 378-388 (1997), the entirety of which is herein incorporated by reference) and dCAPS analysis (Neff *et al.*, *Plant J.* 14:387-392 (1998), the entirety of which is herein incorporated by reference).

Additional markers, such as AFLP markers, RFLP markers and RAPD markers, can be utilized (Walton, *Seed World* 22-29 (July, 1993), the entirety of which is herein incorporated by reference; Burow and Blake, *Molecular Dissection of Complex Traits*, 13-29, Paterson (ed.), CRC Press, New York (1988), the entirety of which is herein incorporated by reference). DNA markers can be developed from nucleic acid molecules using restriction endonucleases, the PCR and/or DNA sequence information. RFLP markers result from single base changes or insertions/deletions. These codominant markers are highly abundant in plant genomes, have a medium level of polymorphism and are developed by a combination of restriction endonuclease digestion and Southern blotting hybridization. CAPS are similarly developed from restriction nuclease digestion but only of specific PCR products. These markers are also codominant, have a medium level of polymorphism and are highly abundant in the genome. The CAPS result from single base changes and insertions/deletions.

Another marker type, RAPDs, are developed from DNA amplification with random primers and result from single base changes and insertions/deletions in plant genomes. They are dominant markers with a medium level of polymorphisms and are highly abundant. AFLP markers require using the PCR on a subset of restriction fragments from extended adapter

primers. These markers are both dominant and codominant are highly abundant in genomes and exhibit a medium level of polymorphism.

SSRs require DNA sequence information. These codominant markers result from repeat length changes, are highly polymorphic and do not exhibit as high a degree of abundance in the genome as CAPS, AFLPs and RAPDs SNPs also require DNA sequence information. These codominant markers result from single base substitutions. They are highly abundant and exhibit a medium of polymorphism (Rafalski *et al.*, In: *Nonmammalian Genomic Analysis*, Birren and Lai (ed.), Academic Press, San Diego, CA, pp. 75-134 (1996), the entirety of which is herein incorporated by reference). It is understood that a nucleic acid molecule of the present invention may be used as a marker.

A PCR probe is a nucleic acid molecule capable of initiating a polymerase activity while in a double-stranded structure to with another nucleic acid. Various methods for determining the structure of PCR probes and PCR techniques exist in the art. Computer generated searches using programs such as Primer3 (www-genome.wi.mit.edu/cgi-bin/primer/primer3.cgi), STSPipeline (www-genome.wi.mit.edu/cgi-bin/www-STS_Pipeline), or GeneUp (Pesole *et al.*, *BioTechniques* 25:112-123 (1998) the entirety of which is herein incorporated by reference), for example, can be used to identify potential PCR primers.

It is understood that a fragment of one or more of the nucleic acid molecules of the present invention may be a probe and specifically a PCR probe.

(b) Protein and Peptide Molecules

A class of agents comprises one or more of the protein or fragments thereof or peptide molecules encoded by SEQ ID NO: 1 through SEQ ID NO: 711 or one or more of the protein or fragment thereof and peptide molecules encoded by other nucleic acid agents of the present

invention. As used herein, the term "protein molecule" or "peptide molecule" includes any molecule that comprises five or more amino acids. It is well known in the art that proteins may undergo modification, including post-translational modifications, such as, but not limited to, disulfide bond formation, glycosylation, phosphorylation, or oligomerization. Thus, as used herein, the term "protein molecule" or "peptide molecule" includes any protein molecule that is modified by any biological or non-biological process. The terms "amino acid" and "amino acids" refer to all naturally occurring L-amino acids. This definition is meant to include norleucine, ornithine, homocysteine and homoserine.

Non-limiting examples of the protein or fragment thereof of the present invention include a maize or a soybean cytokinin pathway protein or fragment thereof; a maize or a soybean adenine phosphoribosyl transferase or fragment thereof, a maize or β glucosidase or fragment thereof or a soybean isopentyltransferase or fragment thereof.

Non-limiting examples of the protein or fragment molecules of the present invention are an cytokinin pathway protein or fragment thereof encoded by: SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof that encode for a cytokinin pathway protein or fragment thereof, SEQ ID NO: 1 through SEQ ID NO: 40 and SEQ ID NO: 480 through SEQ ID NO: 515 or fragment thereof that encode for an adenine phosphoribosyl transferase protein or fragment thereof, SEQ ID NO: 41 through SEQ ID NO: 479 and SEQ ID NO: 516 through SEQ ID NO: 710 or fragment thereof that encode for a β glucosidase protein or fragment thereof and SEQ ID NO: 711 or fragment thereof that encodes for an isopentyltransferase protein or fragment thereof.

One or more of the protein or fragment of peptide molecules may be produced via chemical synthesis, or more preferably, by expressing in a suitable bacterial or eucaryotic host.

Suitable methods for expression are described by Sambrook *et al.*, (In: *Molecular Cloning, A Laboratory Manual, 2nd Edition, Cold Spring Harbor Press, Cold Spring Harbor, New York* (1989)), or similar texts. For example, the protein may be expressed in, for example, Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants; Section (b) Fungal Constructs and Fungal Transformants; Section (c) Mammalian Constructs and Transformed Mammalian Cells; Section (d) Insect Constructs and Transformed Insect Cells; and Section (e) Bacterial Constructs and Transformed Bacterial Cells.

A “protein fragment” is a peptide or polypeptide molecule whose amino acid sequence comprises a subset of the amino acid sequence of that protein. A protein or fragment thereof that comprises one or more additional peptide regions not derived from that protein is a “fusion” protein. Such molecules may be derivatized to contain carbohydrate or other moieties (such as keyhole limpet hemocyanin, etc.). Fusion protein or peptide molecules of the present invention are preferably produced via recombinant means.

Another class of agents comprise protein or peptide molecules or fragments or fusions thereof encoded by SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof in which conservative, non-essential or non-relevant amino acid residues have been added, replaced or deleted. Computerized means for designing modifications in protein structure are known in the art (Dahiyat and Mayo, *Science* 278:82-87 (1997), the entirety of which is herein incorporated by reference).

The protein molecules of the present invention include plant homologue proteins. An example of such a homologue is a homologue protein of a non-maize or non soybean plant species, that include but not limited to alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut,

pepper, potato, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm, *Phaseolus* etc. Particularly preferred non-maize or non-soybean for use for the isolation of homologs would include, *Arabidopsis*, barley, cotton, oat, oilseed rape, rice, canola, ornamentals, sugarcane, sugarbeet, tomato, potato, wheat and turf grasses. Such a homologue can be obtained by any of a variety of methods. Most preferably, as indicated above, one or more of the disclosed sequences (SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof) will be used to define a pair of primers that may be used to isolate the homologue-encoding nucleic acid molecules from any desired species. Such molecules can be expressed to yield homologues by recombinant means.

(c) Antibodies

One aspect of the present invention concerns antibodies, single-chain antigen binding molecules, or other proteins that specifically bind to one or more of the protein or peptide molecules of the present invention and their homologues, fusions or fragments. Such antibodies may be used to quantitatively or qualitatively detect the protein or peptide molecules of the present invention. As used herein, an antibody or peptide is said to “specifically bind” to a protein or peptide molecule of the present invention if such binding is not competitively inhibited by the presence of non-related molecules.

Nucleic acid molecules that encode all or part of the protein of the present invention can be expressed, via recombinant means, to yield protein or peptides that can in turn be used to elicit antibodies that are capable of binding the expressed protein or peptide. Such antibodies may be used in immunoassays for that protein. Such protein-encoding molecules, or their fragments may be a “fusion” molecule (i.e., a part of a larger nucleic acid molecule) such that, upon expression,

a fusion protein is produced. It is understood that any of the nucleic acid molecules of the present invention may be expressed, via recombinant means, to yield proteins or peptides encoded by these nucleic acid molecules.

The antibodies that specifically bind proteins and protein fragments of the present invention may be polyclonal or monoclonal and may comprise intact immunoglobulins, or antigen binding portions of immunoglobulins fragments (such as $(F(ab'))$, $F(ab')_2$), or single-chain immunoglobulins producible, for example, via recombinant means. It is understood that practitioners are familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of antibodies (*see*, for example, Harlow and Lane, In: *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York (1988), the entirety of which is herein incorporated by reference).

Murine monoclonal antibodies are particularly preferred. BALB/c mice are preferred for this purpose, however, equivalent strains may also be used. The animals are preferably immunized with approximately 25 μ g of purified protein (or fragment thereof) that has been emulsified in a suitable adjuvant (such as TiterMax adjuvant (Vaxcel, Norcross, GA)). Immunization is preferably conducted at two intramuscular sites, one intraperitoneal site and one subcutaneous site at the base of the tail. An additional i.v. injection of approximately 25 μ g of antigen is preferably given in normal saline three weeks later. After approximately 11 days following the second injection, the mice may be bled and the blood screened for the presence of anti-protein or peptide antibodies. Preferably, a direct binding Enzyme-Linked Immunoassay (ELISA) is employed for this purpose.

More preferably, the mouse having the highest antibody titer is given a third i.v. injection of approximately 25 μ g of the same protein or fragment. The splenic leukocytes from this animal may be recovered 3 days later and then permitted to fuse, most preferably, using polyethylene glycol, with cells of a suitable myeloma cell line (such as, for example, the P3X63Ag8.653 myeloma cell line). Hybridoma cells are selected by culturing the cells under "HAT" (hypoxanthine-aminopterin-thymine) selection for about one week. The resulting clones may then be screened for their capacity to produce monoclonal antibodies ("mAbs"), preferably by direct ELISA.

In one embodiment, anti-protein or peptide monoclonal antibodies are isolated using a fusion of a protein or peptide of the present invention, or conjugate of a protein or peptide of the present invention, as immunogens. Thus, for example, a group of mice can be immunized using a fusion protein emulsified in Freund's complete adjuvant (*e.g.* approximately 50 μ g of antigen per immunization). At three week intervals, an identical amount of antigen is emulsified in Freund's incomplete adjuvant and used to immunize the animals. Ten days following the third immunization, serum samples are taken and evaluated for the presence of antibody. If antibody titers are too low, a fourth booster can be employed. Polysera capable of binding the protein or peptide can also be obtained using this method.

In a preferred procedure for obtaining monoclonal antibodies, the spleens of the above-described immunized mice are removed, disrupted and immune splenocytes are isolated over a ficoll gradient. The isolated splenocytes are fused, using polyethylene glycol with BALB/c-derived HGPRT (hypoxanthine guanine phosphoribosyl transferase) deficient P3x63xAg8.653 plasmacytoma cells. The fused cells are plated into 96 well microtiter plates and screened for

hybridoma fusion cells by their capacity to grow in culture medium supplemented with hypoxanthine, aminopterin and thymidine for approximately 2-3 weeks.

Hybridoma cells that arise from such incubation are preferably screened for their capacity to produce an immunoglobulin that binds to a protein of interest. An indirect ELISA may be used for this purpose. In brief, the supernatants of hybridomas are incubated in microtiter wells that contain immobilized protein. After washing, the titer of bound immunoglobulin can be determined using, for example, a goat anti-mouse antibody conjugated to horseradish peroxidase. After additional washing, the amount of immobilized enzyme is determined (for example through the use of a chromogenic substrate). Such screening is performed as quickly as possible after the identification of the hybridoma in order to ensure that a desired clone is not overgrown by non-secreting neighbor cells. Desirably, the fusion plates are screened several times since the rates of hybridoma growth vary. In a preferred embodiment, a different antigenic form may be used to screen the hybridoma. Thus, for example, the splenocytes may be immunized with one immunogen, but the resulting hybridomas can be screened using a different immunogen. It is understood that any of the protein or peptide molecules of the present invention may be used to raise antibodies.

As discussed below, such antibody molecules or their fragments may be used for diagnostic purposes. Where the antibodies are intended for diagnostic purposes, it may be desirable to derivatize them, for example with a ligand group (such as biotin) or a detectable marker group (such as a fluorescent group, a radioisotope or an enzyme).

The ability to produce antibodies that bind the protein or peptide molecules of the present invention permits the identification of mimetic compounds of those molecules. A "mimetic

compound" is a compound that is not that compound, or a fragment of that compound, but which nonetheless exhibits an ability to specifically bind to antibodies directed against that compound.

It is understood that any of the agents of the present invention can be substantially purified and/or be biologically active and/or recombinant.

Uses of the Agents of the Invention

Nucleic acid molecules and fragments thereof of the present invention may be employed to obtain other nucleic acid molecules from the same species (e.g., ESTs or fragment thereof from maize may be utilized to obtain other nucleic acid molecules from maize). Such nucleic acid molecules include the nucleic acid molecules that encode the complete coding sequence of a protein and promoters and flanking sequences of such molecules. In addition, such nucleic acid molecules include nucleic acid molecules that encode for other isozymes or gene family members. Such molecules can be readily obtained by using the above-described nucleic acid molecules or fragments thereof to screen cDNA or genomic libraries obtained from maize or soybean. Methods for forming such libraries are well known in the art.

Nucleic acid molecules and fragments thereof of the present invention may also be employed to obtain nucleic acid homologues. Such homologues include the nucleic acid molecule of other plants or other organisms (e.g., alfalfa, *Arabidopsis*, barley, *Brassica*, broccoli, cabbage, citrus, cotton, garlic, oat, oilseed rape, onion, canola, flax, an ornamental plant, pea, peanut, pepper, potato, rice, rye, sorghum, strawberry, sugarcane, sugarbeet, tomato, wheat, poplar, pine, fir, eucalyptus, apple, lettuce, lentils, grape, banana, tea, turf grasses, sunflower, oil palm, *Phaseolus*, etc.) including the nucleic acid molecules that encode, in whole or in part, protein homologues of other plant species or other organisms, sequences of genetic elements such as promoters and transcriptional regulatory elements. Such molecules can be readily

obtained by using the above-described nucleic acid molecules or fragments thereof to screen cDNA or genomic libraries obtained from such plant species. Methods for forming such libraries are well known in the art. Such homologue molecules may differ in their nucleotide sequences from those found in one or more of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof because complete complementarity is not needed for stable hybridization. The nucleic acid molecules of the present invention therefore also include molecules that, although capable of specifically hybridizing with the nucleic acid molecules may lack "complete complementarity."

Any of a variety of methods may be used to obtain one or more of the above-described nucleic acid molecules (Zamechik *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 83:4143-4146 (1986), the entirety of which is herein incorporated by reference; Goodchild *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:5507-5511 (1988), the entirety of which is herein incorporated by reference; Wickstrom *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:1028-1032 (1988), the entirety of which is herein incorporated by reference; Holt *et al.*, *Molec. Cell. Biol.* 8:963-973 (1988), the entirety of which is herein incorporated by reference; Gerwirtz *et al.*, *Science* 242:1303-1306 (1988), the entirety of which is herein incorporated by reference; Anfossi *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:3379-3383 (1989), the entirety of which is herein incorporated by reference; Becker *et al.*, *EMBO J.* 8:3685-3691 (1989); the entirety of which is herein incorporated by reference). Automated nucleic acid synthesizers may be employed for this purpose. In lieu of such synthesis, the disclosed nucleic acid molecules may be used to define a pair of primers that can be used with the polymerase chain reaction (Mullis *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich *et al.*, European Patent 50,424; European Patent 84,796; European Patent 258,017; European Patent 237,362; Mullis, European Patent 201,184; Mullis *et al.*, U.S. Patent 4,683,202; Erlich, U.S. Patent 4,582,788; and Saiki *et al.*, U.S. Patent 4,683,194,

all of which are herein incorporated by reference in their entirety) to amplify and obtain any desired nucleic acid molecule or fragment.

Promoter sequence(s) and other genetic elements, including but not limited to transcriptional regulatory flanking sequences, associated with one or more of the disclosed nucleic acid sequences can also be obtained using the disclosed nucleic acid sequence provided herein. In one embodiment, such sequences are obtained by incubating EST nucleic acid molecules or preferably fragments thereof with members of genomic libraries (*e.g.* maize and soybean) and recovering clones that hybridize to the EST nucleic acid molecule or fragment thereof. In a second embodiment, methods of "chromosome walking," or inverse PCR may be used to obtain such sequences (Frohman *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:8998-9002 (1988); Ohara *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:5673-5677 (1989); Pang *et al.*, *Biotechniques* 22:1046-1048 (1977); Huang *et al.*, *Methods Mol. Biol.* 69:89-96 (1997); Huang *et al.*, *Method Mol. Biol.* 67:287-294 (1997); Benkel *et al.*, *Genet. Anal.* 13:123-127 (1996); Hartl *et al.*, *Methods Mol. Biol.* 58:293-301 (1996), all of which are herein incorporated by reference in their entirety).

The nucleic acid molecules of the present invention may be used to isolate promoters of cell enhanced, cell specific, tissue enhanced, tissue specific, developmentally or environmentally regulated expression profiles. Isolation and functional analysis of the 5' flanking promoter sequences of these genes from genomic libraries, for example, using genomic screening methods and PCR techniques would result in the isolation of useful promoters and transcriptional regulatory elements. These methods are known to those of skill in the art and have been described (See, for example, Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, (1997), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., the entirety of which is herein

incorporated by reference). Promoters obtained utilizing the nucleic acid molecules of the present invention could also be modified to affect their control characteristics. Examples of such modifications would include but are not limited to enhanced sequences as reported in Uses of the Agents of the Invention, Section (a) Plant Constructs and Plant Transformants. Such genetic elements could be used to enhance gene expression of new and existing traits for crop improvements.

In one sub-aspect, such an analysis is conducted by determining the presence and/or identity of polymorphism(s) by one or more of the nucleic acid molecules of the present invention and more preferably one or more of the EST nucleic acid molecule or fragment thereof which are associated with a phenotype, or a predisposition to that phenotype.

Any of a variety of molecules can be used to identify such polymorphism(s). In one embodiment, one or more of the EST nucleic acid molecules (or a sub-fragment thereof) may be employed as a marker nucleic acid molecule to identify such polymorphism(s). Alternatively, such polymorphisms can be detected through the use of a marker nucleic acid molecule or a marker protein that is genetically linked to (i.e., a polynucleotide that co-segregates with) such polymorphism(s).

In an alternative embodiment, such polymorphisms can be detected through the use of a marker nucleic acid molecule that is physically linked to such polymorphism(s). For this purpose, marker nucleic acid molecules comprising a nucleotide sequence of a polynucleotide located within 1mb of the polymorphism(s) and more preferably within 100kb of the polymorphism(s) and most preferably within 10kb of the polymorphism(s) can be employed.

The genomes of animals and plants naturally undergo spontaneous mutation in the course of their continuing evolution (Gusella, *Ann. Rev. Biochem.* 55:831-854 (1986)). A

“polymorphism” is a variation or difference in the sequence of the gene or its flanking regions that arises in some of the members of a species. The variant sequence and the “original” sequence co-exist in the species’ population. In some instances, such co-existence is in stable or quasi-stable equilibrium.

A polymorphism is thus said to be “allelic,” in that, due to the existence of the polymorphism, some members of a species may have the original sequence (i.e., the original “allele”) whereas other members may have the variant sequence (i.e., the variant “allele”). In the simplest case, only one variant sequence may exist and the polymorphism is thus said to be di-allelic. In other cases, the species’ population may contain multiple alleles and the polymorphism is termed tri-allelic, etc. A single gene may have multiple different unrelated polymorphisms. For example, it may have a di-allelic polymorphism at one site and a multi-allelic polymorphism at another site.

The variation that defines the polymorphism may range from a single nucleotide variation to the insertion or deletion of extended regions within a gene. In some cases, the DNA sequence variations are in regions of the genome that are characterized by short tandem repeats (STRs) that include tandem di- or tri-nucleotide repeated motifs of nucleotides. Polymorphisms characterized by such tandem repeats are referred to as "variable number tandem repeat" ("VNTR") polymorphisms. VNTRs have been used in identity analysis (Weber, U.S. Patent 5,075,217; Armour *et al.*, *FEBS Lett.* 307:113-115 (1992); Jones *et al.*, *Eur. J. Haematol.* 39:144-147 (1987); Horn *et al.*, PCT Patent Application WO91/14003; Jeffreys, European Patent Application 370,719; Jeffreys, U.S. Patent 5,175,082; Jeffreys *et al.*, *Amer. J. Hum. Genet.* 39:11-24 (1986); Jeffreys *et al.*, *Nature* 316:76-79 (1985); Gray *et al.*, *Proc. R. Acad. Soc. Lond.* 243:241-253 (1991); Moore *et al.*, *Genomics* 10:654-660 (1991); Jeffreys *et al.*, *Anim. Genet.*

18:1-15 (1987); Hillel *et al.*, *Anim. Genet.* 20:145-155 (1989); Hillel *et al.*, *Genet.* 124:783-789 (1990), all of which are herein incorporated by reference in their entirety).

The detection of polymorphic sites in a sample of DNA may be facilitated through the use of nucleic acid amplification methods. Such methods specifically increase the concentration of polynucleotides that span the polymorphic site, or include that site and sequences located either distal or proximal to it. Such amplified molecules can be readily detected by gel electrophoresis or other means.

The most preferred method of achieving such amplification employs the polymerase chain reaction ("PCR") (Mullis *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich *et al.*, European Patent Appln. 50,424; European Patent Appln. 84,796; European Patent Application 258,017; European Patent Appln. 237,362; Mullis, European Patent Appln. 201,184; Mullis *et al.*, U.S. Patent No. 4,683,202; Erlich, U.S. Patent No. 4,582,788; and Saiki *et al.*, U.S. Patent No. 4,683,194), using primer pairs that are capable of hybridizing to the proximal sequences that define a polymorphism in its double-stranded form.

In lieu of PCR, alternative methods, such as the "Ligase Chain Reaction" ("LCR") may be used (Barany, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:189-193 (1991), the entirety of which is herein incorporated by reference). LCR uses two pairs of oligonucleotide probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a substrate for a template-dependent ligase. As with PCR, the resulting products thus serve as a template in subsequent cycles and an exponential amplification of the desired sequence is obtained.

LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a polymorphic site. In one embodiment, either oligonucleotide will be designed to include the actual polymorphic site of the polymorphism. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the polymorphic site present on the oligonucleotide. Alternatively, the oligonucleotides may be selected such that they do not include the polymorphic site (see, Segev, PCT Application WO 90/01069, the entirety of which is herein incorporated by reference).

The "Oligonucleotide Ligation Assay" ("OLA") may alternatively be employed (Landegren *et al.*, *Science* 241:1077-1080 (1988), the entirety of which is herein incorporated by reference). The OLA protocol uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target. OLA, like LCR, is particularly suited for the detection of point mutations. Unlike LCR, however, OLA results in "linear" rather than exponential amplification of the target sequence.

Nickerson *et al.*, have described a nucleic acid detection assay that combines attributes of PCR and OLA (Nickerson *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:8923-8927 (1990), the entirety of which is herein incorporated by reference). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA. In addition to requiring multiple and separate, processing steps, one problem associated with such combinations is that they inherit all of the problems associated with PCR and OLA.

Schemes based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide", thereby amplifying the di-oligonucleotide, are also known (Wu *et al.*, *Genomics* 4:560-569 (1989), the entirety of which is

herein incorporated by reference) and may be readily adapted to the purposes of the present invention.

Other known nucleic acid amplification procedures, such as allele-specific oligomers, branched DNA technology, transcription-based amplification systems, or isothermal amplification methods may also be used to amplify and analyze such polymorphisms (Malek *et al.*, U.S. Patent 5,130,238; Davey *et al.*, European Patent Application 329,822; Schuster *et al.*, U.S. Patent 5,169,766; Miller *et al.*, PCT Patent Application WO 89/06700; Kwoh *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:1173-1177 (1989); Gingeras *et al.*, PCT Patent Application WO 88/10315; Walker *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:392-396 (1992), all of which are herein incorporated by reference in their entirety).

The identification of a polymorphism can be determined in a variety of ways. By correlating the presence or absence of it in a plant with the presence or absence of a phenotype, it is possible to predict the phenotype of that plant. If a polymorphism creates or destroys a restriction endonuclease cleavage site, or if it results in the loss or insertion of DNA (e.g., a VNTR polymorphism), it will alter the size or profile of the DNA fragments that are generated by digestion with that restriction endonuclease. As such, individuals that possess a variant sequence can be distinguished from those having the original sequence by restriction fragment analysis. Polymorphisms that can be identified in this manner are termed "restriction fragment length polymorphisms" ("RFLPs"). RFLPs have been widely used in human and plant genetic analyses (Glassberg, UK Patent Application 2135774; Skolnick *et al.*, *Cytogen. Cell Genet.* 32:58-67 (1982); Botstein *et al.*, *Ann. J. Hum. Genet.* 32:314-331 (1980); Fischer *et al.*, (PCT Application WO90/13668); Uhlen, PCT Application WO90/11369).

Polymorphisms can also be identified by Single Strand Conformation Polymorphism (SSCP) analysis. SSCP is a method capable of identifying most sequence variations in a single strand of DNA, typically between 150 and 250 nucleotides in length (Elles, *Methods in Molecular Medicine: Molecular Diagnosis of Genetic Diseases*, Humana Press (1996), the entirety of which is herein incorporated by reference); Orita *et al.*, *Genomics* 5:874-879 (1989), the entirety of which is herein incorporated by reference). Under denaturing conditions a single strand of DNA will adopt a conformation that is uniquely dependent on its sequence conformation. This conformation usually will be different, even if only a single base is changed. Most conformations have been reported to alter the physical configuration or size sufficiently to be detectable by electrophoresis. A number of protocols have been described for SSCP including, but not limited to, Lee *et al.*, *Anal. Biochem.* 205:289-293 (1992), the entirety of which is herein incorporated by reference; Suzuki *et al.*, *Anal. Biochem.* 192:82-84 (1991), the entirety of which is herein incorporated by reference; Lo *et al.*, *Nucleic Acids Research* 20:1005-1009 (1992), the entirety of which is herein incorporated by reference; Sarkar *et al.*, *Genomics* 13:441-443 (1992), the entirety of which is herein incorporated by reference. It is understood that one or more of the nucleic acids of the present invention, may be utilized as markers or probes to detect polymorphisms by SSCP analysis.

Polymorphisms may also be found using a DNA fingerprinting technique called amplified fragment length polymorphism (AFLP), which is based on the selective PCR amplification of restriction fragments from a total digest of genomic DNA to profile that DNA (Vos *et al.*, *Nucleic Acids Res.* 23:4407-4414 (1995), the entirety of which is herein incorporated by reference). This method allows for the specific co-amplification of high numbers of

restriction fragments, which can be visualized by PCR without knowledge of the nucleic acid sequence.

AFLP employs basically three steps. Initially, a sample of genomic DNA is cut with restriction enzymes and oligonucleotide adapters are ligated to the restriction fragments of the DNA. The restriction fragments are then amplified using PCR by using the adapter and restriction sequence as target sites for primer annealing. The selective amplification is achieved by the use of primers that extend into the restriction fragments, amplifying only those fragments in which the primer extensions match the nucleotide flanking the restriction sites. These amplified fragments are then visualized on a denaturing polyacrylamide gel.

AFLP analysis has been performed on *Salix* (Beismann *et al.*, *Mol. Ecol.* 6:989-993 (1997), the entirety of which is herein incorporated by reference), *Acinetobacter* (Janssen *et al.*, *Int. J. Syst. Bacteriol.* 47:1179-1187 (1997), the entirety of which is herein incorporated by reference), *Aeromonas popoffi* (Huys *et al.*, *Int. J. Syst. Bacteriol.* 47:1165-1171 (1997), the entirety of which is herein incorporated by reference), rice (McCouch *et al.*, *Plant Mol. Biol.* 35:89-99 (1997), the entirety of which is herein incorporated by reference; Nandi *et al.*, *Mol. Gen. Genet.* 255:1-8 (1997), the entirety of which is herein incorporated by reference; Cho *et al.*, *Genome* 39:373-378 (1996), the entirety of which is herein incorporated by reference), barley (*Hordeum vulgare*) (Simons *et al.*, *Genomics* 44:61-70 (1997), the entirety of which is herein incorporated by reference; Waugh *et al.*, *Mol. Gen. Genet.* 255:311-321 (1997), the entirety of which is herein incorporated by reference; Qi *et al.*, *Mol. Gen. Genet.* 254:330-336 (1997), the entirety of which is herein incorporated by reference; Becker *et al.*, *Mol. Gen. Genet.* 249:65-73 (1995), the entirety of which is herein incorporated by reference), potato (Van der Voort *et al.*, *Mol. Gen. Genet.* 255:438-447 (1997), the entirety of which is herein incorporated by reference;

Meksem *et al.*, *Mol. Gen. Genet.* 249:74-81 (1995), the entirety of which is herein incorporated by reference), *Phytophthora infestans* (Van der Lee *et al.*, *Fungal Genet. Biol.* 21:278-291 (1997), the entirety of which is herein incorporated by reference), *Bacillus anthracis* (Keim *et al.*, *J. Bacteriol.* 179:818-824 (1997), the entirety of which is herein incorporated by reference), *Astragalus cremnophylax* (Travis *et al.*, *Mol. Ecol.* 5:735-745 (1996), the entirety of which is herein incorporated by reference), *Arabidopsis* (Cnops *et al.*, *Mol. Gen. Genet.* 253:32-41 (1996), the entirety of which is herein incorporated by reference), *Escherichia coli* (Lin *et al.*, *Nucleic Acids Res.* 24:3649-3650 (1996), the entirety of which is herein incorporated by reference), *Aeromonas* (Huys *et al.*, *Int. J. Syst. Bacteriol.* 46:572-580 (1996), the entirety of which is herein incorporated by reference), nematode (Folkertsma *et al.*, *Mol. Plant Microbe Interact.* 9:47-54 (1996), the entirety of which is herein incorporated by reference), tomato (Thomas *et al.*, *Plant J.* 8:785-794 (1995), the entirety of which is herein incorporated by reference) and human (Latorra *et al.*, *PCR Methods Appl.* 3:351-358 (1994), the entirety of which is herein incorporated by reference). AFLP analysis has also been used for fingerprinting mRNA (Money *et al.*, *Nucleic Acids Res.* 24:2616-2617 (1996), the entirety of which is herein incorporated by reference; Bachem *et al.*, *Plant J.* 9:745-753 (1996), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acids of the present invention, may be utilized as markers or probes to detect polymorphisms by AFLP analysis or for fingerprinting RNA.

Polymorphisms may also be found using random amplified polymorphic DNA (RAPD) (Williams *et al.*, *Nucl. Acids Res.* 18:6531-6535 (1990), the entirety of which is herein incorporated by reference) and cleaveable amplified polymorphic sequences (CAPS) (Lyamichev *et al.*, *Science* 260:778-783 (1993), the entirety of which is herein incorporated by reference). It

is understood that one or more of the nucleic acid molecules of the present invention, may be utilized as markers or probes to detect polymorphisms by RAPD or CAPS analysis.

Through genetic mapping, a fine scale linkage map can be developed using DNA markers and, then, a genomic DNA library of large-sized fragments can be screened with molecular markers linked to the desired trait. Molecular markers are advantageous for agronomic traits that are otherwise difficult to tag, such as resistance to pathogens, insects and nematodes, tolerance to abiotic stress, quality parameters and quantitative traits such as high yield potential.

The essential requirements for marker-assisted selection in a plant breeding program are: (1) the marker(s) should co-segregate or be closely linked with the desired trait; (2) an efficient means of screening large populations for the molecular marker(s) should be available; and (3) the screening technique should have high reproducibility across laboratories and preferably be economical to use and be user-friendly.

The genetic linkage of marker molecules can be established by a gene mapping model such as, without limitation, the flanking marker model reported by Lander and Botstein, *Genetics* 121:185-199 (1989) and the interval mapping, based on maximum likelihood methods described by Lander and Botstein, *Genetics* 121:185-199 (1989) and implemented in the software package MAPMAKER/QTL (Lincoln and Lander, *Mapping Genes Controlling Quantitative Traits Using MAPMAKER/QTL*, Whitehead Institute for Biomedical Research, Massachusetts, (1990).

Additional software includes Qgene, Version 2.23 (1996), Department of Plant Breeding and Biometry, 266 Emerson Hall, Cornell University, Ithaca, NY, the manual of which is herein incorporated by reference in its entirety). Use of Qgene software is a particularly preferred approach.

A maximum likelihood estimate (MLE) for the presence of a marker is calculated, together with an MLE assuming no QTL effect, to avoid false positives. A \log_{10} of an odds ratio (LOD) is then calculated as: $\text{LOD} = \log_{10}(\text{MLE for the presence of a QTL} / \text{MLE given no linked QTL})$.

The LOD score essentially indicates how much more likely the data are to have arisen assuming the presence of a QTL than in its absence. The LOD threshold value for avoiding a false positive with a given confidence, say 95%, depends on the number of markers and the length of the genome. Graphs indicating LOD thresholds are set forth in Lander and Botstein, *Genetics* 121:185-199 (1989) the entirety of which is herein incorporated by reference and further described by Arús and Moreno-González, *Plant Breeding*, Hayward *et al.*, (eds.) Chapman & Hall, London, pp. 314-331 (1993), the entirety of which is herein incorporated by reference.

Additional models can be used. Many modifications and alternative approaches to interval mapping have been reported, including the use non-parametric methods (Kruglyak and Lander, *Genetics* 139:1421-1428 (1995), the entirety of which is herein incorporated by reference). Multiple regression methods or models can be also be used, in which the trait is regressed on a large number of markers (Jansen, *Biometrics in Plant Breeding*, van Oijen and Jansen (eds.), Proceedings of the Ninth Meeting of the Eucarpia Section Biometrics in Plant Breeding, The Netherlands, pp. 116-124 (1994); Weber and Wricke, *Advances in Plant Breeding*, Blackwell, Berlin, 16 (1994), both of which is herein incorporated by reference in their entirety). Procedures combining interval mapping with regression analysis, whereby the phenotype is regressed onto a single putative QTL at a given marker interval and at the same time onto a number of markers that serve as 'cofactors,' have been reported by Jansen and Stam, *Genetics*

136:1447-1455 (1994), the entirety of which is herein incorporated by reference and Zeng, *Genetics* 136:1457-1468 (1994) the entirety of which is herein incorporated by reference. Generally, the use of cofactors reduces the bias and sampling error of the estimated QTL positions (Utz and Melchinger, *Biometrics in Plant Breeding*, van Oijen and Jansen (eds.) Proceedings of the Ninth Meeting of the Eucarpia Section Biometrics in Plant Breeding, The Netherlands, pp.195-204 (1994), the entirety of which is herein incorporated by reference, thereby improving the precision and efficiency of QTL mapping (Zeng, *Genetics* 136:1457-1468 (1994)). These models can be extended to multi-environment experiments to analyze genotype-environment interactions (Jansen *et al.*, *Theo. Appl. Genet.* 91:33-37 (1995), the entirety of which is herein incorporated by reference).

Selection of an appropriate mapping populations is important to map construction. The choice of appropriate mapping population depends on the type of marker systems employed (Tanksley *et al.*, *Molecular mapping plant chromosomes. Chromosome structure and function: Impact of new concepts*, Gustafson and Appels (eds.), Plenum Press, New York, pp 157-173 (1988), the entirety of which is herein incorporated by reference). Consideration must be given to the source of parents (adapted vs. exotic) used in the mapping population. Chromosome pairing and recombination rates can be severely disturbed (suppressed) in wide crosses (adapted x exotic) and generally yield greatly reduced linkage distances. Wide crosses will usually provide segregating populations with a relatively large array of polymorphisms when compared to progeny in a narrow cross (adapted x adapted).

An F_2 population is the first generation of selfing after the hybrid seed is produced. Usually a single F_1 plant is selfed to generate a population segregating for all the genes in Mendelian (1:2:1) fashion. Maximum genetic information is obtained from a completely

classified F_2 population using a codominant marker system (Mather, *Measurement of Linkage in Heredity*, Methuen and Co., (1938), the entirety of which is herein incorporated by reference). In the case of dominant markers, progeny tests (e.g. F_3 , BCF_2) are required to identify the heterozygotes, thus making it equivalent to a completely classified F_2 population. However, this procedure is often prohibitive because of the cost and time involved in progeny testing. Progeny testing of F_2 individuals is often used in map construction where phenotypes do not consistently reflect genotype (e.g. disease resistance) or where trait expression is controlled by a QTL. Segregation data from progeny test populations (e.g. F_3 or BCF_2) can be used in map construction. Marker-assisted selection can then be applied to cross progeny based on marker-trait map associations (F_2 , F_3), where linkage groups have not been completely disassociated by recombination events (i.e., maximum disequilibrium).

Recombinant inbred lines (RIL) (genetically related lines; usually $>F_5$, developed from continuously selfing F_2 lines towards homozygosity) can be used as a mapping population. Information obtained from dominant markers can be maximized by using RIL because all loci are homozygous or nearly so. Under conditions of tight linkage (i.e., about $<10\%$ recombination), dominant and co-dominant markers evaluated in RIL populations provide more information per individual than either marker type in backcross populations (Reiter *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:1477-1481 (1992), the entirety of which is herein incorporated by reference). However, as the distance between markers becomes larger (i.e., loci become more independent), the information in RIL populations decreases dramatically when compared to codominant markers.

Backcross populations (e.g., generated from a cross between a successful variety (recurrent parent) and another variety (donor parent) carrying a trait not present in the former)

can be utilized as a mapping population. A series of backcrosses to the recurrent parent can be made to recover most of its desirable traits. Thus a population is created consisting of individuals nearly like the recurrent parent but each individual carries varying amounts or mosaic of genomic regions from the donor parent. Backcross populations can be useful for mapping dominant markers if all loci in the recurrent parent are homozygous and the donor and recurrent parent have contrasting polymorphic marker alleles (Reiter *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 89:1477-1481 (1992)). Information obtained from backcross populations using either codominant or dominant markers is less than that obtained from F₂ populations because one, rather than two, recombinant gametes are sampled per plant. Backcross populations, however, are more informative (at low marker saturation) when compared to RILs as the distance between linked loci increases in RIL populations (i.e. about 15% recombination). Increased recombination can be beneficial for resolution of tight linkages, but may be undesirable in the construction of maps with low marker saturation.

Near-isogenic lines (NIL) created by many backcrosses to produce an array of individuals that are nearly identical in genetic composition except for the trait or genomic region under interrogation can be used as a mapping population. In mapping with NILs, only a portion of the polymorphic loci are expected to map to a selected region.

Bulk segregant analysis (BSA) is a method developed for the rapid identification of linkage between markers and traits of interest (Michelmore *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:9828-9832 (1991), the entirety of which is herein incorporated by reference). In BSA, two bulked DNA samples are drawn from a segregating population originating from a single cross. These bulks contain individuals that are identical for a particular trait (resistant or susceptible to particular disease) or genomic region but arbitrary at unlinked regions (i.e. heterozygous).

Regions unlinked to the target region will not differ between the bulked samples of many individuals in BSA.

It is understood that one or more of the nucleic acid molecules of the present invention may be used as molecular markers. It is also understood that one or more of the protein molecules of the present invention may be used as molecular markers.

In accordance with this aspect of the present invention, a sample nucleic acid is obtained from plants cells or tissues. Any source of nucleic acid may be used. Preferably, the nucleic acid is genomic DNA. The nucleic acid is subjected to restriction endonuclease digestion. For example, one or more nucleic acid molecule or fragment thereof of the present invention can be used as a probe in accordance with the above-described polymorphic methods. The polymorphism obtained in this approach can then be cloned to identify the mutation at the coding region which alters the protein's structure or regulatory region of the gene which affects its expression level.

In an aspect of the present invention, one or more of the nucleic molecules of the present invention are used to determine the level (i.e., the concentration of mRNA in a sample, etc.) in a plant (preferably maize or soybean) or pattern (i.e., the kinetics of expression, rate of decomposition, stability profile, etc.) of the expression of a protein encoded in part or whole by one or more of the nucleic acid molecule of the present invention (collectively, the "Expression Response" of a cell or tissue). As used herein, the Expression Response manifested by a cell or tissue is said to be "altered" if it differs from the Expression Response of cells or tissues of plants not exhibiting the phenotype. To determine whether a Expression Response is altered, the Expression Response manifested by the cell or tissue of the plant exhibiting the phenotype is compared with that of a similar cell or tissue sample of a plant not exhibiting the phenotype. As

will be appreciated, it is not necessary to re-determine the Expression Response of the cell or tissue sample of plants not exhibiting the phenotype each time such a comparison is made; rather, the Expression Response of a particular plant may be compared with previously obtained values of normal plants. As used herein, the phenotype of the organism is any of one or more characteristics of an organism (e.g. disease resistance, pest tolerance, environmental tolerance such as tolerance to abiotic stress, male sterility, quality improvement or yield etc.). A change in genotype or phenotype may be transient or permanent. Also as used herein, a tissue sample is any sample that comprises more than one cell. In a preferred aspect, a tissue sample comprises cells that share a common characteristic (e.g. derived from root, seed, flower, leaf, stem or pollen etc.).

In one aspect of the present invention, an evaluation can be conducted to determine whether a particular mRNA molecule is present. One or more of the nucleic acid molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention are utilized to detect the presence or quantity of the mRNA species. Such molecules are then incubated with cell or tissue extracts of a plant under conditions sufficient to permit nucleic acid hybridization. The detection of double-stranded probe-mRNA hybrid molecules is indicative of the presence of the mRNA; the amount of such hybrid formed is proportional to the amount of mRNA. Thus, such probes may be used to ascertain the level and extent of the mRNA production in a plant's cells or tissues. Such nucleic acid hybridization may be conducted under quantitative conditions (thereby providing a numerical value of the amount of the mRNA present). Alternatively, the assay may be conducted as a qualitative assay that indicates either that the mRNA is present, or that its level exceeds a user set, predefined value.

A principle of *in situ* hybridization is that a labeled, single-stranded nucleic acid probe will hybridize to a complementary strand of cellular DNA or RNA and, under the appropriate conditions, these molecules will form a stable hybrid. When nucleic acid hybridization is combined with histological techniques, specific DNA or RNA sequences can be identified within a single cell. An advantage of *in situ* hybridization over more conventional techniques for the detection of nucleic acids is that it allows an investigator to determine the precise spatial population (Angerer *et al.*, *Dev. Biol.* 101:477-484 (1984), the entirety of which is herein incorporated by reference; Angerer *et al.*, *Dev. Biol.* 112:157-166 (1985), the entirety of which is herein incorporated by reference; Dixon *et al.*, *EMBO J.* 10:1317-1324 (1991), the entirety of which is herein incorporated by reference). *In situ* hybridization may be used to measure the steady-state level of RNA accumulation. It is a sensitive technique and RNA sequences present in as few as 5-10 copies per cell can be detected (Hardin *et al.*, *J. Mol. Biol.* 202:417-431 (1989), the entirety of which is herein incorporated by reference). A number of protocols have been devised for *in situ* hybridization, each with tissue preparation, hybridization and washing conditions (Meyerowitz, *Plant Mol. Biol. Rep.* 5:242-250 (1987), the entirety of which is herein incorporated by reference; Cox and Goldberg, In: *Plant Molecular Biology: A Practical Approach*, Shaw (ed.), pp 1-35, IRL Press, Oxford (1988), the entirety of which is herein incorporated by reference; Raikhel *et al.*, *In situ RNA hybridization in plant tissues*, In: *Plant Molecular Biology Manual*, vol. B9:1-32, Kluwer Academic Publisher, Dordrecht, Belgium (1989), the entirety of which is herein incorporated by reference).

In situ hybridization also allows for the localization of proteins within a tissue or cell (Wilkinson, *In Situ Hybridization*, Oxford University Press, Oxford (1992), the entirety of which is herein incorporated by reference; Langdale, *In Situ Hybridization* In: *The Maize Handbook*,

Freeling and Walbot (eds.), pp 165-179, Springer-Verlag, New York (1994), the entirety of which is herein incorporated by reference). It is understood that one or more of the molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention or one or more of the antibodies of the present invention may be utilized to detect the level or pattern of a cytokinin pathway protein or mRNA thereof by *in situ* hybridization.

Fluorescent *in situ* hybridization allows the localization of a particular DNA sequence along a chromosome which is useful, among other uses, for gene mapping, following chromosomes in hybrid lines or detecting chromosomes with translocations, transversions or deletions. *In situ* hybridization has been used to identify chromosomes in several plant species (Griffor *et al.*, *Plant Mol. Biol.* 17:101-109 (1991), the entirety of which is herein incorporated by reference; Gustafson *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:1899-1902 (1990), herein incorporated by reference; Mukai and Gill, *Genome* 34:448-452 (1991), the entirety of which is herein incorporated by reference; Schwarzacher and Heslop-Harrison, *Genome* 34:317-323 (1991); Wang *et al.*, *Jpn. J. Genet.* 66:313-316 (1991), the entirety of which is herein incorporated by reference; Parra and Windle, *Nature Genetics* 5:17-21 (1993), the entirety of which is herein incorporated by reference). It is understood that the nucleic acid molecules of the present invention may be used as probes or markers to localize sequences along a chromosome.

Another method to localize the expression of a molecule is tissue printing. Tissue printing provides a way to screen, at the same time on the same membrane many tissue sections from different plants or different developmental stages. Tissue-printing procedures utilize films designed to immobilize proteins and nucleic acids. In essence, a freshly cut section of a tissue is pressed gently onto nitrocellulose paper, nylon membrane or polyvinylidene difluoride

membrane. Such membranes are commercially available (e.g. Millipore, Bedford, Massachusetts U.S.A.). The contents of the cut cell transfer onto the membrane and the contents are immobilized to the membrane. The immobilized contents form a latent print that can be visualized with appropriate probes. When a plant tissue print is made on nitrocellulose paper, the cell walls leave a physical print that makes the anatomy visible without further treatment (Varner and Taylor, *Plant Physiol.* 91:31-33 (1989), the entirety of which is herein incorporated by reference).

Tissue printing on substrate films is described by Daoust, *Exp. Cell Res.* 12:203-211 (1957), the entirety of which is herein incorporated by reference, who detected amylase, protease, ribonuclease and deoxyribonuclease in animal tissues using starch, gelatin and agar films. These techniques can be applied to plant tissues (Yomo and Taylor, *Planta* 112:35-43 (1973); the entirety of which is herein incorporated by reference; Harris and Chrispeels, *Plant Physiol.* 56:292-299 (1975), the entirety of which is herein incorporated by reference). Advances in membrane technology have increased the range of applications of Daoust's tissue-printing techniques allowing (Cassab and Varner, *J. Cell. Biol.* 105:2581-2588 (1987), the entirety of which is herein incorporated by reference) the histochemical localization of various plant enzymes and deoxyribonuclease on nitrocellulose paper and nylon (Spruce *et al.*, *Phytochemistry* 26:2901-2903 (1987), the entirety of which is herein incorporated by reference; Barres *et al.*, *Neuron* 5:527-544 (1990), the entirety of which is herein incorporated by reference; Reid and Pont-Lezica, *Tissue Printing: Tools for the Study of Anatomy, Histochemistry and Gene Expression*, Academic Press, New York, New York (1992), the entirety of which is herein incorporated by reference; Reid *et al.*, *Plant Physiol.* 93:160-165 (1990), the entirety of which is

herein incorporated by reference; Ye *et al.*, *Plant J.* 1:175-183 (1991), the entirety of which is herein incorporated by reference).

It is understood that one or more of the molecules of the present invention, preferably one or more of the EST nucleic acid molecules or fragments thereof of the present invention or one or more of the antibodies of the present invention may be utilized to detect the presence or quantity of a cytokinin pathway protein by tissue printing.

Further it is also understood that any of the nucleic acid molecules of the present invention may be used as marker nucleic acids and or probes in connection with methods that require probes or marker nucleic acids. As used herein, a probe is an agent that is utilized to determine an attribute or feature (e.g. presence or absence, location, correlation, etc.) of a molecule, cell, tissue or plant. As used herein, a marker nucleic acid is a nucleic acid molecule that is utilized to determine an attribute or feature (e.g., presence or absence, location, correlation, etc.) of a molecule, cell, tissue or plant.

A microarray-based method for high-throughput monitoring of plant gene expression may be utilized to measure gene-specific hybridization targets. This 'chip'-based approach involves using microarrays of nucleic acid molecules as gene-specific hybridization targets to quantitatively measure expression of the corresponding plant genes (Schena *et al.*, *Science* 270:467-470 (1995), the entirety of which is herein incorporated by reference; Shalon, Ph.D. Thesis, Stanford University (1996), the entirety of which is herein incorporated by reference). Every nucleotide in a large sequence can be queried at the same time. Hybridization can be used to efficiently analyze nucleotide sequences.

Several microarray methods have been described. One method compares the sequences to be analyzed by hybridization to a set of oligonucleotides representing all possible

subsequences (Bains and Smith, *J. Theor. Biol.* 135:303-307 (1989), the entirety of which is herein incorporated by reference). A second method hybridizes the sample to an array of oligonucleotide or cDNA molecules. An array consisting of oligonucleotides complementary to subsequences of a target sequence can be used to determine the identity of a target sequence, measure its amount and detect differences between the target and a reference sequence. Nucleic acid molecules microarrays may also be screened with protein molecules or fragments thereof to determine nucleic acid molecules that specifically bind protein molecules or fragments thereof.

The microarray approach may be used with polypeptide targets (U.S. Patent No. 5,445,934; U.S. Patent No: 5,143,854; U.S. Patent No. 5,079,600; U.S. Patent No. 4,923,901, all of which are herein incorporated by reference in their entirety). Essentially, polypeptides are synthesized on a substrate (microarray) and these polypeptides can be screened with either protein molecules or fragments thereof or nucleic acid molecules in order to screen for either protein molecules or fragments thereof or nucleic acid molecules that specifically bind the target polypeptides. (Fodor *et al.*, *Science* 251:767-773 (1991), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acid molecules or protein or fragments thereof of the present invention may be utilized in a microarray based method.

In a preferred embodiment of the present invention microarrays may be prepared that comprise nucleic acid molecules where such nucleic acid molecules encode at least one, preferably at least two, more preferably at least three cytokinin pathway enzymes. In a preferred embodiment the nucleic acid molecules are selected from the group consisting of a nucleic acid molecule that encodes a maize or a soybean adenine phosphoribosyl transferase enzyme or

fragment thereof, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or fragment thereof and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or fragment thereof.

Site directed mutagenesis may be utilized to modify nucleic acid sequences, particularly as it is a technique that allows one or more of the amino acids encoded by a nucleic acid molecule to be altered (e.g. a threonine to be replaced by a methionine). Three basic methods for site directed mutagenesis are often employed. These are cassette mutagenesis (Wells *et al.*, *Gene* 34:315-323 (1985), the entirety of which is herein incorporated by reference), primer extension (Gilliam *et al.*, *Gene* 12:129-137 (1980), the entirety of which is herein incorporated by reference; Zoller and Smith, *Methods Enzymol.* 100:468-500 (1983), the entirety of which is herein incorporated by reference; Dalbadie-McFarland *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 79:6409-6413 (1982), the entirety of which is herein incorporated by reference) and methods based upon PCR (Scharf *et al.*, *Science* 233:1076-1078 (1986), the entirety of which is herein incorporated by reference; Higuchi *et al.*, *Nucleic Acids Res.* 16:7351-7367 (1988), the entirety of which is herein incorporated by reference). Site directed mutagenesis approaches are also described in European Patent 0 385 962, the entirety of which is herein incorporated by reference; European Patent 0 359 472, the entirety of which is herein incorporated by reference; and PCT Patent Application WO 93/07278, the entirety of which is herein incorporated by reference.

Site directed mutagenesis strategies have been applied to plants for both *in vitro* as well as *in vivo* site directed mutagenesis (Lanz *et al.*, *J. Biol. Chem.* 266:9971-9976 (1991), the entirety of which is herein incorporated by reference; Kovgan and Zhdanov, *Biotekhnologiya*

5:148-154, No. 207160n, Chemical Abstracts 110:225 (1989), the entirety of which is herein incorporated by reference; Ge *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:4037-4041 (1989), the entirety of which is herein incorporated by reference; Zhu *et al.*, *J. Biol. Chem.* 271:18494-18498 (1996), the entirety of which is herein incorporated by reference; Chu *et al.*, *Biochemistry* 33:6150-6157 (1994), the entirety of which is herein incorporated by reference; Small *et al.*, *EMBO J.* 11:1291-1296 (1992), the entirety of which is herein incorporated by reference; Cho *et al.*, *Mol. Biotechnol.* 8:13-16 (1997), the entirety of which is herein incorporated by reference; Kita *et al.*, *J. Biol. Chem.* 271:26529-26535 (1996), the entirety of which is herein incorporated by reference; Jin *et al.*, *Mol. Microbiol.* 7:555-562 (1993), the entirety of which is herein incorporated by reference; Hatfield and Vierstra, *J. Biol. Chem.* 267:14799-14803 (1992), the entirety of which is herein incorporated by reference; Zhao *et al.*, *Biochemistry* 31:5093-5099 (1992), the entirety of which is herein incorporated by reference).

Any of the nucleic acid molecules of the present invention may either be modified by site directed mutagenesis or used as, for example, nucleic acid molecules that are used to target other nucleic acid molecules for modification. It is understood that mutants with more than one altered nucleotide can be constructed using techniques that practitioners are familiar with such as isolating restriction fragments and ligating such fragments into an expression vector (*see, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989)).

Sequence-specific DNA-binding proteins play a role in the regulation of transcription. The isolation of recombinant cDNAs encoding these proteins facilitates the biochemical analysis of their structural and functional properties. Genes encoding such DNA-binding proteins have been isolated using classical genetics (Vollbrecht *et al.*, *Nature* 350: 241-243 (1991), the entirety

of which is herein incorporated by reference) and molecular biochemical approaches, including the screening of recombinant cDNA libraries with antibodies (Landschulz *et al.*, *Genes Dev.* 2:786-800 (1988), the entirety of which is herein incorporated by reference) or DNA probes (Bodner *et al.*, *Cell* 55:505-518 (1988), the entirety of which is herein incorporated by reference). In addition, an *in situ* screening procedure has been used and has facilitated the isolation of sequence-specific DNA-binding proteins from various plant species (Gilmartin *et al.*, *Plant Cell* 4:839-849 (1992), the entirety of which is herein incorporated by reference; Schindler *et al.*, *EMBO J.* 11:1261-1273 (1992), the entirety of which is herein incorporated by reference). An *in situ* screening protocol does not require the purification of the protein of interest (Vinson *et al.*, *Genes Dev.* 2:801-806 (1988), the entirety of which is herein incorporated by reference; Singh *et al.*, *Cell* 52:415-423 (1988), the entirety of which is herein incorporated by reference).

Two steps may be employed to characterize DNA-protein interactions. The first is to identify promoter fragments that interact with DNA-binding proteins, to titrate binding activity, to determine the specificity of binding and to determine whether a given DNA-binding activity can interact with related DNA sequences (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989)). Electrophoretic mobility-shift assay is a widely used assay. The assay provides a rapid and sensitive method for detecting DNA-binding proteins based on the observation that the mobility of a DNA fragment through a nondenaturing, low-ionic strength polyacrylamide gel is retarded upon association with a DNA-binding protein (Fried and Crother, *Nucleic Acids Res.* 9:6505-6525 (1981), the entirety of which is herein incorporated by reference). When one or more specific binding activities have been identified, the exact sequence of the DNA bound by

the protein may be determined. Several procedures for characterizing protein/DNA-binding sites are used, including methylation and ethylation interference assays (Maxam and Gilbert, *Methods Enzymol.* 65:499-560 (1980), the entirety of which is herein incorporated by reference; Wissman and Hillen, *Methods Enzymol.* 208:365-379 (1991), the entirety of which is herein incorporated by reference), footprinting techniques employing DNase I (Galas and Schmitz, *Nucleic Acids Res.* 5:3157-3170 (1978), the entirety of which is herein incorporated by reference), 1,10-phenanthroline-copper ion methods (Sigman *et al.*, *Methods Enzymol.* 208:414-433 (1991), the entirety of which is herein incorporated by reference) and hydroxyl radicals methods (Dixon *et al.*, *Methods Enzymol.* 208:414-433 (1991), the entirety of which is herein incorporated by reference). It is understood that one or more of the nucleic acid molecules of the present invention may be utilized to identify a protein or fragment thereof that specifically binds to a nucleic acid molecule of the present invention. It is also understood that one or more of the protein molecules or fragments thereof of the present invention may be utilized to identify a nucleic acid molecule that specifically binds to it.

A two-hybrid system is based on the fact that many cellular functions are carried out by proteins, such as transcription factors, that interact (physically) with one another. Two-hybrid systems have been used to probe the function of new proteins (Chien *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 88:9578-9582 (1991) the entirety of which is herein incorporated by reference; Durfee *et al.*, *Genes Dev.* 7:555-569 (1993) the entirety of which is herein incorporated by reference; Choi *et al.*, *Cell* 78:499-512 (1994), the entirety of which is herein incorporated by reference; Kranz *et al.*, *Genes Dev.* 8:313-327 (1994), the entirety of which is herein incorporated by reference).

Interaction mating techniques have facilitated a number of two-hybrid studies of protein-protein interaction. Interaction mating has been used to examine interactions between small sets

of tens of proteins (Finley and Brent, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:12098-12984 (1994), the entirety of which is herein incorporated by reference), larger sets of hundreds of proteins (Bendixen *et al.*, *Nucl. Acids Res.* 22:1778-1779 (1994), the entirety of which is herein incorporated by reference) and to comprehensively map proteins encoded by a small genome (Bartel *et al.*, *Nature Genetics* 12:72-77 (1996), the entirety of which is herein incorporated by reference). This technique utilizes proteins fused to the DNA-binding domain and proteins fused to the activation domain. They are expressed in two different haploid yeast strains of opposite mating type and the strains are mated to determine if the two proteins interact. Mating occurs when haploid yeast strains come into contact and result in the fusion of the two haploids into a diploid yeast strain. An interaction can be determined by the activation of a two-hybrid reporter gene in the diploid strain. An advantage of this technique is that it reduces the number of yeast transformations needed to test individual interactions. It is understood that the protein-protein interactions of protein or fragments thereof of the present invention may be investigated using the two-hybrid system and that any of the nucleic acid molecules of the present invention that encode such proteins or fragments thereof may be used to transform yeast in the two-hybrid system.

(a) Plant Constructs and Plant Transformants

One or more of the nucleic acid molecules of the present invention may be used in plant transformation or transfection. Exogenous genetic material may be transferred into a plant cell and the plant cell regenerated into a whole, fertile or sterile plant. Exogenous genetic material is any genetic material, whether naturally occurring or otherwise, from any source that is capable of being inserted into any organism. Such genetic material may be transferred into either monocotyledons and dicotyledons including, but not limited to maize (pp 63-69), soybean (pp

50-60), *Arabidopsis* (p 45), phaseolus (pp 47-49), peanut (pp 49-50), alfalfa (p 60), wheat (pp 69-71), rice (pp 72-79), oat (pp 80-81), sorghum (p 83), rye (p 84), tritordeum (p 84), millet (p85), fescue (p 85), perennial ryegrass (p 86), sugarcane (p87), cranberry (p101), papaya (pp 101-102), banana (p 103), banana (p 103), muskmelon (p 104), apple (p 104), cucumber (p 105), dendrobium (p 109), gladiolus (p 110), chrysanthemum (p 110), liliacea (p 111), cotton (pp113-114), eucalyptus (p 115), sunflower (p 118), canola (p 118), turfgrass (p121), sugarbeet (p 122), coffee (p 122) and dioscorea (p 122), (Christou, In: *Particle Bombardment for Genetic Engineering of Plants*, Biotechnology Intelligence Unit. Academic Press, San Diego, California (1996), the entirety of which is herein incorporated by reference).

Transfer of a nucleic acid that encodes for a protein can result in overexpression of that protein in a transformed cell or transgenic plant. One or more of the proteins or fragments thereof encoded by nucleic acid molecules of the present invention may be overexpressed in a transformed cell or transformed plant. Particularly, any of the cytokinin pathway proteins or fragments thereof may be overexpressed in a transformed cell or transgenic plant. Such overexpression may be the result of transient or stable transfer of the exogenous genetic material.

Exogenous genetic material may be transferred into a plant cell and the plant cell by the use of a DNA vector or construct designed for such a purpose. Design of such a vector is generally within the skill of the art (See, *Plant Molecular Biology: A Laboratory Manual*, Clark (ed.), Springer, New York (1997), the entirety of which is herein incorporated by reference).

A construct or vector may include a plant promoter to express the protein or protein fragment of choice. A number of promoters which are active in plant cells have been described in the literature. These include the nopaline synthase (NOS) promoter (Ebert *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:5745-5749 (1987), the entirety of which is herein incorporated by

reference), the octopine synthase (OCS) promoter (which are carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter (Lawton *et al.*, *Plant Mol. Biol.* 9:315-324 (1987), the entirety of which is herein incorporated by reference) and the CAMV 35S promoter (Odell *et al.*, *Nature* 313:810-812 (1985), the entirety of which is herein incorporated by reference), the figwort mosaic virus 35S-promoter, the light-inducible promoter from the small subunit of ribulose-1,5-bis-phosphate carboxylase (ssRUBISCO), the Adh promoter (Walker *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 84:6624-6628 (1987), the entirety of which is herein incorporated by reference), the sucrose synthase promoter (Yang *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:4144-4148 (1990), the entirety of which is herein incorporated by reference), the R gene complex promoter (Chandler *et al.*, *The Plant Cell* 1:1175-1183 (1989), the entirety of which is herein incorporated by reference) and the chlorophyll a/b binding protein gene promoter, etc. These promoters have been used to create DNA constructs which have been expressed in plants; *see, e.g.*, PCT publication WO 84/02913, herein incorporated by reference in its entirety.

Promoters which are known or are found to cause transcription of DNA in plant cells can be used in the present invention. Such promoters may be obtained from a variety of sources such as plants and plant viruses. It is preferred that the particular promoter selected should be capable of causing sufficient expression to result in the production of an effective amount of the cytokinin pathway protein to cause the desired phenotype. In addition to promoters that are known to cause transcription of DNA in plant cells, other promoters may be identified for use in the current invention by screening a plant cDNA library for genes which are selectively or preferably expressed in the target tissues or cells.

For the purpose of expression in source tissues of the plant, such as the leaf, seed, root or stem, it is preferred that the promoters utilized in the present invention have relatively high expression in these specific tissues. For this purpose, one may choose from a number of promoters for genes with tissue- or cell-specific or -enhanced expression. Examples of such promoters reported in the literature include the chloroplast glutamine synthetase GS2 promoter from pea (Edwards *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:3459-3463 (1990), herein incorporated by reference in its entirety), the chloroplast fructose-1,6-biphosphatase (FBPase) promoter from wheat (Lloyd *et al.*, *Mol. Gen. Genet.* 225:209-216 (1991), herein incorporated by reference in its entirety), the nuclear photosynthetic ST-LS1 promoter from potato (Stockhaus *et al.*, *EMBO J.* 8:2445-2451 (1989), herein incorporated by reference in its entirety), the serine/threonine kinase (PAL) promoter and the glucoamylase (CHS) promoter from *Arabidopsis thaliana*. Also reported to be active in photosynthetically active tissues are the ribulose-1,5-bisphosphate carboxylase (RbcS) promoter from eastern larch (*Larix laricina*), the promoter for the *cab* gene, *cab6*, from pine (Yamamoto *et al.*, *Plant Cell Physiol.* 35:773-778 (1994), herein incorporated by reference in its entirety), the promoter for the *Cab-1* gene from wheat (Fejes *et al.*, *Plant Mol. Biol.* 15:921-932 (1990), herein incorporated by reference in its entirety), the promoter for the *CAB-1* gene from spinach (Lubberstedt *et al.*, *Plant Physiol.* 104:997-1006 (1994), herein incorporated by reference in its entirety), the promoter for the *cab1R* gene from rice (Luan *et al.*, *Plant Cell.* 4:971-981 (1992), the entirety of which is herein incorporated by reference), the pyruvate, orthophosphate dikinase (PPDK) promoter from maize (Matsuoka *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 90: 9586-9590 (1993), herein incorporated by reference in its entirety), the promoter for the tobacco *Lhcb1*2* gene (Cerdan *et al.*, *Plant Mol. Biol.* 33:245-255 (1997), herein incorporated by reference in its entirety), the *Arabidopsis thaliana* SUC2 sucrose-

H⁺ symporter promoter (Truernit *et al.*, *Planta*. 196:564-570 (1995), herein incorporated by reference in its entirety) and the promoter for the thylakoid membrane proteins from spinach (psaD, psaF, psaE, PC, FNR, atpC, atpD, cab, rbcS). Other promoters for the chlorophyll a/b-binding proteins may also be utilized in the present invention, such as the promoters for LhcB gene and PsbP gene from white mustard (*Sinapis alba*; Kretsch *et al.*, *Plant Mol. Biol.* 28:219-229 (1995), the entirety of which is herein incorporated by reference).

For the purpose of expression in sink tissues of the plant, such as the tuber of the potato plant, the fruit of tomato, or the seed of maize, wheat, rice and barley, it is preferred that the promoters utilized in the present invention have relatively high expression in these specific tissues. A number of promoters for genes with tuber-specific or -enhanced expression are known, including the class I patatin promoter (Bevan *et al.*, *EMBO J.* 8:1899-1906 (1986); Jefferson *et al.*, *Plant Mol. Biol.* 14:995-1006 (1990), both of which are herein incorporated by reference in its entirety), the promoter for the potato tuber ADPGPP genes, both the large and small subunits, the sucrose synthase promoter (Salanoubat and Belliard, *Gene*. 60:47-56 (1987), Salanoubat and Belliard, *Gene*. 84:181-185 (1989), both of which are incorporated by reference in their entirety), the promoter for the major tuber proteins including the 22 kd protein complexes and proteinase inhibitors (Hannapel, *Plant Physiol.* 101:703-704 (1993), herein incorporated by reference in its entirety), the promoter for the granule bound starch synthase gene (GBSS) (Visser *et al.*, *Plant Mol. Biol.* 17:691-699 (1991), herein incorporated by reference in its entirety) and other class I and II patatins promoters (Koster-Topfer *et al.*, *Mol Gen Genet.* 219:390-396 (1989); Mignery *et al.*, *Gene*. 62:27-44 (1988), both of which are herein incorporated by reference in their entirety).

Other promoters can also be used to express a cytokinin pathway protein or fragment thereof in specific tissues, such as seeds or fruits. The promoter for β -conglycinin (Chen *et al.*, *Dev. Genet.* 10: 112-122 (1989), herein incorporated by reference in its entirety) or other seed-specific promoters such as the napin and phaseolin promoters, can be used. The zeins are a group of storage proteins found in maize endosperm. Genomic clones for zein genes have been isolated (Pedersen *et al.*, *Cell* 29:1015-1026 (1982), herein incorporated by reference in its entirety) and the promoters from these clones, including the 15 kD, 16 kD, 19 kD, 22 kD, 27 kD and genes, could also be used. Other promoters known to function, for example, in maize include the promoters for the following genes: *waxy*, *Brittle*, *Shrunken 2*, Branching enzymes I and II, starch synthases, debranching enzymes, oleosins, glutelins and sucrose synthases. A particularly preferred promoter for maize endosperm expression is the promoter for the glutelin gene from rice, more particularly the Osgt-1 promoter (Zheng *et al.*, *Mol. Cell Biol.* 13:5829-5842 (1993), herein incorporated by reference in its entirety). Examples of promoters suitable for expression in wheat include those promoters for the ADPGlucose pyrosynthase (ADPGPP) subunits, the granule bound and other starch synthase, the branching and debranching enzymes, the embryogenesis-abundant proteins, the gliadins and the glutenins. Examples of such promoters in rice include those promoters for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases and the glutelins. A particularly preferred promoter is the promoter for rice glutelin, Osgt-1. Examples of such promoters for barley include those for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases, the hordeins, the embryo globulins and the aleurone specific proteins.

Root specific promoters may also be used. An example of such a promoter is the promoter for the acid chitinase gene (Samac *et al.*, *Plant Mol. Biol.* 25:587-596 (1994), the entirety of which is herein incorporated by reference). Expression in root tissue could also be accomplished by utilizing the root specific subdomains of the CaMV35S promoter that have been identified (Lam *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:7890-7894 (1989), herein incorporated by reference in its entirety). Other root cell specific promoters include those reported by Conkling *et al.* (Conkling *et al.*, *Plant Physiol.* 93:1203-1211 (1990), the entirety of which is herein incorporated by reference).

Additional promoters that may be utilized are described, for example, in U.S. Patent Nos. 5,378,619; 5,391,725; 5,428,147; 5,447,858; 5,608,144; 5,608,144; 5,614,399; 5,633,441; 5,633,435; and 4,633,436, all of which are herein incorporated in their entirety. In addition, a tissue specific enhancer may be used (Fromm *et al.*, *The Plant Cell* 1:977-984 (1989), the entirety of which is herein incorporated by reference).

Constructs or vectors may also include with the coding region of interest a nucleic acid sequence that acts, in whole or in part, to terminate transcription of that region. For example, such sequences have been isolated including the Tr7 3' sequence and the NOS 3' sequence (Ingelbrecht *et al.*, *The Plant Cell* 1:671-680 (1989), the entirety of which is herein incorporated by reference; Bevan *et al.*, *Nucleic Acids Res.* 11:369-385 (1983), the entirety of which is herein incorporated by reference), or the like.

A vector or construct may also include regulatory elements. Examples of such include the Adh intron 1 (Callis *et al.*, *Genes and Develop.* 1:1183-1200 (1987), the entirety of which is herein incorporated by reference), the sucrose synthase intron (Vasil *et al.*, *Plant Physiol.* 91:1575-1579 (1989), the entirety of which is herein incorporated by reference) and the TMV

omega element (Gallie *et al.*, *The Plant Cell* 1:301-311 (1989), the entirety of which is herein incorporated by reference). These and other regulatory elements may be included when appropriate.

A vector or construct may also include a selectable marker. Selectable markers may also be used to select for plants or plant cells that contain the exogenous genetic material. Examples of such include, but are not limited to, a neo gene (Potrykus *et al.*, *Mol. Gen. Genet.* 199:183-188 (1985), the entirety of which is herein incorporated by reference) which codes for kanamycin resistance and can be selected for using kanamycin, G418, etc.; a bar gene which codes for bialaphos resistance; a mutant EPSP synthase gene (Hinchey *et al.*, *Bio/Technology* 6:915-922 (1988), the entirety of which is herein incorporated by reference) which encodes glyphosate resistance; a nitrilase gene which confers resistance to bromoxynil (Stalker *et al.*, *J. Biol. Chem.* 263:6310-6314 (1988), the entirety of which is herein incorporated by reference); a mutant acetolactate synthase gene (ALS) which confers imidazolinone or sulphonylurea resistance (European Patent Application 154,204 (Sept. 11, 1985), the entirety of which is herein incorporated by reference); and a methotrexate resistant DHFR gene (Thillet *et al.*, *J. Biol. Chem.* 263:12500-12508 (1988), the entirety of which is herein incorporated by reference).

A vector or construct may also include a transit peptide. Incorporation of a suitable chloroplast transit peptide may also be employed (European Patent Application Publication Number 0218571, the entirety of which is herein incorporated by reference). Translational enhancers may also be incorporated as part of the vector DNA. DNA constructs could contain one or more 5' non-translated leader sequences which may serve to enhance expression of the gene products from the resulting mRNA transcripts. Such sequences may be derived from the promoter selected to express the gene or can be specifically modified to increase translation of

the mRNA. Such regions may also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence. For a review of optimizing expression of transgenes, see Koziel *et al.*, *Plant Mol. Biol.* 32:393-405 (1996), the entirety of which is herein incorporated by reference.

A vector or construct may also include a screenable marker. Screenable markers may be used to monitor expression. Exemplary screenable markers include a β -glucuronidase or uidA gene (GUS) which encodes an enzyme for which various chromogenic substrates are known (Jefferson, *Plant Mol. Biol. Rep.* 5:387-405 (1987), the entirety of which is herein incorporated by reference; Jefferson *et al.*, *EMBO J.* 6:3901-3907 (1987), the entirety of which is herein incorporated by reference); an R-locus gene, which encodes a product that regulates the production of anthocyanin pigments (red color) in plant tissues (Dellaporta *et al.*, *Stadler Symposium* 11:263-282 (1988), the entirety of which is herein incorporated by reference); a β -lactamase gene (Sutcliffe *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 75:3737-3741 (1978), the entirety of which is herein incorporated by reference), a gene which encodes an enzyme for which various chromogenic substrates are known (e.g., PADAC, a chromogenic cephalosporin); a luciferase gene (Ow *et al.*, *Science* 234:856-859 (1986), the entirety of which is herein incorporated by reference); a xylE gene (Zukowsky *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 80:1101-1105 (1983), the entirety of which is herein incorporated by reference) which encodes a catechol dioxygenase that can convert chromogenic catechols; an α -amylase gene (Ikatu *et al.*, *Bio/Technol.* 8:241-242 (1990), the entirety of which is herein incorporated by reference); a tyrosinase gene (Katz *et al.*, *J. Gen. Microbiol.* 129:2703-2714 (1983), the entirety of which is herein incorporated by reference) which encodes an enzyme capable of oxidizing tyrosine to

DOPA and dopaquinone which in turn condenses to melanin; an α -galactosidase, which will turn a chromogenic α -galactose substrate.

Included within the terms "selectable or screenable marker genes" are also genes which encode a secretable marker whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers which encode a secretable antigen that can be identified by antibody interaction, or even secretable enzymes which can be detected catalytically. Secretable proteins fall into a number of classes, including small, diffusible proteins which are detectable, (*e.g.*, by ELISA), small active enzymes which are detectable in extracellular solution (*e.g.*, α -amylase, β -lactamase, phosphinothricin transferase), or proteins which are inserted or trapped in the cell wall (such as proteins which include a leader sequence such as that found in the expression unit of extension or tobacco PR-S). Other possible selectable and/or screenable marker genes will be apparent to those of skill in the art.

There are many methods for introducing transforming nucleic acid molecules into plant cells. Suitable methods are believed to include virtually any method by which nucleic acid molecules may be introduced into a cell, such as by *Agrobacterium* infection or direct delivery of nucleic acid molecules such as, for example, by PEG-mediated transformation, by electroporation or by acceleration of DNA coated particles, etc (Potrykus, *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42:205-225 (1991), the entirety of which is herein incorporated by reference; Vasil, *Plant Mol. Biol.* 25:925-937 (1994), the entirety of which is herein incorporated by reference). For example, electroporation has been used to transform maize protoplasts (Fromm *et al.*, *Nature* 312:791-793 (1986), the entirety of which is herein incorporated by reference).

Other vector systems suitable for introducing transforming DNA into a host plant cell include but are not limited to binary artificial chromosome (BIBAC) vectors (Hamilton *et al.*, *Gene* 200:107-116 (1997), the entirety of which is herein incorporated by reference); and transfection with RNA viral vectors (Della-Cioppa *et al.*, *Ann. N.Y. Acad. Sci.* (1996), 792 (Engineering Plants for Commercial Products and Applications), 57-61, the entirety of which is herein incorporated by reference). Additional vector systems also include plant selectable YAC vectors such as those described in Mullen *et al.*, *Molecular Breeding* 4:449-457 (1988), the entirety of which is herein incorporated by reference).

Technology for introduction of DNA into cells is well known to those of skill in the art. Four general methods for delivering a gene into cells have been described: (1) chemical methods (Graham and van der Eb, *Virology* 54:536-539 (1973), the entirety of which is herein incorporated by reference); (2) physical methods such as microinjection (Capecchi, *Cell* 22:479-488 (1980), the entirety of which is herein incorporated by reference), electroporation (Wong and Neumann, *Biochem. Biophys. Res. Commun.* 107:584-587 (1982); Fromm *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 82:5824-5828 (1985); U.S. Patent No. 5,384,253, all of which are herein incorporated in their entirety); and the gene gun (Johnston and Tang, *Methods Cell Biol.* 43:353-365 (1994), the entirety of which is herein incorporated by reference); (3) viral vectors (Clapp, *Clin. Perinatol.* 20:155-168 (1993); Lu *et al.*, *J. Exp. Med.* 178:2089-2096 (1993); Eglitis and Anderson, *Biotechniques* 6:608-614 (1988), all of which are herein incorporated in their entirety); and (4) receptor-mediated mechanisms (Curiel *et al.*, *Hum. Gen. Ther.* 3:147-154 (1992), Wagner *et al.*, *Proc. Natl. Acad. Sci. (USA)* 89:6099-6103 (1992), both of which are incorporated by reference in their entirety).

Acceleration methods that may be used include, for example, microprojectile bombardment and the like. One example of a method for delivering transforming nucleic acid molecules to plant cells is microprojectile bombardment. This method has been reviewed by Yang and Christou (eds.), *Particle Bombardment Technology for Gene Transfer*, Oxford Press, Oxford, England (1994), the entirety of which is herein incorporated by reference). Non-biological particles (microprojectiles) that may be coated with nucleic acids and delivered into cells by a propelling force. Exemplary particles include those comprised of tungsten, gold, platinum and the like.

A particular advantage of microprojectile bombardment, in addition to it being an effective means of reproducibly transforming monocots, is that neither the isolation of protoplasts (Cristou *et al.*, *Plant Physiol.* 87:671-674 (1988), the entirety of which is herein incorporated by reference) nor the susceptibility of *Agrobacterium* infection are required. An illustrative embodiment of a method for delivering DNA into maize cells by acceleration is a biolistics α -particle delivery system, which can be used to propel particles coated with DNA through a screen, such as a stainless steel or Nytex screen, onto a filter surface covered with corn cells cultured in suspension. Gordon-Kamm *et al.*, describes the basic procedure for coating tungsten particles with DNA (Gordon-Kamm *et al.*, *Plant Cell* 2:603-618 (1990), the entirety of which is herein incorporated by reference). The screen disperses the tungsten nucleic acid particles so that they are not delivered to the recipient cells in large aggregates. A particle delivery system suitable for use with the present invention is the helium acceleration PDS-1000/He gun is available from Bio-Rad Laboratories (Bio-Rad, Hercules, California)(Sanford *et al.*, *Technique* 3:3-16 (1991), the entirety of which is herein incorporated by reference).

For the bombardment, cells in suspension may be concentrated on filters. Filters containing the cells to be bombarded are positioned at an appropriate distance below the microprojectile stopping plate. If desired, one or more screens are also positioned between the gun and the cells to be bombarded.

Alternatively, immature embryos or other target cells may be arranged on solid culture medium. The cells to be bombarded are positioned at an appropriate distance below the microprojectile stopping plate. If desired, one or more screens are also positioned between the acceleration device and the cells to be bombarded. Through the use of techniques set forth herein one may obtain up to 1000 or more foci of cells transiently expressing a marker gene. The number of cells in a focus which express the exogenous gene product 48 hours post-bombardment often range from one to ten and average one to three.

In bombardment transformation, one may optimize the pre-bombardment culturing conditions and the bombardment parameters to yield the maximum numbers of stable transformants. Both the physical and biological parameters for bombardment are important in this technology. Physical factors are those that involve manipulating the DNA/microprojectile precipitate or those that affect the flight and velocity of either the macro- or microprojectiles. Biological factors include all steps involved in manipulation of cells before and immediately after bombardment, the osmotic adjustment of target cells to help alleviate the trauma associated with bombardment and also the nature of the transforming DNA, such as linearized DNA or intact supercoiled plasmids. It is believed that pre-bombardment manipulations are especially important for successful transformation of immature embryos.

In another alternative embodiment, plastids can be stably transformed. Methods disclosed for plastid transformation in higher plants include the particle gun delivery of DNA

containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination (Svab *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 87:8526-8530 (1990); Svab and Maliga, *Proc. Natl. Acad. Sci. (U.S.A.)* 90:913-917 (1993); Staub and Maliga, *EMBO J.* 12:601-606 (1993); U.S. Patents 5,451,513 and 5,545,818, all of which are herein incorporated by reference in their entirety).

Accordingly, it is contemplated that one may wish to adjust various aspects of the bombardment parameters in small scale studies to fully optimize the conditions. One may particularly wish to adjust physical parameters such as gap distance, flight distance, tissue distance and helium pressure. One may also minimize the trauma reduction factors by modifying conditions which influence the physiological state of the recipient cells and which may therefore influence transformation and integration efficiencies. For example, the osmotic state, tissue hydration and the subculture stage or cell cycle of the recipient cells may be adjusted for optimum transformation. The execution of other routine adjustments will be known to those of skill in the art in light of the present disclosure.

Agrobacterium-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated plant integrating vectors to introduce DNA into plant cells is well known in the art. See, for example the methods described by Fraley *et al.*, *Bio/Technology* 3:629-635 (1985) and Rogers *et al.*, *Methods Enzymol.* 153:253-277 (1987), both of which are herein incorporated by reference in their entirety. Further, the integration of the Ti-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is defined by the border sequences and

intervening DNA is usually inserted into the plant genome as described (Spielmann *et al.*, *Mol. Gen. Genet.* 205:34 (1986), the entirety of which is herein incorporated by reference).

Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations as described (Klee *et al.*, In: *Plant DNA Infectious Agents*, Hohn and Schell (eds.), Springer-Verlag, New York, pp. 179-203 (1985), the entirety of which is herein incorporated by reference. Moreover, technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. The vectors described have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes (Rogers *et al.*, *Methods Enzymol.* 153:253-277 (1987)). In addition, *Agrobacterium* containing both armed and disarmed Ti genes can be used for the transformations. In those plant strains where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer.

A transgenic plant formed using *Agrobacterium* transformation methods typically contains a single gene on one chromosome. Such transgenic plants can be referred to as being heterozygous for the added gene. More preferred is a transgenic plant that is homozygous for the added structural gene; *i.e.*, a transgenic plant that contains two added genes, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) an independent segregant transgenic plant that contains a single added gene, germinating some of the seed produced and analyzing the resulting plants produced for the gene of interest.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes that encode a polypeptide of interest. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation.

Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation and combinations of these treatments (*See, for example, Potrykus et al., Mol. Gen. Genet.* 205:193-200 (1986); Lorz *et al., Mol. Gen. Genet.* 199:178 (1985); Fromm *et al., Nature* 319:791 (1986); Uchimiya *et al., Mol. Gen. Genet.* 204:204 (1986); Marcotte *et al., Nature* 335:454-457 (1988), all of which are herein incorporated by reference in their entirety).

Application of these systems to different plant strains depends upon the ability to regenerate that particular plant strain from protoplasts. Illustrative methods for the regeneration of cereals from protoplasts are described (Fujimura *et al., Plant Tissue Culture Letters* 2:74 (1985); Toriyama *et al., Theor Appl. Genet.* 205:34 (1986); Yamada *et al., Plant Cell Rep.* 4:85 (1986); Abdullah *et al., Biotechnolog* 4:1087 (1986), all of which are herein incorporated by reference in their entirety).

To transform plant strains that cannot be successfully regenerated from protoplasts, other ways to introduce DNA into intact cells or tissues can be utilized. For example, regeneration of cereals from immature embryos or explants can be effected as described (Vasil, *Biotechnology* 6:397 (1988), the entirety of which is herein incorporated by reference). In addition, "particle gun" or high-velocity microprojectile technology can be utilized (Vasil *et al., Bio/Technology* 10:667 (1992), the entirety of which is herein incorporated by reference).

Using the latter technology, DNA is carried through the cell wall and into the cytoplasm on the surface of small metal particles as described (Klein *et al.*, *Nature* 328:70 (1987); Klein *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 85:8502-8505 (1988); McCabe *et al.*, *Bio/Technology* 6:923 (1988), all of which are herein incorporated by reference in their entirety). The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants.

Other methods of cell transformation can also be used and include but are not limited to introduction of DNA into plants by direct DNA transfer into pollen (Zhou *et al.*, *Methods Enzymol.* 101:433 (1983); Hess *et al.*, *Intern Rev. Cytol.* 107:367 (1987); Luo *et al.*, *Plant Mol Biol. Reporter* 6:165 (1988), all of which are herein incorporated by reference in their entirety), by direct injection of DNA into reproductive organs of a plant (Pena *et al.*, *Nature* 325:274 (1987), the entirety of which is herein incorporated by reference), or by direct injection of DNA into the cells of immature embryos followed by the rehydration of desiccated embryos (Neuhaus *et al.*, *Theor. Appl. Genet.* 75:30 (1987), the entirety of which is herein incorporated by reference).

The regeneration, development and cultivation of plants from single plant protoplast transformants or from various transformed explants is well known in the art (Weissbach and Weissbach, In: *Methods for Plant Molecular Biology*, Academic Press, San Diego, CA, (1988), the entirety of which is herein incorporated by reference). This regeneration and growth process typically includes the steps of selection of transformed cells, culturing those individualized cells through the usual stages of embryonic development through the rooted plantlet stage. Transgenic embryos and seeds are similarly regenerated. The resulting transgenic rooted shoots are thereafter planted in an appropriate plant growth medium such as soil.

The development or regeneration of plants containing the foreign, exogenous gene that encodes a protein of interest is well known in the art. Preferably, the regenerated plants are self-pollinated to provide homozygous transgenic plants. Otherwise, pollen obtained from the regenerated plants is crossed to seed-grown plants of agronomically important lines. Conversely, pollen from plants of these important lines is used to pollinate regenerated plants. A transgenic plant of the present invention containing a desired polypeptide is cultivated using methods well known to one skilled in the art.

There are a variety of methods for the regeneration of plants from plant tissue. The particular method of regeneration will depend on the starting plant tissue and the particular plant species to be regenerated.

Methods for transforming dicots, primarily by use of *Agrobacterium tumefaciens* and obtaining transgenic plants have been published for cotton (U.S. Patent No. 5,004,863; U.S. Patent No. 5,159,135; U.S. Patent No. 5,518,908, all of which are herein incorporated by reference in their entirety); soybean (U.S. Patent No. 5,569,834; U.S. Patent No. 5,416,011; McCabe *et al.*, *Biotechnology* 6:923 (1988); Christou *et al.*, *Plant Physiol.* 87:671-674 (1988); all of which are herein incorporated by reference in their entirety); *Brassica* (U.S. Patent No. 5,463,174, the entirety of which is herein incorporated by reference); peanut (Cheng *et al.*, *Plant Cell Rep.* 15:653-657 (1996), McKently *et al.*, *Plant Cell Rep.* 14:699-703 (1995), all of which are herein incorporated by reference in their entirety); papaya; and pea (Grant *et al.*, *Plant Cell Rep.* 15:254-258 (1995), the entirety of which is herein incorporated by reference).

Transformation of monocotyledons using electroporation, particle bombardment and *Agrobacterium* have also been reported. Transformation and plant regeneration have been achieved in asparagus (Bytebier *et al.*, *Proc. Natl. Acad. Sci. (USA)* 84:5354 (1987), the entirety

of which is herein incorporated by reference); barley (Wan and Lemaux, *Plant Physiol* 104:37 (1994), the entirety of which is herein incorporated by reference); maize (Rhodes *et al.*, *Science* 240:204 (1988); Gordon-Kamm *et al.*, *Plant Cell* 2:603-618 (1990); Fromm *et al.*, *Bio/Technology* 8:833 (1990); Koziel *et al.*, *Bio/Technology* 11:194 (1993); Armstrong *et al.*, *Crop Science* 35:550-557 (1995); all of which are herein incorporated by reference in their entirety); oat (Somers *et al.*, *Bio/Technology* 10:1589 (1992), the entirety of which is herein incorporated by reference); orchard grass (Horn *et al.*, *Plant Cell Rep.* 7:469 (1988), the entirety of which is herein incorporated by reference); rice (Toriyama *et al.*, *Theor Appl. Genet.* 205:34 (1986); Part *et al.*, *Plant Mol. Biol.* 32:1135-1148 (1996); Abedinia *et al.*, *Aust. J. Plant Physiol.* 24:133-141 (1997); Zhang and Wu, *Theor. Appl. Genet.* 76:835 (1988); Zhang *et al.*, *Plant Cell Rep.* 7:379 (1988); Battraw and Hall, *Plant Sci.* 86:191-202 (1992); Christou *et al.*, *Bio/Technology* 9:957 (1991), all of which are herein incorporated by reference in their entirety); rye (De la Pena *et al.*, *Nature* 325:274 (1987), the entirety of which is herein incorporated by reference); sugarcane (Bower and Birch, *Plant J.* 2:409 (1992), the entirety of which is herein incorporated by reference); tall fescue (Wang *et al.*, *Bio/Technology* 10:691 (1992), the entirety of which is herein incorporated by reference) and wheat (Vasil *et al.*, *Bio/Technology* 10:667 (1992), the entirety of which is herein incorporated by reference; U.S. Patent No. 5,631,152, the entirety of which is herein incorporated by reference.)

Assays for gene expression based on the transient expression of cloned nucleic acid constructs have been developed by introducing the nucleic acid molecules into plant cells by polyethylene glycol treatment, electroporation, or particle bombardment (Marcotte *et al.*, *Nature* 335:454-457 (1988), the entirety of which is herein incorporated by reference; Marcotte *et al.*, *Plant Cell* 1:523-532 (1989), the entirety of which is herein incorporated by reference; McCarty

et al., *Cell* 66:895-905 (1991), the entirety of which is herein incorporated by reference; Hattori *et al.*, *Genes Dev.* 6:609-618 (1992), the entirety of which is herein incorporated by reference; Goff *et al.*, *EMBO J.* 9:2517-2522 (1990), the entirety of which is herein incorporated by reference). Transient expression systems may be used to functionally dissect gene constructs (see generally, Mailga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995)).

Any of the nucleic acid molecules of the present invention may be introduced into a plant cell in a permanent or transient manner in combination with other genetic elements such as vectors, promoters, enhancers etc. Further, any of the nucleic acid molecules of the present invention may be introduced into a plant cell in a manner that allows for overexpression of the protein or fragment thereof encoded by the nucleic acid molecule.

Cosuppression is the reduction in expression levels, usually at the level of RNA, of a particular endogenous gene or gene family by the expression of a homologous sense construct that is capable of transcribing mRNA of the same strandedness as the transcript of the endogenous gene (Napoli *et al.*, *Plant Cell* 2:279-289 (1990), the entirety of which is herein incorporated by reference; van der Krol *et al.*, *Plant Cell* 2:291-299 (1990), the entirety of which is herein incorporated by reference). Cosuppression may result from stable transformation with a single copy nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell (Prolls and Meyer, *Plant J.* 2:465-475 (1992), the entirety of which is herein incorporated by reference) or with multiple copies of a nucleic acid molecule that is homologous to a nucleic acid sequence found with the cell (Mittlesten *et al.*, *Mol. Gen. Genet.* 244:325-330 (1994), the entirety of which is herein incorporated by reference). Genes, even though different, linked to

homologous promoters may result in the cosuppression of the linked genes (Vaucheret, *C.R. Acad. Sci. III* 316:1471-1483 (1993), the entirety of which is herein incorporated by reference).

This technique has, for example, been applied to generate white flowers from red petunia and tomatoes that do not ripen on the vine. Up to 50% of petunia transformants that contained a sense copy of the glucoamylase (CHS) gene produced white flowers or floral sectors; this was as a result of the post-transcriptional loss of mRNA encoding CHS (Flavell, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:3490-3496 (1994), the entirety of which is herein incorporated by reference); van Blokland *et al.*, *Plant J.* 6:861-877 (1994), the entirety of which is herein incorporated by reference). Cosuppression may require the coordinate transcription of the transgene and the endogenous gene and can be reset by a developmental control mechanism (Jorgensen, *Trends Biotechnol.* 8:340-344 (1990), the entirety of which is herein incorporated by reference; Meins and Kunz, In: *Gene Inactivation and Homologous Recombination in Plants*, Paszkowski (ed.), pp. 335-348, Kluwer Academic, Netherlands (1994), the entirety of which is herein incorporated by reference).

It is understood that one or more of the nucleic acids of the present invention may be introduced into a plant cell and transcribed using an appropriate promoter with such transcription resulting in the cosuppression of an endogenous cytokinin pathway protein.

Antisense approaches are a way of preventing or reducing gene function by targeting the genetic material (Mol *et al.*, *FEBS Lett.* 268:427-430 (1990), the entirety of which is herein incorporated by reference). The objective of the antisense approach is to use a sequence complementary to the target gene to block its expression and create a mutant cell line or organism in which the level of a single chosen protein is selectively reduced or abolished. Antisense techniques have several advantages over other 'reverse genetic' approaches. The site

of inactivation and its developmental effect can be manipulated by the choice of promoter for antisense genes or by the timing of external application or microinjection. Antisense can manipulate its specificity by selecting either unique regions of the target gene or regions where it shares homology to other related genes (Hiatt *et al.*, In: *Genetic Engineering*, Setlow (ed.), Vol. 11, New York: Plenum 49-63 (1989), the entirety of which is herein incorporated by reference).

The principle of regulation by antisense RNA is that RNA that is complementary to the target mRNA is introduced into cells, resulting in specific RNA:RNA duplexes being formed by base pairing between the antisense substrate and the target mRNA (Green *et al.*, *Annu. Rev. Biochem.* 55:569-597 (1986), the entirety of which is herein incorporated by reference). Under one embodiment, the process involves the introduction and expression of an antisense gene sequence. Such a sequence is one in which part or all of the normal gene sequences are placed under a promoter in inverted orientation so that the 'wrong' or complementary strand is transcribed into a noncoding antisense RNA that hybridizes with the target mRNA and interferes with its expression (Takayama and Inouye, *Crit. Rev. Biochem. Mol. Biol.* 25:155-184 (1990), the entirety of which is herein incorporated by reference). An antisense vector is constructed by standard procedures and introduced into cells by transformation, transfection, electroporation, microinjection, infection, etc. The type of transformation and choice of vector will determine whether expression is transient or stable. The promoter used for the antisense gene may influence the level, timing, tissue, specificity, or inducibility of the antisense inhibition.

It is understood that the activity of a cytokinin pathway protein in a plant cell may be reduced or depressed by growing a transformed plant cell containing a nucleic acid molecule whose non-transcribed strand encodes a cytokinin pathway protein or fragment thereof.

Antibodies have been expressed in plants (Hiatt *et al.*, *Nature* 342:76-78 (1989), the entirety of which is herein incorporated by reference; Conrad and Fielder, *Plant Mol. Biol.* 26:1023-1030 (1994), the entirety of which is herein incorporated by reference). Cytoplasmic expression of a scFv (single-chain Fv antibodies) has been reported to delay infection by artichoke mottled crinkle virus. Transgenic plants that express antibodies directed against endogenous proteins may exhibit a physiological effect (Philips *et al.*, *EMBO J.* 16:4489-4496 (1997), the entirety of which is herein incorporated by reference; Marion-Poll, *Trends in Plant Science* 2:447-448 (1997), the entirety of which is herein incorporated by reference). For example, expressed anti-abscisic antibodies have been reported to result in a general perturbation of seed development (Philips *et al.*, *EMBO J.* 16: 4489-4496 (1997)).

Antibodies that are catalytic may also be expressed in plants (abzymes). The principle behind abzymes is that since antibodies may be raised against many molecules, this recognition ability can be directed toward generating antibodies that bind transition states to force a chemical reaction forward (Persidas, *Nature Biotechnology* 15:1313-1315 (1997), the entirety of which is herein incorporated by reference; Baca *et al.*, *Ann. Rev. Biophys. Biomol. Struct.* 26:461-493 (1997), the entirety of which is herein incorporated by reference). The catalytic abilities of abzymes may be enhanced by site directed mutagenesis. Examples of abzymes are, for example, set forth in U.S. Patent No: 5,658,753; U.S. Patent No. 5,632,990; U.S. Patent No. 5,631,137; U.S. Patent 5,602,015; U.S. Patent No. 5,559,538; U.S. Patent No. 5,576,174; U.S. Patent No. 5,500,358; U.S. Patent 5,318,897; U.S. Patent No. 5,298,409; U.S. Patent No. 5,258,289 and U.S. Patent No. 5,194,585, all of which are herein incorporated in their entirety.

It is understood that any of the antibodies of the present invention may be expressed in plants and that such expression can result in a physiological effect. It is also understood that any of the expressed antibodies may be catalytic.

(b) Fungal Constructs and Fungal Transformants

The present invention also relates to a fungal recombinant vector comprising exogenous genetic material. The present invention also relates to a fungal cell comprising a fungal recombinant vector. The present invention also relates to methods for obtaining a recombinant fungal host cell comprising introducing into a fungal host cell exogenous genetic material.

Exogenous genetic material may be transferred into a fungal cell. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either or other nucleic acid molecule of the present invention. The fungal recombinant vector may be any vector which can be conveniently subjected to recombinant DNA procedures. The choice of a vector will typically depend on the compatibility of the vector with the fungal host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the fungal host.

The fungal vector may be an autonomously replicating vector, *i.e.*, a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the fungal cell, is integrated into the genome and

09976094-101501

replicated together with the chromosome(s) into which it has been integrated. For integration, the vector may rely on the nucleic acid sequence of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the fungal host. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, there should be preferably two nucleic acid sequences which individually contain a sufficient number of nucleic acids, preferably 400bp to 1500bp, more preferably 800bp to 1000bp, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. These nucleic acid sequences may be any sequence that is homologous with a target sequence in the genome of the fungal host cell and, furthermore, may be non-encoding or encoding sequences.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Examples of origin of replications for use in a yeast host cell are the 2 micron origin of replication and the combination of CEN3 and ARS 1. Any origin of replication may be used which is compatible with the fungal host cell of choice.

The fungal vectors of the present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides, for example biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs and the like. The selectable marker may be selected from the group including, but not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar*

(phosphinothricin acetyltransferase), *hygB* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase) and *sC* (sulfate adenylyltransferase) and *trpC* (anthranilate synthase). Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* markers of *Aspergillus nidulans* or *Aspergillus oryzae* and the bar marker of *Streptomyces hygroscopicus*. Furthermore, selection may be accomplished by co-transformation, *e.g.*, as described in WO 91/17243, the entirety of which is herein incorporated by reference. A nucleic acid sequence of the present invention may be operably linked to a suitable promoter sequence. The promoter sequence is a nucleic acid sequence which is recognized by the fungal host cell for expression of the nucleic acid sequence. The promoter sequence contains transcription and translation control sequences which mediate the expression of the protein or fragment thereof.

A promoter may be any nucleic acid sequence which shows transcriptional activity in the fungal host cell of choice and may be obtained from genes encoding polypeptides either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of a nucleic acid construct of the invention in a filamentous fungal host are promoters obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase and hybrids thereof. In a yeast host, a useful promoter is the *Saccharomyces cerevisiae* enolase (*eno-1*) promoter. Particularly preferred promoters are the TAKA amylase, NA2-tpi (a hybrid of the promoters from the genes encoding *Aspergillus niger* neutral alpha -amylase and *Aspergillus oryzae* triose phosphate isomerase) and *glaA* promoters.

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be operably linked to a terminator sequence at its 3' terminus. The terminator sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any terminator which is functional in the fungal host cell of choice may be used in the present invention, but particularly preferred terminators are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* alpha-glucosidase and *Saccharomyces cerevisiae* enolase.

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be operably linked to a suitable leader sequence. A leader sequence is a nontranslated region of a mRNA which is important for translation by the fungal host. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the protein or fragment thereof. The leader sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any leader sequence which is functional in the fungal host cell of choice may be used in the present invention, but particularly preferred leaders are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase and *Aspergillus oryzae* triose phosphate isomerase.

A polyadenylation sequence may also be operably linked to the 3' terminus of the nucleic acid sequence of the present invention. The polyadenylation sequence is a sequence which when transcribed is recognized by the fungal host to add polyadenosine residues to transcribed mRNA. The polyadenylation sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any polyadenylation sequence which is functional in the fungal host of choice may be used in the present invention, but particularly

preferred polyadenylation sequences are obtained from the genes encoding *Aspergillus oryzae* TKA amyase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase and *Aspergillus niger* alpha-glucosidase.

To avoid the necessity of disrupting the cell to obtain the protein or fragment thereof and to minimize the amount of possible degradation of the expressed protein or fragment thereof within the cell, it is preferred that expression of the protein or fragment thereof gives rise to a product secreted outside the cell. To this end, a protein or fragment thereof of the present invention may be linked to a signal peptide linked to the amino terminus of the protein or fragment thereof. A signal peptide is an amino acid sequence which permits the secretion of the protein or fragment thereof from the fungal host into the culture medium. The signal peptide may be native to the protein or fragment thereof of the invention or may be obtained from foreign sources. The 5' end of the coding sequence of the nucleic acid sequence of the present invention may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted protein or fragment thereof. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region which is foreign to that portion of the coding sequence which encodes the secreted protein or fragment thereof. The foreign signal peptide may be required where the coding sequence does not normally contain a signal peptide coding region. Alternatively, the foreign signal peptide may simply replace the natural signal peptide to obtain enhanced secretion of the desired protein or fragment thereof. The foreign signal peptide coding region may be obtained from a glucoamylase or an amyase gene from an *Aspergillus* species, a lipase or proteinase gene from *Rhizomucor miehei*, the gene for the alpha-factor from *Saccharomyces cerevisiae*, or the calf preprochymosin gene. An effective signal peptide for fungal host cells is the *Aspergillus*

oryzae TKA amylase signal, *Aspergillus niger* neutral amylase signal, the *Rhizomucor miehei* aspartic proteinase signal, the *Humicola lanuginosus* cellulase signal, or the *Rhizomucor miehei* lipase signal. However, any signal peptide capable of permitting secretion of the protein or fragment thereof in a fungal host of choice may be used in the present invention.

A protein or fragment thereof encoding nucleic acid molecule of the present invention may also be linked to a propeptide coding region. A propeptide is an amino acid sequence found at the amino terminus of a protein or proenzyme. Cleavage of the propeptide from the proprotein yields a mature biochemically active protein. The resulting polypeptide is known as a propolypeptide or proenzyme (or a zymogen in some cases). Propolypeptides are generally inactive and can be converted to mature active polypeptides by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide or proenzyme. The propeptide coding region may be native to the protein or fragment thereof or may be obtained from foreign sources. The foreign propeptide coding region may be obtained from the *Saccharomyces cerevisiae* alpha-factor gene or *Myceliophthora thermophila* laccase gene (WO 95/33836, the entirety of which is herein incorporated by reference).

The procedures used to ligate the elements described above to construct the recombinant expression vector of the present invention are well known to one skilled in the art (see, for example, Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor, N.Y., (1989)).

The present invention also relates to recombinant fungal host cells produced by the methods of the present invention which are advantageously used with the recombinant vector of the present invention. The cell is preferably transformed with a vector comprising a nucleic acid sequence of the invention followed by integration of the vector into the host chromosome. The

choice of fungal host cells will to a large extent depend upon the gene encoding the protein or fragment thereof and its source. The fungal host cell may, for example, be a yeast cell or a filamentous fungal cell.

"Yeast" as used herein includes *Ascosporogenous* yeast (*Endomycetales*), *Basidiosporogenous* yeast and yeast belonging to the *Fungi Imperfecti* (*Blastomycetes*). The *Ascosporogenous* yeasts are divided into the families *Spermophthoraceae* and *Saccharomycetaceae*. The latter is comprised of four subfamilies, *Schizosaccharomycoideae* (for example, genus *Schizosaccharomyces*), *Nadsonioideae*, *Lipomycoideae* and *Saccharomycoideae* (for example, genera *Pichia*, *Kluyveromyces* and *Saccharomyces*). The *Basidiosporogenous* yeasts include the genera *Leucosporidium*, *Rhodospodium*, *Sporidiobolus*, *Filobasidium* and *Filobasidiella*. Yeast belonging to the *Fungi Imperfecti* are divided into two families, *Sporobolomycetaceae* (for example, genera *Sorobolomyces* and *Bullera*) and *Cryptococcaceae* (for example, genus *Candida*). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (Skinner *et al.*, *Soc. App. Bacteriol. Symposium Series* No. 9, (1980), the entirety of which is herein incorporated by reference). The biology of yeast and manipulation of yeast genetics are well known in the art (see, for example, *Biochemistry and Genetics of Yeast*, Bacil *et al.* (ed.), 2nd edition, 1987; *The Yeasts*, Rose and Harrison (eds.), 2nd ed., (1987); and *The Molecular Biology of the Yeast Saccharomyces*, Strathern *et al.* (eds.), (1981), all of which are herein incorporated by reference in their entirety).

"Fungi" as used herein includes the phyla *Ascomycota*, *Basidiomycota*, *Chytridiomycota* and *Zygomycota* (as defined by Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK; the entirety of

which is herein incorporated by reference) as well as the *Oomycota* (as cited in Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK) and all mitosporic fungi (Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK). Representative groups of *Ascomycota* include, for example, *Neurospora*, *Eupenicillium* (= *Penicillium*), *Emericella* (= *Aspergillus*), *Eurotium* (= *Aspergillus*) and the true yeasts listed above. Examples of *Basidiomycota* include mushrooms, rusts and smuts. Representative groups of *Chytridiomycota* include, for example, *Allomyces*, *Blastocladiella*, *Coelomomyces* and aquatic fungi. Representative groups of *Oomycota* include, for example, *Saprolegniomycetous* aquatic fungi (water molds) such as *Achlya*. Examples of mitosporic fungi include *Aspergillus*, *Penicillium*, *Candida* and *Alternaria*. Representative groups of *Zygomycota* include, for example, *Rhizopus* and *Mucor*.

"Filamentous fungi" include all filamentous forms of the subdivision *Eumycota* and *Oomycota* (as defined by Hawksworth *et al.*, In: Ainsworth and Bisby's *Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK). The filamentous fungi are characterized by a vegetative mycelium composed of chitin, cellulose, glucan, chitosan, mannan and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

In one embodiment, the fungal host cell is a yeast cell. In a preferred embodiment, the yeast host cell is a cell of the species of *Candida*, *Kluyveromyces*, *Saccharomyces*, *Schizosaccharomyces*, *Pichia* and *Yarrowia*. In a preferred embodiment, the yeast host cell is a

Saccharomyces cerevisiae cell, a *Saccharomyces carlsbergensis*, *Saccharomyces diastaticus* cell, a *Saccharomyces douglasii* cell, a *Saccharomyces kluyveri* cell, a *Saccharomyces norbensis* cell, or a *Saccharomyces oviformis* cell. In another preferred embodiment, the yeast host cell is a *Kluyveromyces lactis* cell. In another preferred embodiment, the yeast host cell is a *Yarrowia lipolytica* cell.

In another embodiment, the fungal host cell is a filamentous fungal cell. In a preferred embodiment, the filamentous fungal host cell is a cell of the species of, but not limited to, *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Myceliophthora*, *Mucor*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium* and *Trichoderma*. In a preferred embodiment, the filamentous fungal host cell is an *Aspergillus* cell. In another preferred embodiment, the filamentous fungal host cell is an *Acremonium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Fusarium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Humicola* cell. In another preferred embodiment, the filamentous fungal host cell is a *Myceliophthora* cell. In another even preferred embodiment, the filamentous fungal host cell is a *Mucor* cell. In another preferred embodiment, the filamentous fungal host cell is a *Neurospora* cell. In another preferred embodiment, the filamentous fungal host cell is a *Penicillium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Thielavia* cell. In another preferred embodiment, the filamentous fungal host cell is a *Tolypocladium* cell. In another preferred embodiment, the filamentous fungal host cell is a *Trichoderma* cell. In a preferred embodiment, the filamentous fungal host cell is an *Aspergillus oryzae* cell, an *Aspergillus niger* cell, an *Aspergillus foetidus* cell, or an *Aspergillus japonicus* cell. In another preferred embodiment, the filamentous fungal host cell is a *Fusarium oxysporum* cell or a *Fusarium graminearum* cell. In another preferred embodiment, the filamentous fungal

host cell is a *Humicola insolens* cell or a *Humicola lanuginosus* cell. In another preferred embodiment, the filamentous fungal host cell is a *Myceliophthora thermophila* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Mucor miehei* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Neurospora crassa* cell. In a most preferred embodiment, the filamentous fungal host cell is a *Penicillium purpurogenum* cell. In another most preferred embodiment, the filamentous fungal host cell is a *Thielavia terrestris* cell. In another most preferred embodiment, the *Trichoderma* cell is a *Trichoderma reesei* cell, a *Trichoderma viride* cell, a *Trichoderma longibrachiatum* cell, a *Trichoderma harzianum* cell, or a *Trichoderma koningii* cell. In a preferred embodiment, the fungal host cell is selected from an *A. nidulans* cell, an *A. niger* cell, an *A. oryzae* cell and an *A. sojae* cell. In a further preferred embodiment, the fungal host cell is an *A. nidulans* cell.

The recombinant fungal host cells of the present invention may further comprise one or more sequences which encode one or more factors that are advantageous in the expression of the protein or fragment thereof, for example, an activator (e.g., a trans-acting factor), a chaperone and a processing protease. The nucleic acids encoding one or more of these factors are preferably not operably linked to the nucleic acid encoding the protein or fragment thereof. An activator is a protein which activates transcription of a nucleic acid sequence encoding a polypeptide (Kudla *et al.*, *EMBO* 9:1355-1364(1990); Jarai and Buxton, *Current Genetics* 26:2238-244(1994); Verdier, *Yeast* 6:271-297(1990), all of which are herein incorporated by reference in their entirety). The nucleic acid sequence encoding an activator may be obtained from the genes encoding *Saccharomyces cerevisiae* heme activator protein 1 (hap1), *Saccharomyces cerevisiae* galactose metabolizing protein 4 (gal4) and *Aspergillus nidulans* ammonia regulation protein (areA). For further examples, see Verdier, *Yeast* 6:271-297 (1990);

MacKenzie *et al.*, *Journal of Gen. Microbiol.* 139:2295-2307 (1993), both of which are herein incorporated by reference in their entirety). A chaperone is a protein which assists another protein in folding properly (Hartl *et al.*, *TIBS* 19:20-25 (1994); Bergeron *et al.*, *TIBS* 19:124-128 (1994); Demolder *et al.*, *J. Biotechnology* 32:179-189 (1994); Craig, *Science* 260:1902-1903 (1993); Gething and Sambrook, *Nature* 355:33-45 (1992); Puig and Gilbert, *J Biol. Chem.* 269:7764-7771 (1994); Wang and Tsou, *FASEB Journal* 7:1515-11157 (1993); Robinson *et al.*, *Bio/Technology* 1:381-384 (1994), all of which are herein incorporated by reference in their entirety). The nucleic acid sequence encoding a chaperone may be obtained from the genes encoding *Aspergillus oryzae* protein disulphide isomerase, *Saccharomyces cerevisiae* calnexin, *Saccharomyces cerevisiae* BiP/GRP78 and *Saccharomyces cerevisiae* Hsp70. For further examples, see Gething and Sambrook, *Nature* 355:33-45 (1992); Hartl *et al.*, *TIBS* 19:20-25 (1994). A processing protease is a protease that cleaves a propeptide to generate a mature biochemically active polypeptide (Enderlin and Ogrydziak, *Yeast* 10:67-79 (1994); Fuller *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 86:1434-1438 (1989); Julius *et al.*, *Cell* 37:1075-1089 (1984); Julius *et al.*, *Cell* 32:839-852 (1983), all of which are incorporated by reference in their entirety). The nucleic acid sequence encoding a processing protease may be obtained from the genes encoding *Aspergillus niger* Kex2, *Saccharomyces cerevisiae* dipeptidylaminopeptidase, *Saccharomyces cerevisiae* Kex2 and *Yarrowia lipolytica* dibasic processing endoprotease (xpr6). Any factor that is functional in the fungal host cell of choice may be used in the present invention.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts and regeneration of the cell wall in a manner known per se. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and

Yelton *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 81:1470-1474 (1984), both of which are herein incorporated by reference in their entirety. A suitable method of transforming *Fusarium* species is described by Malardier *et al.*, *Gene* 78:147-156 (1989), the entirety of which is herein incorporated by reference. Yeast may be transformed using the procedures described by Becker and Guarente, In: Abelson and Simon, (eds.), *Guide to Yeast Genetics and Molecular Biology, Methods Enzymol.* Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, *J. Bacteriology* 153:163 (1983); Hinnen *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 75:1920 (1978), all of which are herein incorporated by reference in their entirety.

The present invention also relates to methods of producing the protein or fragment thereof comprising culturing the recombinant fungal host cells under conditions conducive for expression of the protein or fragment thereof. The fungal cells of the present invention are cultivated in a nutrient medium suitable for production of the protein or fragment thereof using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the protein or fragment thereof to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art (*see, e.g.*, Bennett and LaSure (eds.), *More Gene Manipulations in Fungi*, Academic Press, CA, (1991), the entirety of which is herein incorporated by reference). Suitable media are available from commercial suppliers or may be prepared according to published compositions (*e.g.*, in catalogues of the American Type Culture Collection, Manassas, VA). If the protein or fragment thereof is secreted into the nutrient

medium, a protein or fragment thereof can be recovered directly from the medium. If the protein or fragment thereof is not secreted, it is recovered from cell lysates.

The expressed protein or fragment thereof may be detected using methods known in the art that are specific for the particular protein or fragment. These detection methods may include the use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, if the protein or fragment thereof has enzymatic activity, an enzyme assay may be used. Alternatively, if polyclonal or monoclonal antibodies specific to the protein or fragment thereof are available, immunoassays may be employed using the antibodies to the protein or fragment thereof. The techniques of enzyme assay and immunoassay are well known to those skilled in the art.

The resulting protein or fragment thereof may be recovered by methods known in the arts. For example, the protein or fragment thereof may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation. The recovered protein or fragment thereof may then be further purified by a variety of chromatographic procedures, e.g., ion exchange chromatography, gel filtration chromatography, affinity chromatography, or the like.

(c) Mammalian Constructs and Transformed Mammalian Cells

The present invention also relates to methods for obtaining a recombinant mammalian host cell, comprising introducing into a mammalian host cell exogenous genetic material. The present invention also relates to a mammalian cell comprising a mammalian recombinant vector. The present invention also relates to methods for obtaining a recombinant mammalian host cell, comprising introducing into a mammalian cell exogenous genetic material. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present

invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines. Suitable promoters for mammalian cells are also known in the art and include viral promoters such as that from Simian Virus 40 (SV40) (Fiers *et al.*, *Nature* 273:113 (1978), the entirety of which is herein incorporated by reference), Rous sarcoma virus (RSV), adenovirus (ADV) and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences and poly-A addition sequences. Enhancer sequences which increase expression may also be included and sequences which promote amplification of the gene may also be desirable (for example methotrexate resistance genes).

Vectors suitable for replication in mammalian cells may include viral replicons, or sequences which insure integration of the appropriate sequences encoding HCV epitopes into the host genome. For example, another vector used to express foreign DNA is vaccinia virus. In this case, for example, a nucleic acid molecule encoding a protein or fragment thereof is inserted into the vaccinia genome. Techniques for the insertion of foreign DNA into the vaccinia virus genome are known in the art and may utilize, for example, homologous recombination. Such heterologous DNA is generally inserted into a gene which is non-essential to the virus, for example, the thymidine kinase gene (tk), which also provides a selectable marker. Plasmid vectors that greatly facilitate the construction of recombinant viruses have been described (*see*, for example, Mackett *et al.*, *J Virol.* 49:857 (1984); Chakrabarti *et al.*, *Mol. Cell. Biol.* 5:3403

(1985); Moss, In: *Gene Transfer Vectors For Mammalian Cells* (Miller and Calos, eds., Cold Spring Harbor Laboratory, N.Y., p. 10, (1987); all of which are herein incorporated by reference in their entirety). Expression of the HCV polypeptide then occurs in cells or animals which are infected with the live recombinant vaccinia virus.

The sequence to be integrated into the mammalian sequence may be introduced into the primary host by any convenient means, which includes calcium precipitated DNA, spheroplast fusion, transformation, electroporation, biolistics, lipofection, microinjection, or other convenient means. Where an amplifiable gene is being employed, the amplifiable gene may serve as the selection marker for selecting hosts into which the amplifiable gene has been introduced.

Alternatively, one may include with the amplifiable gene another marker, such as a drug resistance marker, e.g. neomycin resistance (G418 in mammalian cells), hygromycin in resistance etc., or an auxotrophy marker (HIS3, TRP1, LEU2, URA3, ADE2, LYS2, etc.) for use in yeast cells.

Depending upon the nature of the modification and associated targeting construct, various techniques may be employed for identifying targeted integration. Conveniently, the DNA may be digested with one or more restriction enzymes and the fragments probed with an appropriate DNA fragment which will identify the properly sized restriction fragment associated with integration.

One may use different promoter sequences, enhancer sequences, or other sequence which will allow for enhanced levels of expression in the expression host. Thus, one may combine an enhancer from one source, a promoter region from another source, a 5'- noncoding region upstream from the initiation cytokinin from the same or different source as the other sequences and the like. One may provide for an intron in the non-coding region with appropriate splice

sites or for an alternative 3'- untranslated sequence or polyadenylation site. Depending upon the particular purpose of the modification, any of these sequences may be introduced, as desired.

Where selection is intended, the sequence to be integrated will have with it a marker gene, which allows for selection. The marker gene may conveniently be downstream from the target gene and may include resistance to a cytotoxic agent, e.g. antibiotics, heavy metals, or the like, resistance or susceptibility to HAT, gancyclovir, etc., complementation to an auxotrophic host, particularly by using an auxotrophic yeast as the host for the subject manipulations, or the like. The marker gene may also be on a separate DNA molecule, particularly with primary mammalian cells. Alternatively, one may screen the various transformants, due to the high efficiency of recombination in yeast, by using hybridization analysis, PCR, sequencing, or the like.

For homologous recombination, constructs can be prepared where the amplifiable gene will be flanked, normally on both sides with DNA homologous with the DNA of the target region. Depending upon the nature of the integrating DNA and the purpose of the integration, the homologous DNA will generally be within 100kb, usually 50kb, preferably about 25kb, of the transcribed region of the target gene, more preferably within 2kb of the target gene. Where modeling of the gene is intended, homology will usually be present proximal to the site of the mutation. The homologous DNA may include the 5'-upstream region outside of the transcriptional regulatory region or comprising any enhancer sequences, transcriptional initiation sequences, adjacent sequences, or the like. The homologous region may include a portion of the coding region, where the coding region may be comprised only of an open reading frame or combination of exons and introns. The homologous region may comprise all or a portion of an intron, where all or a portion of one or more exons may also be present. Alternatively, the

homologous region may comprise the 3'-region, so as to comprise all or a portion of the transcriptional termination region, or the region 3' of this region. The homologous regions may extend over all or a portion of the target gene or be outside the target gene comprising all or a portion of the transcriptional regulatory regions and/or the structural gene.

The integrating constructs may be prepared in accordance with conventional ways, where sequences may be synthesized, isolated from natural sources, manipulated, cloned, ligated, subjected to in vitro mutagenesis, primer repair, or the like. At various stages, the joined sequences may be cloned and analyzed by restriction analysis, sequencing, or the like. Usually during the preparation of a construct where various fragments are joined, the fragments, intermediate constructs and constructs will be carried on a cloning vector comprising a replication system functional in a prokaryotic host, e.g., *E. coli* and a marker for selection, e.g., biocide resistance, complementation to an auxotrophic host, etc. Other functional sequences may also be present, such as polylinkers, for ease of introduction and excision of the construct or portions thereof, or the like. A large number of cloning vectors are available such as pBR322, the pUC series, etc. These constructs may then be used for integration into the primary mammalian host.

In the case of the primary mammalian host, a replicating vector may be used. Usually, such vector will have a viral replication system, such as SV40, bovine papilloma virus, adenovirus, or the like. The linear DNA sequence vector may also have a selectable marker for identifying transfected cells. Selectable markers include the neo gene, allowing for selection with G418, the herpes tk gene for selection with HAT medium, the gpt gene with mycophenolic acid, complementation of an auxotrophic host, etc.

The vector may or may not be capable of stable maintenance in the host. Where the vector is capable of stable maintenance, the cells will be screened for homologous integration of the vector into the genome of the host, where various techniques for curing the cells may be employed. Where the vector is not capable of stable maintenance, for example, where a temperature sensitive replication system is employed, one may change the temperature from the permissive temperature to the non-permissive temperature, so that the cells may be cured of the vector. In this case, only those cells having integration of the construct comprising the amplifiable gene and, when present, the selectable marker, will be able to survive selection.

Where a selectable marker is present, one may select for the presence of the targeting construct by means of the selectable marker. Where the selectable marker is not present, one may select for the presence of the construct by the amplifiable gene. For the neo gene or the herpes tk gene, one could employ a medium for growth of the transformants of about 0.1-1 mg/ml of G418 or may use HAT medium, respectively. Where DHFR is the amplifiable gene, the selective medium may include from about 0.01-0.5 M of methotrexate or be deficient in glycine-hypoxanthine-thymidine and have dialysed serum (GHT media).

The DNA can be introduced into the expression host by a variety of techniques that include calcium phosphate/DNA co-precipitates, microinjection of DNA into the nucleus, electroporation, yeast protoplast fusion with intact cells, transfection, polycations, e.g., polybrene, polyornithine, etc., or the like. The DNA may be single or double stranded DNA, linear or circular. The various techniques for transforming mammalian cells are well known (see Keown *et al.*, *Methods Enzymol.* (1989); Keown *et al.*, *Methods Enzymol.* 185:527-537 (1990); Mansour *et al.*, *Nature* 336:348-352, (1988); all of which are herein incorporated by reference in their entirety).

(d) Insect Constructs and Transformed Insect Cells

The present invention also relates to an insect recombinant vectors comprising exogenous genetic material. The present invention also relates to an insect cell comprising an insect recombinant vector. The present invention also relates to methods for obtaining a recombinant insect host cell, comprising introducing into an insect cell exogenous genetic material. In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

The insect recombinant vector may be any vector which can be conveniently subjected to recombinant DNA procedures and can bring about the expression of the nucleic acid sequence. The choice of a vector will typically depend on the compatibility of the vector with the insect host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the insect host. In addition, the insect vector may be an expression vector. Nucleic acid molecules can be suitably inserted into a replication vector for expression in the insect cell under a suitable promoter for insect cells. Many vectors are available for this purpose and selection of the appropriate vector will depend mainly on the size of the nucleic acid molecule to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the particular host cell with which it is compatible. The vector components for insect cell

transformation generally include, but are not limited to, one or more of the following: a signal sequence, origin of replication, one or more marker genes and an inducible promoter.

The insect vector may be an autonomously replicating vector, *i.e.*, a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the insect cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. For integration, the vector may rely on the nucleic acid sequence of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the insect host. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, there should be preferably two nucleic acid sequences which individually contain a sufficient number of nucleic acids, preferably 400bp to 1500bp, more preferably 800bp to 1000bp, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. These nucleic acid sequences may be any sequence that is homologous with a target sequence in the genome of the insect host cell and, furthermore, may be non-encoding or encoding sequences.

Baculovirus expression vectors (BEVs) have become important tools for the expression of foreign genes, both for basic research and for the production of proteins with direct clinical applications in human and veterinary medicine (Doerfler, *Curr. Top. Microbiol. Immunol.*

131:51-68 (1968); Luckow and Summers, *Bio/Technology* 6:47-55 (1988a); Miller, *Annual Review of Microbiol.* 42:177-199 (1988); Summers, *Curr. Comm. Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); all of which are herein incorporated by reference in their entirety). BEVs are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter in place of the viral gene, e.g., polyhedrin (Smith and Summers, U.S. Pat. No., 4,745,051, the entirety of which is incorporated herein by reference).

The use of baculovirus vectors relies upon the host cells being derived from *Lepidopteran* insects such as *Spodoptera frugiperda* or *Trichoplusia ni*. The preferred *Spodoptera frugiperda* cell line is the cell line Sf9. The *Spodoptera frugiperda* Sf9 cell line was obtained from American Type Culture Collection (Manassas, VA.) and is assigned accession number ATCC CRL 1711 (Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), the entirety of which is herein incorporated by reference). Other insect cell systems, such as the silkworm *B. mori* may also be used.

The proteins expressed by the BEVs are, therefore, synthesized, modified and transported in host cells derived from *Lepidopteran* insects. Most of the genes that have been inserted and produced in the baculovirus expression vector system have been derived from vertebrate species. Other baculovirus genes in addition to the polyhedrin promoter may be employed to advantage in a baculovirus expression system. These include immediate-early (alpha), delayed-early (), late (), or very late (delta), according to the phase of the viral infection during which they are expressed. The expression of these genes occurs sequentially, probably as the result of a "cascade" mechanism of transcriptional regulation. (Guarino and Summers, *J. Virol.* 57:563-571

(1986); Guarino and Summers, *J. Virol.* 61:2091-2099 (1987); Guarino and Summers, *Virol.* 162:444-451 (1988); all of which are herein incorporated by reference in their entirety).

Insect recombinant vectors are useful as intermediates for the infection or transformation of insect cell systems. For example, an insect recombinant vector containing a nucleic acid molecule encoding a baculovirus transcriptional promoter followed downstream by an insect signal DNA sequence is capable of directing the secretion of the desired biologically active protein from the insect cell. The vector may utilize a baculovirus transcriptional promoter region derived from any of the over 500 baculoviruses generally infecting insects, such as for example the Orders *Lepidoptera*, *Diptera*, *Orthoptera*, *Coleoptera* and *Hymenoptera*, including for example but not limited to the viral DNAs of *Autographa californica* MNPV, *Bombyx mori* NPV, *Trichoplusia ni* MNPV, *Rachiplusia ou* MNPV or *Galleria mellonella* MNPV, wherein said baculovirus transcriptional promoter is a baculovirus immediate-early gene IEL or IEN promoter; an immediate-early gene in combination with a baculovirus delayed-early gene promoter region selected from the group consisting of 39K and a *HindIII-k* fragment delayed-early gene; or a baculovirus late gene promoter. The immediate-early or delayed-early promoters can be enhanced with transcriptional enhancer elements. The insect signal DNA sequence may code for a signal peptide of a *Lepidopteran* adipokinetic hormone precursor or a signal peptide of the *Manduca sexta* adipokinetic hormone precursor (Summers, U.S. Patent No. 5,155,037; the entirety of which is herein incorporated by reference). Other insect signal DNA sequences include a signal peptide of the *Orthoptera Schistocerca gregaria* locust adipokinetic hormone precursor and the *Drosophila melanogaster* cuticle genes CP1, CP2, CP3 or CP4 or for an insect signal peptide having substantially a similar chemical composition and function (Summers, U.S. Patent No. 5,155,037).

05976094-104604

Insect cells are distinctly different from animal cells. Insects have a unique life cycle and have distinct cellular properties such as the lack of intracellular plasminogen activators in which are present in vertebrate cells. Another difference is the high expression levels of protein products ranging from 1 to greater than 500 mg/liter and the ease at which cDNA can be cloned into cells (Frasier, *In Vitro Cell. Dev. Biol.* 25:225 (1989); Summers and Smith, In: *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), both of which are incorporated by reference in their entirety).

Recombinant protein expression in insect cells is achieved by viral infection or stable transformation. For viral infection, the desired gene is cloned into baculovirus at the site of the wild-type polyhedron gene (Webb and Summers, *Technique* 2:173 (1990); Bishop and Posse, *Adv. Gene Technol.* 1:55 (1990); both of which are incorporated by reference in their entirety). The polyhedron gene is a component of a protein coat in occlusions which encapsulate virus particles. Deletion or insertion in the polyhedron gene results the failure to form occlusion bodies. Occlusion negative viruses are morphologically different from occlusion positive viruses and enable one skilled in the art to identify and purify recombinant viruses.

The vectors of present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides, for example biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs and the like. Selection may be accomplished by co-transformation, *e.g.*, as described in WO 91/17243, a nucleic acid sequence of the present invention may be operably linked to a suitable promoter sequence. The promoter sequence is a nucleic acid sequence which is recognized by the insect host cell for expression of the nucleic acid sequence. The promoter sequence contains transcription and translation control sequences which mediate the expression

of the protein or fragment thereof. The promoter may be any nucleic acid sequence which shows transcriptional activity in the insect host cell of choice and may be obtained from genes encoding polypeptides either homologous or heterologous to the host cell.

For example, a nucleic acid molecule encoding a protein or fragment thereof may also be operably linked to a suitable leader sequence. A leader sequence is a nontranslated region of a mRNA which is important for translation by the fungal host. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the protein or fragment thereof. The leader sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any leader sequence which is functional in the insect host cell of choice may be used in the present invention.

A polyadenylation sequence may also be operably linked to the 3' terminus of the nucleic acid sequence of the present invention. The polyadenylation sequence is a sequence which when transcribed is recognized by the insect host to add polyadenosine residues to transcribed mRNA. The polyadenylation sequence may be native to the nucleic acid sequence encoding the protein or fragment thereof or may be obtained from foreign sources. Any polyadenylation sequence which is functional in the fungal host of choice may be used in the present invention.

To avoid the necessity of disrupting the cell to obtain the protein or fragment thereof and to minimize the amount of possible degradation of the expressed polypeptide within the cell, it is preferred that expression of the polypeptide gene gives rise to a product secreted outside the cell. To this end, the protein or fragment thereof of the present invention may be linked to a signal peptide linked to the amino terminus of the protein or fragment thereof. A signal peptide is an amino acid sequence which permits the secretion of the protein or fragment thereof from the insect host into the culture medium. The signal peptide may be native to the protein or fragment

thereof of the invention or may be obtained from foreign sources. The 5' end of the coding sequence of the nucleic acid sequence of the present invention may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted protein or fragment thereof.

At present, a mode of achieving secretion of a foreign gene product in insect cells is by way of the foreign gene's native signal peptide. Because the foreign genes are usually from non-insect organisms, their signal sequences may be poorly recognized by insect cells and hence, levels of expression may be suboptimal. However, the efficiency of expression of foreign gene products seems to depend primarily on the characteristics of the foreign protein. On average, nuclear localized or non-structural proteins are most highly expressed, secreted proteins are intermediate and integral membrane proteins are the least expressed. One factor generally affecting the efficiency of the production of foreign gene products in a heterologous host system is the presence of native signal sequences (also termed presequences, targeting signals, or leader sequences) associated with the foreign gene. The signal sequence is generally coded by a DNA sequence immediately following (5' to 3') the translation start site of the desired foreign gene.

The expression dependence on the type of signal sequence associated with a gene product can be represented by the following example: If a foreign gene is inserted at a site downstream from the translational start site of the baculovirus polyhedrin gene so as to produce a fusion protein (containing the N-terminus of the polyhedrin structural gene), the fused gene is highly expressed. But less expression is achieved when a foreign gene is inserted in a baculovirus expression vector immediately following the transcriptional start site and totally replacing the polyhedrin structural gene.

Insertions into the region -50 to -1 significantly alter (reduce) steady state transcription which, in turn, reduces translation of the foreign gene product. Use of the pVL941 vector optimizes transcription of foreign genes to the level of the polyhedrin gene transcription. Even though the transcription of a foreign gene may be optimal, optimal translation may vary because of several factors involving processing: signal peptide recognition, mRNA and ribosome binding, glycosylation, disulfide bond formation, sugar processing, oligomerization, for example.

The properties of the insect signal peptide are expected to be more optimal for the efficiency of the translation process in insect cells than those from vertebrate proteins. This phenomenon can generally be explained by the fact that proteins secreted from cells are synthesized as precursor molecules containing hydrophobic N-terminal signal peptides. The signal peptides direct transport of the select protein to its target membrane and are then cleaved by a peptidase on the membrane, such as the endoplasmic reticulum, when the protein passes through it.

Another exemplary insect signal sequence is the sequence encoding for *Drosophila* cuticle proteins such as CP1, CP2, CP3 or CP4 (Summers, U.S. Patent No. 5,278,050; the entirety of which is herein incorporated by reference). Most of a 9kb region of the *Drosophila* genome containing genes for the cuticle proteins has been sequenced. Four of the five cuticle genes contains a signal peptide coding sequence interrupted by a short intervening sequence (about 60 base pairs) at a conserved site. Conserved sequences occur in the 5' mRNA untranslated region, in the adjacent 35 base pairs of upstream flanking sequence and at -200 base pairs from the mRNA start position in each of the cuticle genes.

Standard methods of insect cell culture, cotransfection and preparation of plasmids are set forth in Summers and Smith (Summers and Smith, *A Manual of Methods for Baculovirus*

Vectors and Insect Cell Culture Procedures, Texas Agricultural Experiment Station Bulletin No. 1555, Texas A&M University (1987)). Procedures for the cultivation of viruses and cells are described in Volkman and Summers, *J. Virol* 19:820-832 (1975) and Volkman *et al.*, *J. Virol* 19:820-832 (1976); both of which are herein incorporated by reference in their entirety.

(e) Bacterial Constructs and Transformed Bacterial Cells

The present invention also relates to a bacterial recombinant vector comprising exogenous genetic material. The present invention also relates to a bacteria cell comprising a bacterial recombinant vector. The present invention also relates to methods for obtaining a recombinant bacteria host cell, comprising introducing into a bacterial host cell exogenous genetic material. . In a preferred embodiment the exogenous genetic material includes a nucleic acid molecule of the present invention having a sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 711 or complements thereof or fragments of either or other nucleic acid molecule of the present invention.

The bacterial recombinant vector may be any vector which can be conveniently subjected to recombinant DNA procedures. The choice of a vector will typically depend on the compatibility of the vector with the bacterial host cell into which the vector is to be introduced. The vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the bacterial host. In addition, the bacterial vector may be an expression vector. Nucleic acid molecules encoding protein homologues or fragments thereof can, for example, be suitably inserted into a replicable vector for expression in the bacterium under the control of a suitable promoter for bacteria. Many vectors are available for this purpose and selection of the appropriate vector will depend mainly on the size of the nucleic acid to be

inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the particular host cell with which it is compatible. The vector components for bacterial transformation generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes and an inducible promoter.

In general, plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell are used in connection with bacterial hosts. The vector ordinarily carries a replication site, as well as marking sequences that are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species (see, e.g., Bolivar *et al.*, *Gene* 2:95 (1977); the entirety of which is herein incorporated by reference). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, or other microbial plasmid or phage, also generally contains, or is modified to contain, promoters that can be used by the microbial organism for expression of the selectable marker genes.

Nucleic acid molecules encoding protein or fragments thereof may be expressed not only directly, but also as a fusion with another polypeptide, preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the polypeptide DNA that is inserted into the vector. The heterologous signal sequence selected should be one that is recognized and processed (i.e., cleaved by a signal peptidase) by the host cell. For bacterial host cells that do not recognize and process the native polypeptide signal

sequence, the signal sequence is substituted by a bacterial signal sequence selected, for example, from the group consisting of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria.

Expression and cloning vectors also generally contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous protein homologue or fragment thereof produce a protein conferring drug resistance and thus survive the selection regimen.

The expression vector for producing a protein or fragment thereof can also contain an inducible promoter that is recognized by the host bacterial organism and is operably linked to the nucleic acid encoding, for example, the nucleic acid molecule encoding the protein homologue or fragment thereof of interest. Inducible promoters suitable for use with bacterial hosts include the

09976064-101501
-lactamase and lactose promoter systems (Chang *et al.*, *Nature* 275:615 (1978); Goeddel *et al.*, *Nature* 281:544 (1979); both of which are herein incorporated by reference in their entirety), the arabinose promoter system (Guzman *et al.*, *J. Bacteriol.* 174:7716-7728 (1992); the entirety of which is herein incorporated by reference), alkaline phosphatase, a tryptophan (trp) promoter system (Goeddel, *Nucleic Acids Res.* 8:4057 (1980); EP 36,776; both of which are herein incorporated by reference in their entirety) and hybrid promoters such as the tac promoter (deBoer *et al.*, *Proc. Natl. Acad. Sci. (USA)* 80:21-25 (1983); the entirety of which is herein incorporated by reference). However, other known bacterial inducible promoters are suitable (Siebenlist *et al.*, *Cell* 20:269 (1980); the entirety of which is herein incorporated by reference).

Promoters for use in bacterial systems also generally contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the polypeptide of interest. The promoter can be removed from the bacterial source DNA by restriction enzyme digestion and inserted into the vector containing the desired DNA.

Construction of suitable vectors containing one or more of the above-listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored and re-ligated in the form desired to generate the plasmids required. Examples of available bacterial expression vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as Bluescript™ (Stratagene, La Jolla, CA), in which, for example, encoding an *A. nidulans* protein homologue or fragment thereof homologue, may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke and Schuster, *J. Biol. Chem.* 264:5503-5509 (1989), the entirety of which is herein incorporated by reference); and the

like. pGEX vectors (Promega, Madison Wisconsin U.S.A.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems are designed to include heparin, thrombin or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Suitable host bacteria for a bacterial vector include archaebacteria and eubacteria, especially eubacteria and most preferably *Enterobacteriaceae*. Examples of useful bacteria include *Escherichia*, *Enterobacter*, *Azotobacter*, *Erwinia*, *Bacillus*, *Pseudomonas*, *Klebsiella*, *Proteus*, *Salmonella*, *Serratia*, *Shigella*, *Rhizobia*, *Vitreoscilla* and *Paracoccus*. Suitable *E. coli* hosts include *E. coli* W3110 (American Type Culture Collection (ATCC) 27,325, Manassas, Virginia U.S.A.), *E. coli* 294 (ATCC 31,446), *E. coli* B and *E. coli* X1776 (ATCC 31,537). These examples are illustrative rather than limiting. Mutant cells of any of the above-mentioned bacteria may also be employed. It is, of course, necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E. coli*, *Serratia*, or *Salmonella* species can be suitably used as the host when well known plasmids such as pBR322, pBR325, pACYC177, or pKN410 are used to supply the replicon. *E. coli* strain W3110 is a preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell should secrete minimal amounts of proteolytic enzymes.

Host cells are transfected and preferably transformed with the above-described vectors and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

09976094-404604

Numerous methods of transfection are known to the ordinarily skilled artisan, for example, calcium phosphate and electroporation. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Laboratory Press, (1989), is generally used for bacterial cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO, as described in Chung and Miller (Chung and Miller, *Nucleic Acids Res.* 16:3580 (1988); the entirety of which is herein incorporated by reference). Yet another method is the use of the technique termed electroporation.

Bacterial cells used to produce the polypeptide of interest for purposes of this invention are cultured in suitable media in which the promoters for the nucleic acid encoding the heterologous polypeptide can be artificially induced as described generally, e.g., in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Laboratory Press, (1989). Examples of suitable media are given in U.S. Pat. Nos. 5,304,472 and 5,342,763; both of which are incorporated by reference in their entirety.

In addition to the above discussed procedures, practitioners are familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of recombinant organisms and the screening and isolating of clones, (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Mailga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995), the entirety of which

is herein incorporated by reference: Birren *et al.*, *Genome Analysis: Analyzing DNA*, 1, Cold Spring Harbor, New York, the entirety of which is herein incorporated by reference).

(f) Computer Readable Media

The nucleotide sequence provided in SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof, or complement thereof, or a nucleotide sequence at least 90% identical, preferably 95%, identical even more preferably 99% or 100% identical to the sequence provided in SEQ ID NO: 1 through SEQ ID NO: 711 or fragment thereof, or complement thereof, can be "provided" in a variety of mediums to facilitate use. Such a medium can also provide a subset thereof in a form that allows a skilled artisan to examine the sequences.

A preferred subset of nucleotide sequences are those nucleic acid sequences that encode a maize or a soybean adenine phosphoribosyl transferase enzyme or complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or complement thereof or fragment of either and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or complement thereof or fragment of either.

A further preferred subset of nucleic acid sequences is where the subset of sequences is two proteins or fragments thereof, more preferably three proteins or fragments thereof and even more preferable four proteins or fragments thereof, these nucleic acid sequences are selected from the group that comprises a maize or a soybean adenine phosphoribosyl transferase enzyme or complement thereof or fragment of either, a nucleic acid molecule that encodes a maize or a soybean β glucosidase enzyme or complement thereof or fragment of either and a nucleic acid molecule that encodes a soybean isopentyltransferase enzyme or complement thereof or fragment of either.

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc, storage medium and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate media comprising the nucleotide sequence information of the present invention. A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (e.g. text file or database) in order to

obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing one or more of nucleotide sequences of the present invention, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990), the entirety of which is herein incorporated by reference) and BLAZE (Brutlag *et al.*, *Comp. Chem.* 17:203-207 (1993), the entirety of which is herein incorporated by reference) search algorithms on a Sybase system can be used to identify open reading frames (ORFs) within the genome that contain homology to ORFs or proteins from other organisms. Such ORFs are protein-encoding fragments within the sequences of the present invention and are useful in producing commercially important proteins such as enzymes used in amino acid biosynthesis, metabolism, transcription, translation, RNA processing, nucleic acid and a protein degradation, protein modification and DNA replication, restriction, modification, recombination and repair.

The present invention further provides systems, particularly computer-based systems, which contain the sequence information described herein. Such systems are designed to identify commercially important fragments of the nucleic acid molecule of the present invention. As used herein, "a computer-based system" refers to the hardware means, software means and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means and data storage means. A skilled

artisan can readily appreciate that any one of the currently available computer-based system are suitable for use in the present invention.

As indicated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory that can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention. As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequence of the present invention that match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are available can be used in the computer-based systems of the present invention. Examples of such software include, but are not limited to, MacPattern (EMBL), BLASTIN and BLASTIX (NCBIA). One of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based systems.

The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that during searches for commercially important fragments of the nucleic acid molecules of the present invention, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequences the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzymatic active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, *cis* elements, hairpin structures and inducible expression elements (protein binding sequences).

Thus, the present invention further provides an input means for receiving a target sequence, a data storage means for storing the target sequences of the present invention sequence identified using a search means as described above and an output means for outputting the identified homologous sequences. A variety of structural formats for the input and output means can be used to input and output information in the computer-based systems of the present invention. A preferred format for an output means ranks fragments of the sequence of the present invention by varying degrees of homology to the target sequence or target motif. Such presentation provides a skilled artisan with a ranking of sequences which contain various amounts of the target sequence or target motif and identifies the degree of homology contained in the identified fragment.

A variety of comparing means can be used to compare a target sequence or target motif with the data storage means to identify sequence fragments sequence of the present invention. For example, implementing software which implement the BLAST and BLAZE algorithms (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990)) can be used to identify open frames within the nucleic acid molecules of the present invention. A skilled artisan can readily recognize that any

one of the publicly available homology search programs can be used as the search means for the computer-based systems of the present invention.

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration and are not intended to be limiting of the present invention, unless specified.

Example 1

The MONN01 cDNA library is a normalized library generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation.

09976034-101501

The SATMON001 cDNA library is generated from maize (B73, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) immature tassels at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant is collected at the V6 stage. At that stage the tassel is an immature tassel of about 2-3 cm in length. The tassels are removed and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON003 library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) roots at the V6 developmental stage. Seeds are planted at a depth of approximately 3 cm in coil into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth, the seedlings are transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and approximately 3 times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting at a concentration of 150 ppm N. Two to three times during the life time of the plant from transplanting to flowering a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in approximately 15hr day/9hr

night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6 leaf development stage. The root system is cut from maize plant and washed with water to free it from the soil. The tissue is then immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON004 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON005 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) root tissue at the V6 development stage. Seeds

are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen and the harvested tissue is then stored at -80°C until RNA preparation.

The SATMON006 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental

lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON007 cDNA library is generated from the primary root tissue of 5 day old maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). After germination, the trays, along with the moist paper, are moved to a greenhouse where the maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles for approximately 5 days. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F . Supplemental lighting is provided by 1000 W sodium vapor lamps. The primary root tissue is collected when the seedlings are 5 days old. At this stage, the primary root (radicle) is pushed through the coleorhiza which itself is pushed through the seed coat. The primary root, which is about 2-3 cm long, is cut and immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON008 cDNA library is generated from the primary shoot (coleoptile 2-3 cm) of maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings which are approximately 5 days old. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). Then the trays containing the seeds are moved to a greenhouse at 15hr daytime/9 hr nighttime cycles and grown until they are 5 days post germination. The daytime temperature is approximately 80°F and the nighttime temperature is approximately

70°F. Tissue is collected when the seedlings are 5 days old. At this stage, the primary shoot (coleoptile) is pushed through the seed coat and is about 2-3 cm long. The coleoptile is dissected away from the rest of the seedling, immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON009 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaves at the 8 leaf stage (V8 plant development stage). Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 8-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical, are cut at the base of the leaves. The leaves are then pooled and then immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON010 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) root tissue at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium.

After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the V8 development stage. The root system is cut from this mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON011 cDNA library is generated from undeveloped maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaf at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The second youngest leaf which is at the base of the apical leaf of V6 stage maize plant is cut at the base and

immediately transferred to liquid nitrogen containers in which the leaf is crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON012 cDNA library is generated from 2 day post germination maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination (one day). Then the trays containing the seeds are moved to the greenhouse and grown at 15hr daytime/9 hr nighttime cycles until 2 days post germination. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Tissue is collected when the seedlings are 2 days old. At the two day stage, the coleorhiza is pushed through the seed coat and the primary root (the radicle) is pierced the coleorhiza but is barely visible. Also, at this two day stage, the coleoptile is just emerging from the seed coat. The 2 days post germination seedlings are then immersed in liquid nitrogen and crushed. The harvested tissue is stored at -80°C until preparation of total RNA.

The SATMON013 cDNA library is generated from apical maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) meristem founder at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F

and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Prior to tissue collection, the plant is at the 4 leaf stage. The lead at the apex of the V4 stage maize plant is referred to as the meristem founder. This apical meristem founder is cut, immediately frozen in liquid nitrogen and crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON014 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) endosperm fourteen days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the maize plant ear shoots are ready for fertilization. At this stage, the ear shoots are enclosed in a paper bag before silk emergence to withhold the pollen. The ear shoots are pollinated and 14 days after pollination, the ears are pulled out and then the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the endosperms are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON016 library is a maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) sheath library collected at the V8 developmental stage. Seeds are planted in a depth of approximately 3 cm in solid into 2-3 inch pots containing Metro growing medium. After 2-3 weeks growth, they are transplanted into 10" pots containing the same. Plants are watered daily before transplantation and approximately the times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting, at a strength of 150 ppm N. Two to three times during the life time of the plant from transplanting to flowering, a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15hr day/9hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. When the maize plants are at the V8 stage the 5th and 6th leaves from the bottom exhibit fully developed leaf blades. At the base of these leaves, the ligule is differentiated and the leaf blade is joined to the sheath. The sheath is dissected away from the base of the leaf then the sheath is frozen in liquid nitrogen and crushed. The tissue is then stored at -80°C until RNA preparation.

The SATMON017 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) embryo seventeen days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth the seeds are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr

day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of maize plant, which are ready for fertilization, are enclosed in a paper bag before silk emergence to withhold the pollen. The ear shoots are fertilized and 21 days after pollination, the ears are pulled out and the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the embryos are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON019 (Lib3054) cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) culm (stem) at the V8 developmental stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. When the maize plant is at the V8 stage, the 5th and 6th leaves from the bottom have fully developed leaf blades. The region between the nodes of the 5th and the sixth leaves from the bottom is the region of the stem that is collected. The leaves are pulled out and the sheath is

also torn away from the stem. This stem tissue is completely free of any leaf and sheath tissue. The stem tissue is then frozen in liquid nitrogen and stored at -80°C until RNA preparation.

The SATMON020 cDNA library is from a maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) Hill Type II-Initiated Callus. Petri plates containing approximately 25 ml of Type II initiation media are prepared. This medium contains N6 salts and vitamins, 3% sucrose, 2.3 g/liter proline 0.1 g/liter enzymatic casein hydrolysate, 2mg/liter 2,4 – dichloro phenoxy-acetic acid (2,4, D), 15.3 mg/liter AgNO₃ and 0.8% bacto agar and is adjusted to pH 6.0 before autoclaving. At 9-11 days after pollination, an ear with immature embryos measuring approximately 1-2 mm in length is chosen. The husks and silks are removed and then the ear is broken into halves and placed in an autoclaved solution of Clorox/TWEEN 20 sterilizing solution. Then the ear is rinsed with deionized water. Then each embryo is extracted from the kernel. Intact embryos are placed in contact with the medium, scutellar side up). Multiple embryos are plated on each plate and the plates are incubated in the dark at 25°C. Type II calluses are friable, can be subcultured with a spatula, frequently regenerate via somatic embryogenesis and are relatively undifferentiated. As seen in the microscope, the Tape II calluses show color ranging from translucent to light yellow and heterogeneity on with respect to embryoid structure as well as stage of embryoid development. Once Type II callus are formed, the calluses is transferred to type II callus maintenance medium without AgNO₃. Every 7-10 days, the callus is subcultured. About 4 weeks after embryo isolation the callus is removed from the plates and then frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

09976094, 101301

The SATMON021 cDNA library is generated from the immature maize (DK604, Dekalb Genetics, Dekalb Illinois, U.S.A.) tassel at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. As the maize plant enters the V8 stage, tassels which are 15-20 cm in length are collected and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

The SATMON022 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) ear (growing silks) at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. *Zea mays* plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W

sodium vapor lamps. Tissue is collected when the plant is in the V8 stage. At this stage, some immature ear shoots are visible. The immature ear shoots (approximately 1 cm in length) are pulled out, frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON23 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) ear (growing silk) at the V8 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. When the tissue is harvested at the V8 stage, the length of the ear that is harvested is about 10-15 cm and the silks are just exposed (approximately 1 inch). The ear along with the silks is frozen in liquid nitrogen and then the tissue is stored at -80°C until RNA preparation.

The SATMON024 cDNA library is generated from the immature maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) tassel at the V9 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a

strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. As a maize plant enters the V9 stage, the tassel is rapidly developing and a 37 cm tassel along with the glume, anthers and pollen is collected and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

The SATMON025 cDNA library is from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) Hill Type II-Regenerated Callus. Type II callus is grown in initiation media as described for SATMON020 and then the embryoids on the surface of the Type II callus are allowed to mature and germinate. The 1-2 gm fresh weight of the soft friable type callus containing numerous embryoids are transferred to 100 x 15 mm petri plates containing 25 ml of regeneration media. Regeneration media consists of Murashige and Skoog (MS) basal salts, modified White's vitamins (0.2 g/liter glycine and 0.5 g/liter myo-inositol and 0.8% bacto agar (6SMS0D)). The plates are then placed in the dark after covering with parafilm. After 1 week, the plates are moved to a lighted growth chamber with 16 hr light and 8 hr dark photoperiod. Three weeks after plating the Type II callus to 6SMS0D, the callus exhibit shoot formation. The callus and the shoots are transferred to fresh 6SMS0D plates for another 2 weeks. The callus and the shoots are then transferred to petri plates with reduced sucrose (3SMS0D). Upon distinct formation of a root and shoot, the newly developed green plants are then removed out with a spatula and frozen in liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.

09976044-104604

The SATMON026 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) juvenile/adult shift leaves at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plants are at the 8-leaf development stage. Leaves are founded sequentially around the meristem over weeks of time and the older, more juvenile leaves arise earlier and in a more basal position than the younger, more adult leaves, which are in a more apical position. In a V8 plant, some leaves which are in the middle portion of the plant exhibit characteristics of both juvenile as well as adult leaves. They exhibit a yellowing color but also exhibit, in part, a green color. These leaves are termed juvenile/adult shift leaves. The juvenile/adult shift leaves (the 4th, 5th leaves from the bottom) are cut at the base, pooled and transferred to liquid nitrogen in which they are then crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON027 cDNA library is generated from 6 day maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaves. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are

transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. *Zea mays* plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Prior to tissue collection, when the plant is at the 8-leaf stage, water is held back for six days. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical, are all cut at the base of the leaves. All the leaves exhibit significant wilting. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are then crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON028 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) roots at the V8 developmental stage that are subject to six days water stress. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the Metro 200 growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is

approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Prior to tissue collection, when the plant is at the 8-leaf stage, water is held back for six days. The root system is cut, shaken and washed to remove soil. Root tissue is then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are then crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON029 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings at the etiolated stage. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark for 4 days at approximately 70°F. Tissue is collected when the seedlings are 4 days old. By 4 days, the primary root has penetrated the coleorhiza and is about 4-5 cm and the secondary lateral roots have also made their appearance. The coleoptile has also pushed through the seed coat and is about 4-5 cm long. The seedlings are frozen in liquid nitrogen and crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON030 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) root tissue at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth, they are transplanted into 10 inch pots containing the same. Plants are watered daily before transplantation and approximately 3 times a week after transplantation. Peters 15-16-17 fertilizer is applied approximately three times per week after transplanting, at a strength of 150 ppm N. Two to three times during the life time of the plant, from transplanting to flowering, a total of approximately 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15hr day/9hr night cycles. The daytime temperature is approximately 80°F

and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 sodium vapor lamps. Tissue is collected when the maize plant is at the 4 leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is then immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON031 cDNA library is generated from the maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) leaf tissue at the V4 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is 80°F and the nighttime temperature is 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 4-leaf development stage. The third leaf from the bottom is cut at the base and immediately frozen in liquid nitrogen and crushed. The tissue is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON033 cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) embryo tissue 13 days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium.

Plants are watered daily before transplantation and three times a week after transplantation.

Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of the maize plant, which are ready for fertilization, are enclosed in a paper bag before silk emergent to withhold the pollen. The ear shoots are pollinated and 13 days after pollination, the ears are pulled out and then the kernels are plucked cut of the ears. Each kernel is then dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the embryos are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The SATMON034 cDNA library is generated from cold stressed maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) seedlings. Seeds are planted on a moist filter paper on a covered tray that is kept on at 10°C for 7 days. After 7 days, the temperature is shifted to 15°C for one day until germination of the seed. Tissue is collected once the seedlings are 1 day old. At this point, the coleorhiza has just pushed out of the seed coat and the primary root is just making its appearance. The coleoptile has not yet pushed completely through the seed coat and is also just making its appearance. These 1 day old cold stressed seedlings are frozen in liquid nitrogen and crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMON~001 (Lib36, Lib83, Lib84) cDNA library is generated from maize leaves at the V8 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3

inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant is collected at the V8 stage. The older more juvenile leaves in a basal position as well as the younger more adult leaves which are more apical are all cut at the base, pooled and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SATMONN01 cDNA library is generated from maize (B73, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) normalized immature tassels at the V6 plant development stage normalized tissue. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in a greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue from the maize plant

is collected at the V6 stage. At that stage the tassel is an immature tassel of about 2-3 cm in length. The tassels are removed and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The SATMONN04 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign, Illinois U.S.A.) normalized total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is

at the 6-leaf development stage. The older, more juvenile leaves, which are in a basal position, as well as the younger, more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The SATMONN05 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) normalized root tissue at the V6 development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is

approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The root system is cut from the mature maize plant and washed with water to free it from the soil. The tissue is immediately frozen in liquid nitrogen and the harvested tissue is then stored at -80°C until RNA preparation. The single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The SATMONN06 cDNA library is generated from maize (B73 x Mo17, Illinois Foundation Seeds, Champaign Illinois, U.S.A.) normalized total leaf tissue at the V6 plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize

plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 6-leaf development stage. The older more juvenile leaves, which are in a basal position, as well as the younger more adult leaves, which are more apical are cut at the base of the leaves. The leaves are then pooled and immediately transferred to liquid nitrogen containers in which the pooled leaves are crushed. The harvested tissue is then stored at -80°C until RNA preparation. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The CMZ029 (SATMON036) cDNA library is generated from maize (DK604, Dekalb Genetics, Dekalb, Illinois U.S.A.) endosperm 22 days after pollination. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a

strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the green house in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. After the V10 stage, the ear shoots of the maize plant, which are ready for fertilization, are enclosed in a paper bag before silk emergent to withhold the pollen. The ear shoots are pollinated and 22 days after pollination, the ears are pulled out and then the kernels are plucked out of the ears. Each kernel is then dissected into the embryo and the endosperm and the alurone layer is removed. After dissection, the endosperms are immediately frozen in liquid nitrogen and then stored at -80°C until RNA preparation.

The CMz030 (Lib143) cDNA library is generated from maize seedling tissue two days post germination. Seeds are planted on a moist filter paper on a covered tray that is kept in the dark until germination. The trays are then moved to the bench top at 15 hr daytime/9 hr nighttime cycles for 2 days post-germination. The day time temperature is 80°F and the nighttime temperature is 70°F. Tissue is collected when the seedlings are 2 days old. At this stage, the coleorrhiza has pushed through the seed coat and the primary root (the radicle) is just piercing the coleorrhiza and is barely visible. The seedlings are placed at 42°C for 1 hour. Following the heat shock treatment, the seedlings are immersed in liquid nitrogen and crushed. The harvested tissue is stored at -80°C until RNA preparation.

The CMz031 (Lib148) cDNA library is generated from maize pollen tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into

10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag to withhold pollen. Twenty-one days after pollination, prior to removing the ears, the paper bag is shaken to collect the mature pollen. The mature pollen is immediately frozen in liquid nitrogen containers and the pollen is crushed. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz033 (Lib189) cDNA library is generated from maize pooled leaf tissue. Samples are harvested from open pollinated plants. Tissue is collected from maize leaves at the anthesis stage. The leaves are collected from 10-12 plants and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz034 (Lib3060) cDNA library is generated from maize mature tissue at 40 days post pollination plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times

during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from leaves located two leaves below the ear leaf. This sample represents those genes expressed during onset and early stages of leaf senescence. The leaves are pooled and immediately transferred to liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz035 (Lib3061) cDNA library is generated from maize endosperm tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag prior to silk emergence to withhold pollen. Thirty-two days after pollination, the ears are pulled out and the kernels are removed from the cob. Each kernel is dissected into the embryo and the endosperm and the aleurone layer is removed. After dissection, the endosperms are immediately

transferred to liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz036 (Lib3062) cDNA library is generated from maize husk tissue at the 8 week old plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from 8 week old plants. The husk is separated from the ear and immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz037 (Lib3059) cDNA library is generated from maize pooled kernal at 12-15 days after pollination plant development stage. Sample were collected from field grown material. Whole kernals from hand pollinated (control pollination) are harvested as whole ears and immediately frozen on dry ice. Kernels from 10-12 ears were pooled and ground together in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz039 (Lib3066) cDNA library is generated from maize immature anther tissue at the 7 week old immature tassel stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are

transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 7 week old immature tassel stage. At this stage, prior to anthesis, the immature anthers are green and enclosed in the staminate spikelet. The developing anthers are dissected away from the 7 week old immature tassel and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz040 (Lib3067) cDNA library is generated from maize kernel tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag

before silk emergence to withhold pollen. Five to eight days after controlled pollination. The ears are pulled and the kernels removed. The kernels are immediately frozen in liquid nitrogen. The harvested kernels tissue is then stored at -80°C until RNA preparation. This sample represents gene expressed in early kernel development, during periods of cell division, amyloplast biogenesis and early carbon flow across the material to filial tissue.

The CMz041 (Lib3068) cDNA library is generated from maize pollen germinating silk tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants when the ear shoots are ready for fertilization at the silk emergence stage. The emerging silks are pollinated with an excess of pollen under controlled pollination conditions in the green house. Eighteen hours after pollination the silks are removed from the ears and immediately frozen in liquid nitrogen containers. This sample represents genes expressed in both pollen and silk tissue early in pollination. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz042 (Lib3069) cDNA library is generated from maize ear tissue excessively pollinated at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ stage plants and the ear shoots which are ready for fertilization are at the silk emergence stage. The immature ears are pollinated with an excess of pollen under controlled pollination conditions. Eighteen hours post-pollination, the ears are removed and immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz044 (Lib3075) cDNA library is generated from maize microspore tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime

temperature is approximately 80°F and the nighttime temperature is approximately 70°F.

Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from immature anthers from 7 week old tassels. The immature anthers are first dissected from the 7 week old tassel with a scalpel on a glass slide covered with water. The microspores (immature pollen) are released into the water and are recovered by centrifugation. The microspore suspension is immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz045 (Lib3076) cDNA library is generated from maize immature ear megaspore tissue. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from immature ear (megaspore) obtained from 7 week old plants. The immature ears are harvested from the 7 week old plants and are approximately 2.5 to 3 cm in length. The kernels are removed from the cob immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

09976094.104504

The CMz047 (Lib3078) cDNA library is generated from maize CO₂ treated high-exposure shoot tissue at the V10+ plant development stage. RX601 maize seeds are sterilized for 1 minute with a 10% clorox solution. The seeds are rolled in germination paper, and germinated in 0.5 mM calcium sulfate solution for two days at 30°C. The seedlings are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium at a rate of 2-3 seedlings per pot. Twenty pots are placed into a high CO₂ environment (approximately 1000 ppm CO₂). Twenty plants were grown under ambient greenhouse CO₂ (approximately 450 ppm CO₂). Plants are watered daily before transplantation and three times a week after transplantation. Peters 20-20-20 fertilizer is also lightly applied. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. At ten days post planting, the shoots from both atmosphere are frozen in liquid nitrogen and lightly ground. The roots are washed in deionized water to remove the support media and the tissue is immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz048 (Lib3079) cDNA library is generated from maize basal endosperm transfer layer tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe

is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected from V10+ maize plants. The ear shoots, which are ready for fertilization, are enclosed in a paper bag prior to silk emergence, to withhold the pollen. Kernels are harvested at 12 days post-pollination and placed on wet ice for dissection. The kernels are cross sectioned laterally, dissecting just above the pedicel region, including 1-2 mm of the lower endosperm and the basal endosperm transfer region. The pedicel and lower endosperm region containing the basal endosperm transfer layer is pooled and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz049(Lib3088) cDNA library is generated from maize immature anther tissue at the 7 week old immature tassel stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is at the 7 week old immature tassel stage. At this stage, prior to anthesis, the immature anthers are green and enclosed in the staminate spikelet. The developing anthers are

dissected away from the 7 week old immature tassel and immediately transferred to liquid nitrogen container. The harvested tissue is then stored at -80°C until RNA preparation.

The CMz050 (Lib3114) cDNA library is generated from maize silk tissue at the V10+ plant development stage. Seeds are planted at a depth of approximately 3 cm into 2-3 inch peat pots containing Metro 200 growing medium. After 2-3 weeks growth they are transplanted into 10 inch pots containing the same growing medium. Plants are watered daily before transplantation and three times a week after transplantation. Peters 15-16-17 fertilizer is applied three times per week after transplanting at a strength of 150 ppm N. Two to three times during the lifetime of the plant, from transplanting to flowering, a total of 900 mg Fe is added to each pot. Maize plants are grown in the greenhouse in 15 hr day/9 hr night cycles. The daytime temperature is approximately 80°F and the nighttime temperature is approximately 70°F. Supplemental lighting is provided by 1000 W sodium vapor lamps. Tissue is collected when the maize plant is beyond the 10-leaf development stage and the ear shoots are approximately 15-20 cm in length. The ears are pulled and silks are separated from the ears and immediately transferred to liquid nitrogen containers. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON001 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) total leaf tissue at the V4 plant development stage. Leaf tissue from 38, field grown V4 stage plants is harvested from the 4th node. Leaf tissue is removed from the plants and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

09976064-1015001
The SOYMON002 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue at the V4 plant development stage. Root tissue from 76, field grown V4 stage plants is harvested. The root systems is cut from the soybean plant and washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON003 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling hypocotyl axis tissue harvested 2 day post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 2 days after the start of imbibition. The 2 days after imbibition samples are separated into 3 collections after removal of any adhering seed coat. At the 2 day stage, the hypocotyl axis is emerging from the soil. A few seedlings have cracked the soil surface and exhibited slight greening of the exposed cotyledons. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON004 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling cotyledon tissue harvested 2 day post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even

moisture conditions. Tissue is collected 2 days after the start of imbibition. The 2 days after imbibition samples are separated into 3 collections after removal of any adhering seed coat. At the 2 day stage, the hypocotyl axis is emerging from the soil. A few seedlings have cracked the soil surface and exhibited slight greening of the exposed cotyledons. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON005 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling hypocotyl axis tissue harvested 6 hour post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 6 hours after the start of imbibition. The 6 hours after imbibition samples are separated into 3 collections after removal of any adhering seed coat. The 6 hours after imbibition sample is collected over the course of approximately 2 hours starting at 6 hours post imbibition. At the 6 hours after imbibition stage, not all cotyledons have become fully hydrated and germination, or radicle protrusion, has not occurred. The seedlings are washed in water to remove soil, hypocotyl axis harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON006 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling cotyledons tissue harvest 6 hour post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. Trays are placed in an environmental chamber and grown at

12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Tissue is collected 6 hours after imbibition. The 6 hours after imbibition samples are separated into 3 collections after removal of any adhering seed coat. The 6 hours after imbibition sample is collected over the course of approximately 2 hours starting at 6 hours post-imbibition. At the 6 hours after imbibition, not all cotyledons have become fully hydrated and germination or radicle protrusion, have not occurred. The seedlings are washed in water to remove soil, cotyledon harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON007 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 and 35 days post-flowering. Seed pods from field grown plants are harvested 25 and 35 days after flowering and the seeds extracted from the pods. Approximately 4.4g and 19.3g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON008 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested from 25 and 35 days post-flowering plants. Total leaf tissue is harvested from field grown plants. Approximately 19g and 29g of leaves are harvested from the fourth node of the plant 25 and 35 days post-flowering and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON009 cDNA library is generated from soybean cutlivar C1944 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) pod and seed tissue harvested 15 days

post-flowering. Pods from field grown plants are harvested 15 days post-flowering.

Approximately 3g of pod tissue is harvested and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON010 cDNA library is generated from soybean cultivar C1944 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) seed tissue harvested 40 days post-flowering. Pods from field grown plants are harvested 40 days post-flowering. Pods and seeds are separated, approximately 19g of seed tissue is harvested and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON011 cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON012 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue. Leaves from field grown plants are harvested from the fourth node 15 days post-flowering. Approximately 12g of leaves are harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON013 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root and nodule tissue. Approximately, 28g of root tissue from field grown plants is harvested 15 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON014 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 and 35 days after flowering. Seed pods from field grown plants are harvested 15 days after flowering and the seeds extracted from the pods. Approximately 5g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON015 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 45 and 55 days post-flowering. Seed pods from field grown plants are harvested 45 and 55 days after flowering and the seeds extracted from the pods. Approximately 19g and 31g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON016 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue. Approximately, 61g and 38g of root tissue from field grown plants is harvested 25 and 35 days post- flowering is harvested. The root system is cut from the soybean plant and washed with water to free it from the soil. The tissue is placed in 14ml polystyrene tubes and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON017 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue. Approximately 28g of root tissue from field grown plants is harvested 45 and 55 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON018 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested 45 and 55 days post-flowering. Leaves from field grown plants are harvested 45 and 55 days after flowering from the fourth node. Approximately 27g and 33g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON019 cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) root tissue. Roots are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 50g and 56g of roots are harvested from each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON020 cDNA is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 65 and 75 days post-flowering. Seed pods from field grown plants are harvested 45 and 55 days after flowering and the seeds extracted from the pods. Approximately 14g and 31g of seeds are harvested from the respective

seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON021 cDNA library is generated from Soybean Cyst Nematode-resistant soybean cultivar Hartwig (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) root tissue. Plants are grown in tissue culture at room temperature. At approximately 6 weeks post-germination, the plants are exposed to sterilized Soybean Cyst Nematode eggs. Infection is then allowed to progress for 10 days. After the 10 day infection process, the tissue is harvested. Agar from the culture medium and nematodes are removed and the root tissue is immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON022 (Lib3030) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) partially opened flower tissue. Partially to fully opened flower tissue is harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. A total of 3g of flower tissue is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON023 cDNA library is generated from soybean genotype BW211S Null (Tohoku University, Morioka, Japan) seed tissue harvested 15 and 40 days post-flowering. Seed pods from field grown plants are harvested 15 and 40 days post-flowering and the seeds extracted from the pods. Approximately 0.7g and 14.2g of seeds are harvested from the respective seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON024 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) internode-2 tissue harvested 18 days post-imbibition. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium. The plants are grown in a greenhouse for 18 days after the start of imbibition at ambient temperature. Soil is checked and watered daily to maintain even moisture conditions. Stem tissue is harvested 18 days after the start of imbibition. The samples are divided into hypocotyl and internodes 1 through 5. The fifth internode contains some leaf bud material. Approximately 3 g of each sample is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON025 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) leaf tissue harvested 65 days post-flowering. Leaves are harvested from the fourth node of field grown plants 65 days post-flowering. Approximately 18.4g of leaf tissue is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

SOYMON026 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) root tissue harvested 65 and 75 days post-flowering. Approximately 27g and 40g of root tissue from field grown plants is harvested 65 and 75 days post-flowering. The root system is cut from the soybean plant, washed with water to free it from the soil and immediately frozen in dry-ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON027 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed tissue harvested 25 days post-flowering. Seed pods from field grown plants are harvested 25 days post-flowering and the seeds

extracted from the pods. Approximately 17g of seeds are harvested from the seed pods and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON028 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought-stressed root tissue. The plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of development, water is withheld from half of the plant collection (drought stressed population). After 3 days, half of the plants from the drought stressed condition and half of the plants from the control population are harvested. After another 3 days (6 days post drought induction) the remaining plants are harvested. A total of 27g and 40g of root tissue is harvested and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON029 cDNA library is generated from Soybean Cyst Nematode-resistant soybean cultivar PI07354 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) root tissue. Late fall to early winter greenhouse grown plants are exposed to Soybean Cyst Nematode eggs. At 10 days post-infection, the plants are uprooted, rinsed briefly and the roots frozen in liquid nitrogen. Approximately 20 grams of root tissue is harvested from the infected plants. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON030 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) flower bud tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The

daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C.

Soil is checked and watered daily to maintain even moisture conditions. Flower buds are removed from the plant at the pedicel. A total of 100mg of flower buds are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON031 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) carpel and stamen tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Flower buds are removed from the plant at the pedicel. Flowers are dissected to separate petals, sepals and reproductive structures (carpels and stamens). A total of 300mg of carpel and stamen tissue are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON032 cDNA library is prepared from the Asgrow cultivar A4922 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) rehydrated dry soybean seed meristem tissue. Surface sterilized seeds are germinated in liquid media for 24 hours. The seed axis is then excised from the barely germinating seed, placed on tissue culture media and incubated overnight at 20°C in the dark. The supportive tissue is removed from the explant prior to harvest. Approximately 570mg of tissue is harvested and frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

0906064-104604

The SOYMON033 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) heat-shocked seedling tissue without cotyledons. Seeds are imbibed and germinated in vermiculite for 2 days under constant illumination. After 48 hours, the seedlings are transferred to an incubator set at 40°C under constant illumination. After 30, 60 and 180 minutes seedlings are harvested and dissected. A portion of the seedling consisting of the root, hypocotyl and apical hook is frozen in liquid nitrogen and stored at -80°C. The seedlings after 2 days of imbibition are beginning to emerge from the vermiculite surface. The apical hooks are dark green in appearance. Total RNA and poly A⁺ RNA is prepared from equal amounts of pooled tissue.

The SOYMON034 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) cold-shocked seedling tissue without cotyledons. Seeds are imbibed and germinated in vermiculite for 2 days under constant illumination. After 48 hours, the seedlings are transferred to a cold room set at 5°C under constant illumination. After 30, 60 and 180 minutes seedlings are harvested and dissected. A portion of the seedling consisting of the root, hypocotyl and apical hook is frozen in liquid nitrogen and stored at -80°C. The seedlings after 2 days of imbibition are beginning to emerge from the vermiculite surface. The apical hooks are dark green in appearance.

The SOYMON035 cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seed coat tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Seeds are harvested from mid to nearly

full maturation (seed coats are not yellowing). The entire embryo proper is removed from the seed coat sample and the seed coat tissue are harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON036 cDNA library is generated from soybean cultivars PI171451, PI227687 and PI229358 (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) insect challenged leaves. Plants from each of the three cultivars are grown in screenhouse conditions. The screenhouse is divided in half and one half of the screenhouse is infested with soybean looper and the other half infested with velvetbean caterpillar. A single leaf is taken from each of the representative plants at 3 different time points, 11 days after infestation, 2 weeks after infestation and 5 weeks after infestation and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation. Total RNA and poly A+ RNA is isolated from pooled tissue consisting of equal quantities of all 18 samples (3 genotypes X 3 sample times X 2 insect genotypes).

The SOYMON037 cDNA library is generated from soybean cultivar A3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) etiolated axis and radical tissue. Seeds are planted in moist vermiculite, wrapped and kept at room temperature in complete darkness until harvest. Etiolated axis and hypocotyl tissue is harvested at 2, 3 and 4 days post-planting. A total of 1 gram of each tissue type is harvested at 2, 3 and 4 days after planting and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The SOYMON038 cDNA library is generated from soybean variety Asgrow A3237 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) rehydrated dry seeds. Explants are prepared for transformation after germination of surface-sterilized seeds on solid tissue media. After 6 days, at 28°C and 18 hours of light per day, the germinated seeds are cold shocked at 4°C for 24

hours. Meristemic tissue and part of the hypocotyl is removed and cotyledon excised. The prepared explant is then wounded for *Agrobacterium* infection. The 2 grams of harvested tissue is frozen in liquid nitrogen and stored at -80°C until RNA preparation.

The Soy51 (LIB3027) cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The Soy52 (LIB3028) cDNA library is generated from normalized flower DNA. Single stranded DNA representing approximately 1×10^6 colony forming units of SOYMON022 harvested tissue is used as the starting material for normalization. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet.

The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

The Soy53 (LIB3039) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) seedling shoot apical meristem tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Apical tissue is harvested from seedling shoot meristem tissue, 7-8 days after the start of imbibition. The apex of each seedling is dissected to include the fifth node to the apical meristem. The fifth node corresponds to the third trifoliolate leaf in the very early stages of development. Stipules completely envelop the leaf primordia at this time. A total of 200mg of apical tissue is harvested and immediately frozen in liquid nitrogen. The harvested tissue is then stored at -80°C until RNA preparation.

The Soy54 (LIB3040) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) heart to torpedo stage embryo tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Seeds are collected and embryos removed from surrounding endosperm and maternal tissues. Embryos from globular to young torpedo stages (by corresponding analogy to *Arabidopsis*) are collected with a bias towards the middle of this spectrum. Embryos which are beginning to show

asymmetric development of cotyledons are considered the upper developmental boundary for the collection and are excluded. A total of 12 mg embryo tissue is frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

Soy55 (LIB3049) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) young seed tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Seeds are collected from very young pods (5 to 15 days after flowering). A total of 100mg of seeds are harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

Soy56 (LIB3029) cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are not converted to double stranded form and represent a non-normalized seed pool for comparison to Soy51 cDNA libraries.

TheSoy58 (LIB3050) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed root tissue subtracted from control root tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days root tissue from both drought stressed and control (watered regularly) plants are collected and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. For subtraction, target cDNA is made from the drought stressed tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 l 2X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.).

The Soy59 (LIB3051) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) endosperm tissue. Seeds are germinated on

paper towels under laboratory ambient light conditions. At 8, 10 and 14 hours after imbibition, the seed coats are harvested. The endosperm consists of a very thin layer of tissue affixed to the inside of the seed coat. The seed coat and endosperm are frozen immediately after harvest in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

The Soy60 (LIB3072) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed seed plus pod subtracted from control seed plus pod tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 26°C and the nighttime temperature 21°C and 70% relative humidity. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days seeds and pods from both drought stressed and control (watered regularly) plants are collected from the fifth and sixth node and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. For subtraction, target cDNA is made from the drought stressed tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 12X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads

oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.).

The Soy61 (LIB3073) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. After 18hours, 24hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. For subtraction, target cDNA is made from the jasmonic acid treated tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 12X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed

from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.). For this library's construction, the eighth fraction of the cDNA size fractionation step was used for ligation.

The Soy62 (LIB3074) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. After 18 hours, 24 hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. For subtraction, target cDNA is made from the jasmonic acid treated tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using

Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 12X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.). For this library's construction, the ninth fraction of the cDNA size fractionation step was used for ligation.

The Soy65 (LIB3107) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought-stressed abscission zone tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature 24°C. Soil is checked and watered daily to maintain even moisture conditions. Plants are irrigated with 15-16-17 Peter's Mix. At the R3 stage of development, drought is imposed by withholding water. At 3, 4, 5 and 6 days, tissue is harvested and wilting is not obvious until the fourth day. Abscission layers from reproductive organs are harvested by cutting less than one millimeter proximal and distal to the layer and immediately frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

The Soy66 (LIB3109) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) non-drought stressed abscission zone tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr

nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Plants are irrigated with 15-16-17 Peter's Mix. At 3, 4, 5 and 6 days, control abscission layer tissue is harvested. Abscission layers from reproductive organs are harvested by cutting less than one millimeter proximal and distal to the layer and immediately frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation.

Soy67 (LIB3065) cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. Captured hybrids are eluted with water.

Soy68 (LIB3052) cDNA library is prepared from equal amounts tissue harvested from SOYMON007, SOYMON015 and SOYMON020 prepared tissue. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280

streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. Captured hybrids are eluted with water.

Soy69 (LIB3053) cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) normalized leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation. Single stranded and double stranded DNA representing approximately 1×10^6 colony forming units are isolated using standard protocols. RNA, complementary to the single stranded DNA, is synthesized using the double stranded DNA as a template. Biotinylated dATP is incorporated into the RNA during the synthesis reaction. The single stranded DNA is mixed with the biotinylated RNA in a 1:10 molar ratio) and allowed to hybridize. DNA-RNA hybrids are captured on Dynabeads M280 streptavidin (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The dynabeads with captured hybrids are collected with a magnet. The non-hybridized single stranded molecules remaining after hybrid capture are converted to double stranded form and represent the primary normalized library.

Soy70 (LIB3055) cDNA library is generated from soybean cultivars Cristalina (USDA Soybean Germplasm Collection, Urbana, Illinois U.S.A.) and FT108 (Monsoy, Brazil) (tropical germ plasma) leaf tissue. Leaves are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and

the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 30g of leaves are harvested from the 4th node of each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

Soy71 (LIB3056) cDNA library is generated from soybean cultivars Cristalina and FT108 (tropical germ plasma) root tissue. Roots are harvested from plants grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 29°C and the nighttime temperature approximately 24°C. Soil is checked and watered daily to maintain even moisture conditions. Approximately 50g and 56g of roots are harvested from each of the Cristalina and FT108 cultivars and immediately frozen in dry ice. The harvested tissue is then stored at -80°C until RNA preparation.

Soy72 (LIB3093) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed leaf control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 26°C and the nighttime temperature 21°C and 70% relative humidity. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days seeds and pods from both drought stressed and control (watered regularly) plants are collected from the fifth and sixth node and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. For subtraction, target cDNA is made from the drought stressed tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to

Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 1 2X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.).

Soy73 (LIB3093) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) drought stressed leaf subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in an environmental chamber under 12hr daytime/12hr nighttime cycles. The daytime temperature is approximately 26°C and the nighttime temperature 21°C and 70% relative humidity. Soil is checked and watered daily to maintain even moisture conditions. At the R3 stage of the plant drought is induced by withholding water. After 3 and 6 days seeds and pods from both drought stressed and control (watered regularly) plants are collected from the fifth and sixth node and frozen in dry-ice. The harvested tissue is stored at -80°C until RNA preparation. For subtraction, target cDNA is made from the drought stressed tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA

is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 1 2X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.).

The Soy76 (Lib3106) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid and arachidonic treated seedling subtracted from control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. Arachidonic treated seedlings are sprayed with 1m/ml arachidonic acid in 0.1% Tween-20. After 18hours, 24hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA from the arachidonic treated seedlings is isolated separately. For subtraction, target cDNA is made from the jasmonic acid treated tissue total RNA using the SMART cDNA

synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 1 2X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.). Fraction 10 of the size fractionated cDNA is ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.) in order to capture some of the smaller transcripts characteristic of antifungal proteins.

Soy77 (LIB3108) cDNA library is generated from soybean cultivar Asgrow 3244 (Asgrow Seed Company, Des Moines, Iowa U.S.A.) jasmonic acid control tissue. Seeds are planted at a depth of approximately 2cm into 2-3 inch peat pots containing Metromix 350 medium and the plants are grown in a greenhouse. The daytime temperature is approximately 29.4°C and the nighttime temperature 20°C. Soil is checked and watered daily to maintain even moisture conditions. At 9 days post planting, the plantlets are sprayed with either control buffer of 0.1% Tween-20 or jasmonic acid (Sigma J-2500, Sigma, St. Louis, Missouri U.S.A.) at 1 mg/ml in 0.1% Tween-20. Plants are sprayed until runoff and the soil and the stem is soaked with the spraying solution. At 18 hours post application of jasmonic acid, the soybean plantlets appear growth retarded. Arachidonic treated seedlings are sprayed with 1m/ml arachidonic acid

in 0.1% Tween-20. After 18 hours, 24 hours and 48 hours post treatment, the cotyledons are removed and the remaining leaf and stem tissue above the soil is harvested and frozen in liquid nitrogen. The harvested tissue is stored at -80°C until RNA preparation. To make RNA, the three sample timepoints were combined and ground. The RNA from the arachidonic treated seedlings is isolated separately. For subtraction, target cDNA is made from the jasmonic acid treated tissue total RNA using the SMART cDNA synthesis system from Clontech (Clontech Laboratories, Palo Alto, California U.S.A.). Driver first strand cDNA is covalently linked to Dynabeads following a protocol similar to that described in the Dynal literature (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.). The target cDNA is then heat denatured and the second strand trapped using Dynabeads oligo-dT. The target second strand cDNA is then hybridized to the driver cDNA in 400 12X SSPE for two rounds of hybridization at 65°C and 20 hours. After each hybridization, the hybridization solution is removed from the system and the hybridized target cDNA removed from the driver by heat denaturation in water. After hybridization, the remaining cDNA is trapped with Dynabeads oligo-dT. The trapped cDNA is then amplified as in previous PCR based libraries and the resulting cDNA ligated into the pSPORT vector (Invitrogen, Carlsbad California U.S.A.). Fraction 10 of the size fractionated cDNA is ligated into the pSPORT vector in order to capture some of the smaller transcripts characteristic of antifungal proteins.

The stored RNA is purified using Trizol reagent from Life Technologies (Gibco BRL, Life Technologies, Gaithersburg, Maryland U.S.A.), essentially as recommended by the manufacturer. Poly A⁺ RNA (mRNA) is purified using magnetic oligo dT beads essentially as recommended by the manufacturer (Dynabeads, Dynal Corporation, Lake Success, New York U.S.A.).

Construction of plant cDNA libraries is well-known in the art and a number of cloning strategies exist. A number of cDNA library construction kits are commercially available. The Superscript™ Plasmid System for cDNA synthesis and Plasmid Cloning (Gibco BRL, Life Technologies, Gaithersburg, Maryland U.S.A.) is used, following the conditions suggested by the manufacturer.

Normalized libraries are made using essentially the Soares procedure (Soares *et al.*, *Proc. Natl. Acad. Sci. (U.S.A.)* 91:9228-9232 (1994), the entirety of which is herein incorporated by reference). This approach is designed to reduce the initial 10,000-fold variation in individual cDNA frequencies to achieve abundances within one order of magnitude while maintaining the overall sequence complexity of the library. In the normalization process, the prevalence of high-abundance cDNA clones decreases dramatically, clones with mid-level abundance are relatively unaffected and clones for rare transcripts are effectively increased in abundance.

Example 2

The cDNA libraries are plated on LB agar containing the appropriate antibiotics for selection and incubated at 37° for a sufficient time to allow the growth of individual colonies. Single colonies are individually placed in each well of a 96-well microtiter plates containing LB liquid including the selective antibiotics. The plates are incubated overnight at approximately 37°C with gentle shaking to promote growth of the cultures. The plasmid DNA is isolated from each clone using Qiaprep plasmid isolation kits, using the conditions recommended by the manufacturer (Qiagen Inc., Santa Clara, California U.S.A.).

Template plasmid DNA clones are used for subsequent sequencing. For sequencing, the ABI PRISM dRhodamine Terminator Cycle Sequencing Ready Reaction Kit with AmpliTaq® DNA Polymerase, FS, is used (PE Applied Biosystems, Foster City, California U.S.A.).

Example 3

Nucleic acid sequences that encode for the following proteins: adenine phosphoribosyl transferase, β glucosidase and isopentyltransferase are identified from the Monsanto EST PhytoSeq database using TBLASTN (default values)(TBLASTN compares a protein query against the six reading frames of a nucleic acid sequence). Matches found with BLAST P values equal or less than 0.001 (probability) or BLAST Score of equal or greater than 90 are classified as hits. If the program used to determine the hit is HMMSW then the score refers to HMMSW score.

In addition, the GenBank database is searched with BLASTN and BLASTX (default values) using ESTs as queries. EST that pass the hit probability threshold of $10e^{-8}$ for the following enzymes are combined with the hits generated by using TBLASTN (described above) and classified by enzyme (see Table A below).

A cluster refers to a set of overlapping clones in the PhytoSeq database. Such an overlapping relationship among clones is designated as a "cluster" when BLAST scores from pairwise sequence comparisons of the member clones meets a predetermined minimum value or product score of 50 or more (Product Score = (BLAST SCORE x Percentage Identity)/(5 x minimum [length (Seq1), length (Seq2)]))

Since clusters are formed on the basis of single-linkage relationships, it is possible for two non-overlapping clones to be members of the same cluster if, for instance, they both overlap

a third clone with at least the predetermined minimum BLAST score (stringency). A cluster ID is arbitrarily assigned to all of those clones which belong to the same cluster at a given stringency and a particular clone will belong to only one cluster at a given stringency. If a cluster contains only a single clone (a "singleton"), then the cluster ID number will be negative, with an absolute value equal to the clone ID number of its single member. Clones grouped in a cluster in most cases represent a contiguous sequence.

TABLE A*

MAIZE ADENINE PHOSPHORIBOSYL TRANSFERASE (EC 2.4.2.7)								
Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	% Ident
1	-700193568	700193568H1	SATMON014	g726304	BLASTN	490	1e-32	73
2	-700432807	700432807H1	SATMONN01	g16164	BLASTX	212	1e-26	68
3	-700475820	700475820H1	SATMON025	g726305	BLASTX	87	1e-11	84
4	-700552966	700552966H1	SATMON022	g726304	BLASTN	927	1e-68	81
5	-L30612612	LIB3061-015-Q1-K1-H2	LIB3061	g726304	BLASTN	447	1e-26	73
6	-L30682155	LIB3068-004-Q1-K1-B3	LIB3068	g726304	BLASTN	320	1e-27	77
7	-L30691613	LIB3069-005-Q1-K1-D1	LIB3069	g726304	BLASTN	478	1e-28	76
8	-L30784520	LIB3078-039-Q1-K1-D9	LIB3078	g726304	BLASTN	374	1e-32	74
9	-L831334	LIB83-003-Q1-E1-F6	LIB83	g1402893	BLASTN	461	1e-27	66
10	10045	LIB3067-006-Q1-K1-H12	LIB3067	g1321681	BLASTX	241	1e-44	71
11	10045	700338620H1	SATMON020	g1321681	BLASTX	173	1e-29	62
12	10045	700335677H1	SATMON019	g1321681	BLASTX	86	1e-19	52
13	5380	LIB3061-047-Q1-K1-B4	LIB3061	g726304	BLASTN	1205	1e-92	82
14	5380	700082054H1	SATMON011	g726304	BLASTN	1032	1e-77	82
15	5380	700242515H1	SATMON010	g726304	BLASTN	1008	1e-75	83
16	5380	700339222H1	SATMON020	g726304	BLASTN	842	1e-74	82
17	5380	700027757H1	SATMON003	g726304	BLASTN	939	1e-69	83
18	5380	700029386H1	SATMON003	g726304	BLASTN	900	1e-66	82
19	5380	700241615H1	SATMON010	g726304	BLASTN	894	1e-65	81
20	5380	700172169H1	SATMON013	g726304	BLASTN	724	1e-51	80
21	5380	700045315H1	SATMON004	g726304	BLASTN	655	1e-45	83
22	5380	700018155H1	SATMON001	g726304	BLASTN	614	1e-42	85
23	5380	700157175H1	SATMON012	g726304	BLASTN	618	1e-42	83
24	5380	700335263H1	SATMON019	g726304	BLASTN	296	1e-36	82
25	5380	700022056H1	SATMON001	g726305	BLASTX	147	1e-13	93
26	5380	700196739H1	SATMON014	g726305	BLASTX	89	1e-10	93
27	6937	LIB189-003-Q1-E1-F5	LIB189	g726304	BLASTN	947	1e-70	80
28	6937	LIB3059-015-Q1-K1-A2	LIB3059	g726304	BLASTN	905	1e-66	82
29	6937	LIB143-061-Q1-E1-B3	LIB143	g726304	BLASTN	779	1e-55	80
30	6937	LIB3067-052-Q1-K1-A1	LIB3067	g726304	BLASTN	532	1e-54	79
31	6937	700334619H1	SATMON019	g726304	BLASTN	711	1e-50	81
32	6937	700219612H1	SATMON011	g726304	BLASTN	588	1e-40	79
33	6937	700405177H1	SATMON028	g726304	BLASTN	589	1e-40	81
34	6937	700236956H1	SATMON010	g726304	BLASTN	573	1e-39	81
35	6937	700238553H1	SATMON010	g726304	BLASTN	387	1e-37	81
36	6937	700104336H1	SATMON010	g726304	BLASTN	537	1e-36	76
37	6937	LIB3068-059-	LIB3068	g726304	BLASTN	385	1e-33	73

		Q1-K1-H7						
38	6937	700238576H1	SATMON010	g726304	BLASTN	407	1e-23	69
39	6937	700142447H1	SATMON012	g16164	BLASTX	135	1e-14	79
40	6937	700204679H1	SATMON003	g726304	BLASTN	185	1e-13	76

MAIZE β GLUCOSIDASE (EC 3.2.1.21)

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
41	-700019404	700019404H1	SATMON001	g1206012	BLASTN	587	1e-40	85
42	-700051621	700051621H1	SATMON003	g1206012	BLASTN	417	1e-55	76
43	-700072125	700072125H1	SATMON007	g1518673	BLASTN	320	1e-16	93
44	-700073309	700073309H1	SATMON007	g21953	BLASTX	97	1e-21	50
45	-700077116	700077116H1	SATMON007	g1518673	BLASTN	297	1e-14	90
46	-700084705	700084705H1	SATMON011	g1206012	BLASTN	235	1e-9	100
47	-700085269	700085269H1	SATMON011	g1143864	BLASTX	151	1e-16	53
48	-700088245	700088245H1	SATMON011	g435312	BLASTN	537	1e-59	75
49	-700094593	700094593H1	SATMON008	g1399389	BLASTN	197	1e-14	85
50	-700104334	700104334H1	SATMON010	g1399389	BLASTN	760	1e-79	96
51	-700160044	700160044H1	SATMON012	g804656	BLASTX	252	1e-37	79
52	-700168880	700168880H1	SATMON013	g435312	BLASTN	703	1e-49	81
53	-700207934	700207934H1	SATMON016	g1155255	BLASTX	172	1e-16	54
54	-700208416	700208416H1	SATMON016	g1518674	BLASTN	459	1e-36	96
55	-700220501	700220501H1	SATMON011	g1399389	BLASTN	598	1e-40	81
56	-700221075	700221075H1	SATMON011	g1143863	BLASTN	640	1e-44	75
57	-700235295	700235295H1	SATMON010	g1399389	BLASTN	1166	1e-91	93
58	-700258664	700258664H1	SATMON017	g804656	BLASTX	195	1e-28	66
59	-700265357	700265357H1	SATMON017	g1143863	BLASTN	367	1e-39	80
60	-700338753	700338753H1	SATMON020	g804655	BLASTN	955	1e-70	82
61	-700343160	700343160H1	SATMON021	g1143863	BLASTN	714	1e-50	82
62	-700352084	700352084H1	SATMON023	g1518673	BLASTN	796	1e-59	90
63	-700353902	700353902H1	SATMON024	g804656	BLASTX	238	1e-25	60
64	-700356246	700356246H1	SATMON024	g1143864	BLASTX	237	1e-26	66
65	-700356858	700356858H1	SATMON024	g804656	BLASTX	101	1e-23	54
66	-700444014	700444014H1	SATMON027	g1399389	BLASTN	426	1e-24	82
67	-700468671	700468671H1	SATMON025	g1155255	BLASTX	63	1e-10	47
68	-700468683	700468683H1	SATMON025	g804655	BLASTN	360	1e-44	81
69	-700468738	700468738H1	SATMON025	g804655	BLASTN	301	1e-47	87
70	-700469144	700469144H1	SATMON025	g1399389	BLASTN	292	1e-45	88
71	-700471979	700471979H1	SATMON025	g804656	BLASTX	172	1e-16	76
72	-700472168	700472168H1	SATMON025	g804656	BLASTX	117	1e-23	66
73	-700477783	700477783H1	SATMON025	g804655	BLASTN	341	1e-59	88
74	-700548872	700548872H1	SATMON022	g804656	BLASTX	234	1e-25	70
75	-700573216	700573216H1	SATMON030	g1399389	BLASTN	472	1e-46	90
76	-700619394	700619394H1	SATMON034	g435312	BLASTN	354	1e-33	84
77	-700621680	700621680H1	SATMON034	g21953	BLASTX	90	1e-22	61
78	-700623741	700623741H1	SATMON034	g1399390	BLASTX	152	1e-13	100
79	-700624575	700624575H1	SATMON034	g804655	BLASTN	345	1e-30	77
80	-701164553	701164553H1	SATMONN04	g1518673	BLASTN	329	1e-19	88
81	-701165120	701165120H1	SATMONN04	g1206012	BLASTN	597	1e-42	84
82	-L1431868	LIB143-029-Q1-E1-H4	LIB143	g804656	BLASTX	308	1e-51	76
83	-L1435738	LIB143-047-Q1-E1-C2	LIB143	g804656	BLASTX	123	1e-25	63

84	-L1486423	LIB148-051-Q1-E1-A8	LIB148	g1518673	BLASTN	466	1e-44	81
85	-L1892203	LIB189-005-Q1-E1-G3	LIB189	g757740	BLASTX	152	1e-28	50
86	-L1893440	LIB189-023-Q1-E1-E2	LIB189	g21953	BLASTX	129	1e-35	44
87	-L30624187	LIB3062-035-Q1-K1-G11	LIB3062	g435312	BLASTN	397	1e-22	67
88	-L30625219	LIB3062-020-Q1-K1-A12	LIB3062	g1143863	BLASTN	221	1e-12	74
89	-L30626353	LIB3062-024-Q1-K1-E2	LIB3062	g1206012	BLASTN	317	1e-30	74
90	-L30626596	LIB3062-038-Q1-K1-A12	LIB3062	g142586	BLASTX	220	1e-39	56
91	-L30665817	LIB3066-006-Q1-K1-B12	LIB3066	g1143864	BLASTX	182	1e-34	84
92	-L30676013	LIB3067-057-Q1-K1-A5	LIB3067	g1143863	BLASTN	663	1e-58	77
93	-L30692578	LIB3069-019-Q1-K1-E5	LIB3069	g1206013	BLASTX	138	1e-26	36
94	-L30692596	LIB3069-019-Q1-K1-A8	LIB3069	g799376	BLASTN	246	1e-9	51
95	-L30694297	LIB3069-051-Q1-K1-C1	LIB3069	g1143864	BLASTX	133	1e-36	54
96	-L30784416	LIB3078-039-Q1-K1-H3	LIB3078	g1206012	BLASTN	630	1e-102	81
97	10283	700356224H1	SATMON024	g804655	BLASTN	436	1e-25	68
98	10283	700354663H1	SATMON024	g804656	BLASTX	186	1e-18	64
99	10343	LIB3062-041-Q1-K1-B1	LIB3062	g1143863	BLASTN	860	1e-81	74
100	10343	700212710H1	SATMON016	g1143863	BLASTN	805	1e-63	79
101	10343	700023123H1	SATMON003	g1143863	BLASTN	834	1e-60	80
102	10343	700168364H1	SATMON013	g1143863	BLASTN	816	1e-59	80
103	10343	700281856H2	SATMON021	g1143863	BLASTN	615	1e-42	73
104	10343	700170973H1	SATMON013	g1143863	BLASTN	587	1e-40	75
105	10343	700222918H1	SATMON011	g1143863	BLASTN	529	1e-35	65
106	10343	700623415H1	SATMON034	g1143864	BLASTX	215	1e-22	52
107	10343	700262090H1	SATMON017	g1143863	BLASTN	165	1e-9	76
108	10564	700572950H1	SATMON030	g1206012	BLASTN	491	1e-85	85
109	10564	700573795H1	SATMON030	g1206012	BLASTN	881	1e-74	85
110	10564	700157129H1	SATMON012	g1206012	BLASTN	523	1e-63	86
111	10712	700073072H1	SATMON007	g1206012	BLASTN	469	1e-59	86
112	10712	700072996H1	SATMON007	g1206012	BLASTN	454	1e-58	85
113	10712	700076579H1	SATMON007	g435312	BLASTN	315	1e-41	80
114	10712	700075075H1	SATMON007	g435312	BLASTN	315	1e-34	80
115	10712	700155128H1	SATMON007	g1399390	BLASTX	136	1e-14	78
116	11895	700169369H1	SATMON013	g1143863	BLASTN	744	1e-53	79
117	11895	700622210H1	SATMON034	g1143863	BLASTN	516	1e-50	79
118	11895	700020586H1	SATMON001	g1143863	BLASTN	646	1e-45	79
119	12484	700473715H1	SATMON025	g804655	BLASTN	474	1e-67	84
120	12484	700474014H1	SATMON025	g804655	BLASTN	338	1e-54	86
121	13406	700202816H1	SATMON003	g804656	BLASTX	223	1e-23	50
122	13553	700345013H1	SATMON021	g1143864	BLASTX	150	1e-13	96
123	13553	700346887H1	SATMON021	g1143864	BLASTX	114	1e-8	95

124	14210	700106119H1	SATMON010	g1206012	BLASTN	936	1e-78	83
125	14210	700236969H1	SATMON010	g435312	BLASTN	945	1e-69	84
126	14210	700569967H1	SATMON030	g1399389	BLASTN	609	1e-41	80
127	14713	LIB3066-054-Q1-K1-H11	LIB3066	g1769814	BLASTX	183	1e-36	63
128	14713	LIB3066-053-Q1-K1-H12	LIB3066	g21955	BLASTX	162	1e-34	71
129	14713	700103716H1	SATMON010	g1769814	BLASTX	111	1e-12	63
130	14713	700096365H1	SATMON008	g21955	BLASTX	135	1e-11	65
131	15366	LIB143-060-Q1-E1-B6	LIB143	g804655	BLASTN	605	1e-85	82
132	15366	700469301H1	SATMON025	g804655	BLASTN	524	1e-55	87
133	15366	700573405H2	SATMON030	g804655	BLASTN	388	1e-45	84
134	15366	700473205H1	SATMON025	g804655	BLASTN	374	1e-27	79
135	15366	700263901H1	SATMON017	g804656	BLASTX	69	1e-24	86
136	15944	LIB3062-042-Q1-K1-D8	LIB3062	g1206012	BLASTN	928	1e-124	83
137	15944	LIB3062-038-Q1-K1-F5	LIB3062	g1206012	BLASTN	1103	1e-121	81
138	15944	LIB3062-010-Q1-K1-F8	LIB3062	g1206012	BLASTN	826	1e-116	83
139	15944	LIB3062-039-Q1-K1-H1	LIB3062	g1206012	BLASTN	842	1e-97	84
140	15944	LIB3062-002-Q1-K2-D6	LIB3062	g1206012	BLASTN	940	1e-97	81
141	15944	700104654H1	SATMON010	g435312	BLASTN	815	1e-89	85
142	15944	LIB3062-015-Q1-K1-H3	LIB3062	g1206012	BLASTN	1060	1e-79	76
143	15944	LIB3062-029-Q1-K1-D9	LIB3062	g1206012	BLASTN	1008	1e-75	81
144	15944	LIB3062-027-Q1-K1-D11	LIB3062	g1206012	BLASTN	848	1e-72	82
145	15944	700221092H1	SATMON011	g1206012	BLASTN	643	1e-71	88
146	15944	700020487H1	SATMON001	g1399389	BLASTN	655	1e-45	83
147	16295	LIB3062-011-Q1-K1-A5	LIB3062	g1206012	BLASTN	1263	1e-101	80
148	16295	LIB3062-005-Q1-K1-B1	LIB3062	g1206012	BLASTN	966	1e-87	79
149	16295	LIB3062-053-Q1-K1-C7	LIB3062	g435312	BLASTN	1142	1e-86	79
150	16295	LIB3062-009-Q1-K1-F11	LIB3062	g1206012	BLASTN	558	1e-71	84
151	16295	700020626H1	SATMON001	g435312	BLASTN	819	1e-59	83
152	16295	700218004H1	SATMON016	g1206012	BLASTN	800	1e-57	81
153	16464	700282508H2	SATMON024	g21955	BLASTX	87	1e-22	58
154	16478	700333375H1	SATMON019	g1399390	BLASTX	76	1e-8	58
155	19731	700282461H2	SATMON024	g1769814	BLASTX	156	1e-14	56
156	22706	700172681H1	SATMON013	g1805413	BLASTX	148	1e-13	62
157	22706	700172682H1	SATMON013	g1805413	BLASTX	133	1e-11	59
158	295	LIB3067-037-Q1-K1-C6	LIB3067	g435312	BLASTN	2303	1e-183	99
159	295	LIB3062-013-Q1-K1-G1	LIB3062	g435312	BLASTN	2294	1e-182	99
160	295	LIB3078-053-	LIB3078	g1206012	BLASTN	2216	1e-179	98

161	295	Q1-K1-B7 LIB3062-036- Q1-K1-G11	LIB3062	g435312	BLASTN	2182	1e-173	99
162	295	LIB3062-004- Q1-K1-B11	LIB3062	g435312	BLASTN	2183	1e-173	98
163	295	LIB3069-038- Q1-K1-D1	LIB3069	g435312	BLASTN	2187	1e-173	95
164	295	LIB3067-046- Q1-K1-C11	LIB3067	g435312	BLASTN	2163	1e-171	99
165	295	LIB3069-051- Q1-K1-G1	LIB3069	g435312	BLASTN	1998	1e-166	97
166	295	LIB3062-038- Q1-K1-G10	LIB3062	g435312	BLASTN	2066	1e-163	98
167	295	LIB3062-050- Q1-K1-E4	LIB3062	g435312	BLASTN	1382	1e-162	92
168	295	LIB3078-039- Q1-K1-H6	LIB3078	g1206012	BLASTN	1603	1e-158	95
169	295	LIB3069-004- Q1-K1-A9	LIB3069	g435312	BLASTN	1724	1e-158	96
170	295	LIB3062-039- Q1-K1-H6	LIB3062	g435312	BLASTN	1712	1e-157	99
171	295	LIB3062-056- Q1-K1-D3	LIB3062	g435312	BLASTN	1990	1e-157	94
172	295	LIB3069-048- Q1-K1-A10	LIB3069	g435312	BLASTN	1851	1e-156	95
173	295	LIB83-004- Q1-E2-F6	LIB83	g1206012	BLASTN	1928	1e-154	98
174	295	LIB3069-033- Q1-K1-G7	LIB3069	g435312	BLASTN	1939	1e-152	98
175	295	LIB3067-058- Q1-K1-B7	LIB3067	g435312	BLASTN	1753	1e-151	99
176	295	LIB3069-017- Q1-K1-F10	LIB3069	g435312	BLASTN	1812	1e-148	94
177	295	LIB143-011- Q1-E1-A1	LIB143	g435312	BLASTN	1822	1e-143	95
178	295	700571031H1	SATMON030	g435312	BLASTN	1348	1e-136	99
179	295	700094755H1	SATMON008	g435312	BLASTN	1703	1e-133	99
180	295	LIB3069-002- Q1-K1-G7	LIB3069	g1399389	BLASTN	1275	1e-132	95
181	295	LIB143-031- Q1-E1-B3	LIB143	g799376	BLASTN	895	1e-131	94
182	295	700623229H1	SATMON034	g435312	BLASTN	1671	1e-130	99
183	295	700572265H1	SATMON030	g435312	BLASTN	1529	1e-129	98
184	295	LIB143-030- Q1-E1-H1	LIB143	g435312	BLASTN	1605	1e-129	100
185	295	LIB3078-018- Q1-K1-C11	LIB3078	g1206012	BLASTN	1643	1e-128	99
186	295	700047584H1	SATMON003	g799376	BLASTN	1645	1e-128	100
187	295	700095023H1	SATMON008	g435312	BLASTN	1653	1e-128	99
188	295	LIB3062-032- Q1-K1-C1	LIB3062	g435312	BLASTN	991	1e-127	81
189	295	700619910H1	SATMON034	g435312	BLASTN	1258	1e-127	95
190	295	700048340H1	SATMON003	g435312	BLASTN	1620	1e-126	100
191	295	700095521H1	SATMON008	g435312	BLASTN	1620	1e-126	100

192	295	700071964H1	SATMON007	g435312	BLASTN	1610	1e-125	100
193	295	LIB143-012-Q1-E1-A8	LIB143	g435312	BLASTN	1615	1e-125	100
194	295	700620843H1	SATMON034	g435312	BLASTN	870	1e-124	100
195	295	700623282H1	SATMON034	g435312	BLASTN	1600	1e-124	100
196	295	700405326H1	SATMON029	g435312	BLASTN	1603	1e-124	99
197	295	700201262H1	SATMON003	g435312	BLASTN	1394	1e-123	99
198	295	700096113H1	SATMON008	g435312	BLASTN	1590	1e-123	100
199	295	700092753H1	SATMON008	g435312	BLASTN	1590	1e-123	100
200	295	700091708H1	SATMON011	g1206012	BLASTN	941	1e-122	98
201	295	700207847H1	SATMON016	g435312	BLASTN	1161	1e-121	99
202	295	700106675H1	SATMON010	g435312	BLASTN	1523	1e-121	99
203	295	700405420H1	SATMON029	g435312	BLASTN	1560	1e-121	100
204	295	700094707H1	SATMON008	g435312	BLASTN	1552	1e-120	99
205	295	700099904H1	SATMON009	g1206012	BLASTN	1556	1e-120	99
206	295	700047601H1	SATMON003	g799376	BLASTN	541	1e-119	99
207	295	700573387H1	SATMON030	g435312	BLASTN	1353	1e-119	96
208	295	700571123H1	SATMON030	g435312	BLASTN	1446	1e-119	98
209	295	700092870H1	SATMON008	g435312	BLASTN	1535	1e-119	100
210	295	700099718H1	SATMON009	g1206012	BLASTN	1535	1e-119	97
211	295	700092358H1	SATMON008	g435312	BLASTN	1540	1e-119	100
212	295	700047779H1	SATMON003	g435312	BLASTN	1526	1e-118	99
213	295	700100960H1	SATMON009	g1206012	BLASTN	1510	1e-117	95
214	295	700072841H1	SATMON007	g435312	BLASTN	1520	1e-117	100
215	295	700094663H1	SATMON008	g1206012	BLASTN	1521	1e-117	99
216	295	700093094H1	SATMON008	g435312	BLASTN	760	1e-116	98
217	295	700103422H1	SATMON010	g435312	BLASTN	1508	1e-116	99
218	295	700093551H1	SATMON008	g435312	BLASTN	1508	1e-116	99
219	295	700075211H1	SATMON007	g435312	BLASTN	1429	1e-115	99
220	295	700095166H1	SATMON008	g435312	BLASTN	1486	1e-115	99
221	295	700075959H1	SATMON007	g435312	BLASTN	1487	1e-115	98
222	295	700334933H1	SATMON019	g435312	BLASTN	1490	1e-115	100
223	295	700093526H1	SATMON008	g435312	BLASTN	1286	1e-114	99
224	295	700623754H1	SATMON034	g435312	BLASTN	1392	1e-114	98
225	295	700205454H1	SATMON003	g435312	BLASTN	1485	1e-114	100
226	295	700623202H1	SATMON034	g435312	BLASTN	789	1e-113	97
227	295	700095616H1	SATMON008	g435312	BLASTN	1229	1e-113	95
228	295	LIB3067-032-Q1-K1-A2	LIB3067	g435312	BLASTN	1376	1e-113	96
229	295	700202823H1	SATMON003	g1399389	BLASTN	1438	1e-113	97
230	295	700096024H1	SATMON008	g435312	BLASTN	1465	1e-113	100
231	295	700096336H1	SATMON008	g435312	BLASTN	1465	1e-113	100
232	295	LIB143-063-Q1-E1-B2	LIB143	g1206012	BLASTN	1000	1e-112	96
233	295	700238549H1	SATMON010	g435312	BLASTN	1450	1e-112	100
234	295	700244114H1	SATMON010	g435312	BLASTN	1450	1e-112	100
235	295	700028461H1	SATMON003	g799376	BLASTN	1455	1e-112	100
236	295	700093584H1	SATMON008	g435312	BLASTN	1440	1e-111	100
237	295	700075942H1	SATMON007	g435312	BLASTN	1440	1e-111	100
238	295	700096338H1	SATMON008	g435312	BLASTN	1443	1e-111	99
239	295	700101061H1	SATMON009	g1206012	BLASTN	1446	1e-111	99
240	295	700072221H1	SATMON007	g435312	BLASTN	1448	1e-111	99
241	295	700202889H1	SATMON003	g435312	BLASTN	1294	1e-110	95
242	295	700096691H1	SATMON008	g435312	BLASTN	1430	1e-110	100

243	295	700092770H1	SATMON008	g435312	BLASTN	1431	1e-110	99
244	295	700048119H1	SATMON003	g435312	BLASTN	1434	1e-110	92
245	295	700220654H1	SATMON011	g1206012	BLASTN	1420	1e-109	100
246	295	700048558H1	SATMON003	g435312	BLASTN	1423	1e-109	98
247	295	700096602H1	SATMON008	g435312	BLASTN	1271	1e-108	99
248	295	700094103H1	SATMON008	g435312	BLASTN	1406	1e-108	99
249	295	700333041H1	SATMON019	g435312	BLASTN	1413	1e-108	99
250	295	700239210H1	SATMON010	g435312	BLASTN	1213	1e-107	97
251	295	700030527H1	SATMON003	g435312	BLASTN	1339	1e-107	95
252	295	700101347H1	SATMON009	g1206012	BLASTN	1401	1e-107	97
253	295	700457234H1	SATMON029	g435312	BLASTN	1379	1e-106	97
254	295	700030281H1	SATMON003	g799376	BLASTN	1381	1e-106	99
255	295	700095440H1	SATMON008	g435312	BLASTN	1387	1e-106	98
256	295	700220367H1	SATMON011	g1206012	BLASTN	1302	1e-105	99
257	295	700573418H2	SATMON030	g435312	BLASTN	1369	1e-105	99
258	295	700469112H1	SATMON025	g1399389	BLASTN	1370	1e-105	100
259	295	700221610H1	SATMON011	g1206012	BLASTN	1370	1e-105	100
260	295	700096680H1	SATMON008	g435312	BLASTN	1374	1e-105	99
261	295	700029615H1	SATMON003	g435312	BLASTN	1377	1e-105	99
262	295	700456615H1	SATMON029	g435312	BLASTN	1356	1e-104	99
263	295	700221707H1	SATMON011	g1206012	BLASTN	1357	1e-104	98
264	295	700105740H1	SATMON010	g435312	BLASTN	1359	1e-104	98
265	295	700235691H1	SATMON010	g435312	BLASTN	1363	1e-104	99
266	295	700348545H1	SATMON023	g435312	BLASTN	1050	1e-103	100
267	295	700028370H1	SATMON003	g435312	BLASTN	1065	1e-103	100
268	295	700027182H1	SATMON003	g435312	BLASTN	1348	1e-103	99
269	295	700106536H1	SATMON010	g435312	BLASTN	1350	1e-103	100
270	295	700105843H1	SATMON010	g435312	BLASTN	1353	1e-103	99
271	295	700106129H1	SATMON010	g435312	BLASTN	1079	1e-102	97
272	295	700242556H1	SATMON010	g435312	BLASTN	1225	1e-102	100
273	295	700027085H1	SATMON003	g435312	BLASTN	1335	1e-102	100
274	295	700085893H1	SATMON011	g1206012	BLASTN	1335	1e-102	98
275	295	700029004H1	SATMON003	g799376	BLASTN	1335	1e-102	100
276	295	700457191H1	SATMON029	g435312	BLASTN	1336	1e-102	99
277	295	700096560H1	SATMON008	g435312	BLASTN	1339	1e-102	99
278	295	700237805H1	SATMON010	g435312	BLASTN	1340	1e-102	100
279	295	700096210H1	SATMON008	g435312	BLASTN	916	1e-101	99
280	295	700222320H1	SATMON011	g1206012	BLASTN	1320	1e-101	100
281	295	700240040H1	SATMON010	g435312	BLASTN	1320	1e-101	100
282	295	700052948H1	SATMON007	g435312	BLASTN	1325	1e-101	100
283	295	700048831H1	SATMON003	g435312	BLASTN	835	1e-100	98
284	295	700041531H1	SATMON004	g1206012	BLASTN	1188	1e-100	98
285	295	700029745H1	SATMON003	g435312	BLASTN	1310	1e-100	100
286	295	700243513H1	SATMON010	g435312	BLASTN	1311	1e-100	99
287	295	700029026H1	SATMON003	g435312	BLASTN	1315	1e-100	100
288	295	700241584H1	SATMON010	g435312	BLASTN	1316	1e-100	99
289	295	700073024H1	SATMON007	g1399389	BLASTN	853	1e-99	96
290	295	700223548H1	SATMON011	g1206012	BLASTN	1301	1e-99	99
291	295	700236626H1	SATMON010	g1399389	BLASTN	1305	1e-99	100
292	295	LIB143-008- Q1-E1-E8	LIB143	g1399389	BLASTN	738	1e-98	92
293	295	700095640H1	SATMON008	g1399389	BLASTN	1285	1e-98	95
294	295	700084078H1	SATMON011	g1206012	BLASTN	1287	1e-98	97
295	295	700422134H1	SATMONN01	g1206012	BLASTN	1290	1e-98	100

296	295	700162241H1	SATMON012	g435312	BLASTN	1290	1e-98	100
297	295	700102975H1	SATMON010	g435312	BLASTN	613	1e-97	98
298	295	700050492H1	SATMON003	g1399389	BLASTN	1010	1e-97	99
299	295	700204531H1	SATMON003	g1399389	BLASTN	1216	1e-96	98
300	295	700103594H1	SATMON010	g435312	BLASTN	1260	1e-96	100
301	295	701163840H1	SATMONN04	g435312	BLASTN	1262	1e-96	99
302	295	700332403H1	SATMON019	g435312	BLASTN	1268	1e-96	99
303	295	700236618H1	SATMON010	g435312	BLASTN	1248	1e-95	99
304	295	700149854H1	SATMON007	g435312	BLASTN	1255	1e-95	100
305	295	700104119H1	SATMON010	g435312	BLASTN	1255	1e-95	92
306	295	700167667H1	SATMON013	g435312	BLASTN	1255	1e-95	100
307	295	700096406H1	SATMON008	g435312	BLASTN	1235	1e-94	100
308	295	700573958H1	SATMON030	g435312	BLASTN	1239	1e-94	98
309	295	700154663H1	SATMON007	g435312	BLASTN	1240	1e-94	100
310	295	700243347H1	SATMON010	g435312	BLASTN	1243	1e-94	99
311	295	700350103H1	SATMON023	g435312	BLASTN	580	1e-93	96
312	295	700106454H1	SATMON010	g435312	BLASTN	803	1e-92	87
313	295	700105284H1	SATMON010	g1399389	BLASTN	1098	1e-92	97
314	295	700053186H1	SATMON008	g435312	BLASTN	1220	1e-92	100
315	295	700050806H1	SATMON003	g1399389	BLASTN	895	1e-90	100
316	295	700153593H1	SATMON007	g435312	BLASTN	1195	1e-90	100
317	295	700156871H1	SATMON012	g435312	BLASTN	1195	1e-90	96
318	295	700152345H1	SATMON007	g435312	BLASTN	1063	1e-89	98
319	295	700237990H1	SATMON010	g435312	BLASTN	1175	1e-89	100
320	295	700152994H1	SATMON007	g435312	BLASTN	1180	1e-89	100
321	295	701158395H1	SATMONN04	g799376	BLASTN	1180	1e-89	98
322	295	700622836H1	SATMON034	g435312	BLASTN	845	1e-88	96
323	295	700454336H1	SATMON029	g1399389	BLASTN	1018	1e-88	91
324	295	700574985H1	SATMON030	g435312	BLASTN	1030	1e-88	91
325	295	700094208H1	SATMON008	g1399389	BLASTN	690	1e-87	93
326	295	700094580H1	SATMON008	g435312	BLASTN	916	1e-87	99
327	295	700088251H1	SATMON011	g1206012	BLASTN	1154	1e-87	94
328	295	700203573H1	SATMON003	g799376	BLASTN	1155	1e-87	100
329	295	700152986H1	SATMON007	g435312	BLASTN	1157	1e-87	98
330	295	700457987H1	SATMON029	g435312	BLASTN	490	1e-86	95
331	295	700156186H1	SATMON007	g435312	BLASTN	1140	1e-86	100
332	295	700162244H1	SATMON012	g435312	BLASTN	1142	1e-86	98
333	295	700161070H1	SATMON012	g435312	BLASTN	1147	1e-86	98
334	295	700241478H1	SATMON010	g435312	BLASTN	790	1e-85	96
335	295	700238324H1	SATMON010	g435312	BLASTN	1126	1e-85	97
336	295	700156421H1	SATMON012	g435312	BLASTN	1127	1e-85	96
337	295	700162187H1	SATMON012	g435312	BLASTN	1130	1e-85	100
338	295	700161129H1	SATMON012	g435312	BLASTN	1135	1e-85	100
339	295	700156075H1	SATMON007	g435312	BLASTN	681	1e-84	98
340	295	700351086H1	SATMON023	g1399389	BLASTN	1045	1e-84	95
341	295	700155517H1	SATMON007	g435312	BLASTN	1115	1e-84	100
342	295	700157475H1	SATMON012	g435312	BLASTN	1115	1e-84	98
343	295	700169796H1	SATMON013	g435312	BLASTN	1120	1e-84	100
344	295	700201343H1	SATMON003	g435312	BLASTN	1104	1e-83	99
345	295	700102333H1	SATMON010	g435312	BLASTN	1106	1e-83	87
346	295	700235624H1	SATMON010	g1206012	BLASTN	1108	1e-83	86
347	295	700622378H1	SATMON034	g435312	BLASTN	412	1e-82	90
348	295	700072123H1	SATMON007	g435312	BLASTN	935	1e-82	89
349	295	700017263H1	SATMON001	g799376	BLASTN	1100	1e-82	100

350	295	LIB3062-051-Q1-K1-G10	LIB3062	g435312	BLASTN	487	1e-81	88
351	295	700457620H1	SATMON029	g1206012	BLASTN	511	1e-81	91
352	295	700164773H1	SATMON013	g1399389	BLASTN	1087	1e-81	96
353	295	700020206H1	SATMON001	g799376	BLASTN	1089	1e-81	98
354	295	700456365H1	SATMON029	g435312	BLASTN	1041	1e-80	99
355	295	700238182H1	SATMON010	g435312	BLASTN	1055	1e-79	100
356	295	700201830H1	SATMON003	g435312	BLASTN	1058	1e-79	99
357	295	700155920H1	SATMON007	g435312	BLASTN	873	1e-77	97
358	295	701158695H1	SATMONN04	g435312	BLASTN	913	1e-77	94
359	295	700029134H1	SATMON003	g435312	BLASTN	1031	1e-77	99
360	295	701185615H1	SATMONN06	g1399389	BLASTN	465	1e-75	100
361	295	700621515H1	SATMON034	g1399389	BLASTN	616	1e-75	88
362	295	700094517H1	SATMON008	g435312	BLASTN	581	1e-74	92
363	295	700162382H1	SATMON012	g435312	BLASTN	846	1e-74	96
364	295	LIB143-028-Q1-E1-C5	LIB143	g1206012	BLASTN	995	1e-74	100
365	295	700158308H1	SATMON012	g435312	BLASTN	1000	1e-74	100
366	295	700242248H1	SATMON010	g1206012	BLASTN	1001	1e-74	86
367	295	700153633H1	SATMON007	g435312	BLASTN	1001	1e-74	99
368	295	701158495H1	SATMONN04	g799376	BLASTN	970	1e-72	87
369	295	LIB143-028-Q1-E1-G8	LIB143	g435312	BLASTN	975	1e-72	100
370	295	700093581H1	SATMON008	g435312	BLASTN	965	1e-71	100
371	295	700051874H1	SATMON003	g435312	BLASTN	738	1e-70	94
372	295	700570142H1	SATMON030	g1206012	BLASTN	559	1e-69	86
373	295	700238661H1	SATMON010	g435312	BLASTN	831	1e-69	98
374	295	700153172H1	SATMON007	g435312	BLASTN	941	1e-69	95
375	295	700153988H1	SATMON007	g435312	BLASTN	921	1e-67	98
376	295	700352641H1	SATMON024	g1206012	BLASTN	842	1e-66	97
377	295	700030142H1	SATMON003	g435312	BLASTN	900	1e-66	100
378	295	701164693H1	SATMONN04	g435312	BLASTN	317	1e-64	94
379	295	700159524H1	SATMON012	g1399389	BLASTN	535	1e-64	86
380	295	700164779H1	SATMON013	g1399389	BLASTN	647	1e-63	93
381	295	700467312H1	SATMON025	g435312	BLASTN	836	1e-63	99
382	295	700161392H1	SATMON012	g1206012	BLASTN	676	1e-62	87
383	295	700334631H1	SATMON019	g435312	BLASTN	823	1e-59	99
384	295	700242893H1	SATMON010	g435312	BLASTN	793	1e-57	94
385	295	700621992H1	SATMON034	g1399389	BLASTN	235	1e-55	93
386	295	700349745H1	SATMON023	g435312	BLASTN	750	1e-53	100
387	295	700050250H1	SATMON003	g435312	BLASTN	483	1e-52	98
388	295	700456624H1	SATMON029	g435312	BLASTN	726	1e-51	98
389	295	700149879H1	SATMON007	g1206012	BLASTN	602	1e-50	83
390	295	700458589H1	SATMON029	g435312	BLASTN	436	1e-48	93
391	295	700168245H1	SATMON013	g435312	BLASTN	690	1e-48	100
392	295	700151362H1	SATMON007	g435312	BLASTN	430	1e-47	100
393	295	700075679H1	SATMON007	g435312	BLASTN	505	1e-44	90
394	295	700456649H1	SATMON029	g1206012	BLASTN	644	1e-44	97
395	295	700236165H1	SATMON010	g435312	BLASTN	645	1e-44	100
396	295	700236174H1	SATMON010	g435312	BLASTN	623	1e-43	99
397	295	700095081H1	SATMON008	g435312	BLASTN	630	1e-43	100
398	295	700150082H1	SATMON007	g1399389	BLASTN	612	1e-42	91
399	295	700456926H1	SATMON029	g435312	BLASTN	331	1e-41	99
400	295	700281403H2	SATMON019	g435312	BLASTN	303	1e-33	92

401	295	700095048H1	SATMON008	g435312	BLASTN	503	1e-33	99
402	295	700053540H1	SATMON010	g1206012	BLASTN	478	1e-31	89
403	295	700241965H1	SATMON010	g435312	BLASTN	313	1e-30	90
404	295	700623494H1	SATMON034	g435312	BLASTN	448	1e-28	91
405	295	700076766H1	SATMON007	g435312	BLASTN	433	1e-27	98
406	3015	LIB3069-005-Q1-K1-F9	LIB3069	g804655	BLASTN	648	1e-74	80
407	3015	700475354H1	SATMON025	g804655	BLASTN	299	1e-46	81
408	3015	700456167H1	SATMON029	g804655	BLASTN	253	1e-38	78
409	3015	700473368H1	SATMON025	g804656	BLASTX	270	1e-37	66
410	3015	700350439H1	SATMON023	g1143863	BLASTN	533	1e-35	65
411	3015	700469778H1	SATMON025	g804656	BLASTX	148	1e-23	74
412	31009	LIB3066-030-Q1-K1-A2	LIB3066	g804656	BLASTX	304	1e-51	56
413	31009	700355331H1	SATMON024	g804656	BLASTX	235	1e-25	51
414	31970	LIB3067-027-Q1-K1-B9	LIB3067	g804656	BLASTX	202	1e-36	58
415	32475	LIB3066-035-Q1-K1-F1	LIB3066	g40644	BLASTX	138	1e-32	56
416	32588	LIB143-030-Q1-E1-G7	LIB143	g804655	BLASTN	463	1e-27	85
417	32588	700096317H1	SATMON008	g804656	BLASTX	119	1e-10	71
418	32784	700027379H1	SATMON003	g21955	BLASTX	118	1e-9	56
419	3311	LIB3062-026-Q1-K1-H8	LIB3062	g1143863	BLASTN	1526	1e-118	81
420	3311	LIB3078-022-Q1-K1-A8	LIB3078	g1143863	BLASTN	1394	1e-107	82
421	3311	LIB3062-021-Q1-K1-C6	LIB3062	g1143863	BLASTN	1396	1e-107	82
422	3311	LIB143-021-Q1-E1-C8	LIB143	g1143863	BLASTN	811	1e-88	78
423	3311	LIB3062-020-Q1-K1-A11	LIB3062	g1143863	BLASTN	404	1e-87	81
424	3311	LIB3062-052-Q1-K1-H2	LIB3062	g1143863	BLASTN	712	1e-86	83
425	3311	700207937H1	SATMON016	g1143863	BLASTN	1127	1e-85	84
426	3311	LIB3078-022-Q1-K1-C8	LIB3078	g1143863	BLASTN	1127	1e-85	79
427	3311	700613304H1	SATMON033	g1143863	BLASTN	709	1e-84	84
428	3311	700083153H1	SATMON011	g1143863	BLASTN	1087	1e-81	84
429	3311	LIB3059-044-Q1-K1-D4	LIB3059	g1143863	BLASTN	1087	1e-81	86
430	3311	700083988H1	SATMON011	g1143863	BLASTN	1074	1e-80	82
431	3311	700091689H1	SATMON011	g1143863	BLASTN	1061	1e-79	83
432	3311	700211382H1	SATMON016	g1143863	BLASTN	1065	1e-79	83
433	3311	700093849H1	SATMON008	g1143863	BLASTN	1065	1e-79	81
434	3311	700265072H1	SATMON017	g1143863	BLASTN	1052	1e-78	84
435	3311	700224821H1	SATMON011	g1143863	BLASTN	1039	1e-77	84
436	3311	LIB3068-010-Q1-K1-E12	LIB3068	g1143863	BLASTN	455	1e-76	74
437	3311	700073004H1	SATMON007	g1143863	BLASTN	540	1e-76	83
438	3311	700351608H1	SATMON023	g1143863	BLASTN	1026	1e-76	82
439	3311	700077236H1	SATMON007	g1143863	BLASTN	1029	1e-76	85
440	3311	700614027H1	SATMON033	g1143863	BLASTN	1017	1e-75	86

441	3311	700333838H1	SATMON019	g1143863	BLASTN	1004	1e-74	82
442	3311	LIB3061-009-Q1-K1-G12	LIB3061	g804656	BLASTX	433	1e-73	62
443	3311	LIB3067-044-Q1-K1-H4	LIB3067	g1143863	BLASTN	587	1e-71	76
444	3311	700469775H1	SATMON025	g1143863	BLASTN	960	1e-71	84
445	3311	700214450H1	SATMON016	g1143863	BLASTN	968	1e-71	85
446	3311	700571763H1	SATMON030	g1143863	BLASTN	702	1e-70	82
447	3311	700256818H1	SATMON017	g1143863	BLASTN	949	1e-70	79
448	3311	700087102H1	SATMON011	g1143863	BLASTN	951	1e-70	79
449	3311	700614486H1	SATMON033	g1143863	BLASTN	774	1e-68	79
450	3311	LIB3069-003-Q1-K1-D3	LIB3069	g1143863	BLASTN	918	1e-67	70
451	3311	700351075H1	SATMON023	g1143863	BLASTN	920	1e-67	84
452	3311	LIB143-034-Q1-E1-F3	LIB143	g804656	BLASTX	406	1e-64	70
453	3311	700469282H1	SATMON025	g1143863	BLASTN	469	1e-61	85
454	3311	700224535H1	SATMON011	g1143863	BLASTN	840	1e-61	80
455	3311	700457618H1	SATMON029	g1143863	BLASTN	845	1e-61	80
456	3311	700469464H1	SATMON025	g1143863	BLASTN	481	1e-60	83
457	3311	700161193H1	SATMON012	g1143863	BLASTN	820	1e-59	84
458	3311	700196802H1	SATMON014	g1143863	BLASTN	777	1e-55	79
459	3311	LIB3068-029-Q1-K1-H10	LIB3068	g1143863	BLASTN	703	1e-50	79
460	3311	700205028H1	SATMON003	g1143863	BLASTN	354	1e-45	79
461	3311	700334602H1	SATMON019	g804656	BLASTX	382	1e-45	77
462	3311	700018146H1	SATMON001	g1143863	BLASTN	629	1e-43	80
463	3311	700351771H1	SATMON023	g1143863	BLASTN	611	1e-42	81
464	3311	700206661H1	SATMON003	g1143863	BLASTN	604	1e-41	79
465	3311	700616407H1	SATMON033	g1143863	BLASTN	522	1e-38	83
466	3311	700053754H1	SATMON011	g804655	BLASTN	401	1e-37	70
467	3311	700469563H1	SATMON025	g1143863	BLASTN	510	1e-33	88
468	3311	700616186H1	SATMON033	g1143864	BLASTX	128	1e-31	72
469	3311	700261867H1	SATMON017	g804656	BLASTX	175	1e-27	79
470	3311	700223387H1	SATMON011	g1143863	BLASTN	427	1e-25	82
471	3311	700334681H1	SATMON019	g804656	BLASTX	226	1e-24	77
472	3311	700086604H1	SATMON011	g1143863	BLASTN	349	1e-20	85
473	3311	LIB3069-018-Q1-K1-B8	LIB3069	g1143863	BLASTN	234	1e-16	86
474	7962	700088807H1	SATMON011	g1143863	BLASTN	977	1e-72	83
475	7962	700582628H1	SATMON031	g1143863	BLASTN	838	1e-61	77
476	8824	700266114H1	SATMON017	g804655	BLASTN	514	1e-40	78
477	8824	700258493H1	SATMON017	g804655	BLASTN	361	1e-21	84
478	8824	LIB143-050-Q1-E1-D9	LIB143	g804655	BLASTN	241	1e-9	83
479	9273	700153802H1	SATMON007	g1143864	BLASTX	151	1e-14	59

SOYBEAN ADENINE PHOSPHORIBOSYL TRANSFERASE (EC 2.4.2.7)

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
480	-700661182	700661182H1	SOYMON005	g16164	BLASTX	151	1e-14	70
481	-700832379	700832379H1	SOYMON019	g16164	BLASTX	134	1e-33	76
482	-700942837	700942837H1	SOYMON024	g16163	BLASTN	472	1e-30	71
483	-701038319	701038319H1	SOYMON029	g16164	BLASTX	196	1e-20	84

484	-701052195	701052195H1	SOYMON032	g1321681	BLASTX	223	1e-33	74
485	-701054342	701054342H1	SOYMON032	g16163	BLASTN	383	1e-21	68
486	-701099058	701099058H1	SOYMON028	g16163	BLASTN	782	1e-56	76
487	-701103094	701103094H1	SOYMON028	g1321681	BLASTX	150	1e-13	71
488	-701205435	701205435H1	SOYMON035	g16164	BLASTX	298	1e-39	65
489	-GM1685	LIB3028-009-Q1-B1-H2	LIB3028	g16163	BLASTN	408	1e-38	73
490	11402	700898558H1	SOYMON027	g1402893	BLASTN	690	1e-48	73
491	11402	700734213H1	SOYMON010	g1321681	BLASTX	194	1e-40	76
492	24349	701055830H1	SOYMON032	g1321681	BLASTX	221	1e-24	81
493	24349	701211541H1	SOYMON035	g1321681	BLASTX	191	1e-19	86
494	26036	701044219H1	SOYMON032	g1402893	BLASTN	474	1e-29	73
495	26036	701053252H1	SOYMON032	g1321681	BLASTX	194	1e-20	81
496	26036	701051166H1	SOYMON032	g1321681	BLASTX	176	1e-17	81
497	26036	700977973H1	SOYMON009	g1321681	BLASTX	177	1e-17	83
498	4852	701121868H1	SOYMON037	g16163	BLASTN	786	1e-56	77
499	4852	701206156H1	SOYMON035	g16163	BLASTN	568	1e-38	77
500	4852	700651407H1	SOYMON003	g16163	BLASTN	492	1e-30	75
501	4852	701127525H1	SOYMON037	g16163	BLASTN	473	1e-29	75
502	4852	700895694H1	SOYMON027	g16163	BLASTN	434	1e-27	77
503	4852	700560253H1	SOYMON001	g16164	BLASTX	199	1e-21	82
504	4852	701138208H1	SOYMON038	g16164	BLASTX	175	1e-17	80
505	4852	701137650H1	SOYMON038	g16163	BLASTN	317	1e-15	75
506	4852	LIB3040-031-Q1-E2-D4	LIB3040	g16163	BLASTN	308	1e-14	80
507	4852	701066549H1	SOYMON034	g16164	BLASTX	142	1e-12	84
508	4852	700893483H1	SOYMON024	g16164	BLASTX	134	1e-11	83
509	8624	700653006H1	SOYMON003	g16163	BLASTN	605	1e-40	70
510	8624	LIB3039-040-Q1-E1-C8	LIB3039	g16163	BLASTN	511	1e-31	67
511	8624	701135156H1	SOYMON038	g16164	BLASTX	207	1e-22	72
512	8624	701054685H1	SOYMON032	g726305	BLASTX	199	1e-21	80
513	8624	700834123H1	SOYMON019	g16164	BLASTX	182	1e-18	67
514	8624	700830624H1	SOYMON019	g1321681	BLASTX	128	1e-10	73
515	8624	700833413H1	SOYMON019	g1321681	BLASTX	129	1e-10	65

SOYBEAN β GLUCOSIDASE (EC 3.2.1.21)

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
516	-700747805	700747805H1	SOYMON013	g1155090	BLASTX	152	1e-13	65
517	-700749102	700749102H1	SOYMON013	g1155255	BLASTX	141	1e-12	69
518	-700749944	700749944H1	SOYMON013	g1155255	BLASTX	169	1e-18	54
519	-700837183	700837183H1	SOYMON020	g804656	BLASTX	161	1e-15	81
520	-700894885	700894885H1	SOYMON024	g142580	BLASTX	114	1e-8	61
521	-700907258	700907258H1	SOYMON022	g804656	BLASTX	187	1e-18	46
522	-700965238	700965238H1	SOYMON022	g1155255	BLASTX	145	1e-23	51
523	-700967625	700967625H1	SOYMON032	g21953	BLASTX	165	1e-15	64
524	-700978836	700978836H1	SOYMON009	g804656	BLASTX	187	1e-18	43
525	-700982375	700982375H1	SOYMON009	g1155090	BLASTX	109	1e-9	44
526	-701043315	701043315H1	SOYMON029	g1155255	BLASTX	170	1e-19	57
527	-701054964	701054964H1	SOYMON032	g757740	BLASTX	173	1e-16	61
528	-701055914	701055914H1	SOYMON032	g21953	BLASTX	176	1e-25	69
529	-701127573	701127573H1	SOYMON037	g21953	BLASTX	121	1e-20	74

530	-701131494	701131494H1	SOYMON038	g1143864	BLASTX	133	1e-21	68
531	-701138450	701138450H1	SOYMON038	g21955	BLASTX	235	1e-25	55
532	-701146550	701146550H1	SOYMON031	g21955	BLASTX	65	1e-9	66
533	-701203719	701203719H2	SOYMON035	g1143864	BLASTX	160	1e-14	46
534	-701213534	701213534H1	SOYMON035	g1483154	BLASTX	78	1e-9	48
535	-GM10346	LIB3049-006-Q1-E1-H5	LIB3049	g3201553	BLASTN	742	1e-51	63
536	-GM12457	LIB3049-044-Q1-E1-A10	LIB3049	g3201553	BLASTN	683	1e-48	65
537	-GM34028	LIB3051-038-Q1-K1-D2	LIB3051	g249262	BLASTX	78	1e-27	53
538	-GM37305	LIB3051-073-Q1-K1-G10	LIB3051	g804656	BLASTX	258	1e-44	68
539	-GM40790	LIB3051-105-Q1-K1-F4	LIB3051	g804656	BLASTX	362	1e-72	69
540	11009	700660118H1	SOYMON004	g804656	BLASTX	244	1e-26	79
541	11009	700746718H1	SOYMON013	g804656	BLASTX	121	1e-11	61
542	11009	701061014H1	SOYMON033	g804656	BLASTX	117	1e-9	72
543	12181	701146853H1	SOYMON031	g21955	BLASTX	162	1e-22	62
544	12181	701040693H1	SOYMON029	g21955	BLASTX	162	1e-21	63
545	12181	701212713H1	SOYMON035	g21955	BLASTX	162	1e-19	67
546	12623	701141106H1	SOYMON038	g142580	BLASTX	163	1e-15	65
547	12623	700975287H1	SOYMON009	g142580	BLASTX	126	1e-10	65
548	12814	LIB3051-053-Q1-K2-H12	LIB3051	g804656	BLASTX	243	1e-42	61
549	12814	LIB3052-007-Q1-B1-G11	LIB3052	g1143863	BLASTN	613	1e-40	66
550	12814	LIB3051-111-Q1-K1-C12	LIB3051	g1143863	BLASTN	608	1e-39	67
551	12814	700656613H1	SOYMON004	g1143863	BLASTN	572	1e-38	67
552	12814	701068027H2	SOYMON034	g804656	BLASTX	205	1e-35	86
553	12814	701211582H1	SOYMON035	g804656	BLASTX	247	1e-35	82
554	12814	700986671H1	SOYMON009	g804656	BLASTX	201	1e-31	58
555	12814	701153690H1	SOYMON031	g804656	BLASTX	270	1e-30	63
556	12814	701066513H1	SOYMON034	g804656	BLASTX	120	1e-28	88
557	12814	700838612H1	SOYMON020	g1143863	BLASTN	348	1e-18	70
558	13173	LIB3049-018-Q1-E1-B7	LIB3049	g1143863	BLASTN	745	1e-51	66
559	13173	LIB3049-001-Q1-E1-G7	LIB3049	g1143863	BLASTN	623	1e-41	65
560	13173	LIB3051-111-Q1-K1-B12	LIB3051	g804656	BLASTX	215	1e-40	64
561	13173	700837342H1	SOYMON020	g804656	BLASTX	337	1e-39	70
562	13173	700838567H1	SOYMON020	g804656	BLASTX	339	1e-39	69
563	13173	LIB3051-111-Q1-K1-B10	LIB3051	g1143863	BLASTN	574	1e-36	67
564	13173	LIB3051-114-Q1-K1-B10	LIB3051	g804656	BLASTX	89	1e-31	53
565	13173	700761996H1	SOYMON015	g804656	BLASTX	279	1e-31	64
566	13173	700971311H1	SOYMON005	g804656	BLASTX	270	1e-30	58
567	13173	700854217H1	SOYMON023	g804656	BLASTX	158	1e-27	58
568	13173	701063334H1	SOYMON033	g1143863	BLASTN	460	1e-27	62
569	13173	700900778H1	SOYMON027	g804656	BLASTX	199	1e-20	76
570	1499	LIB3039-014-	LIB3039	g1155090	BLASTX	144	1e-44	53

Q1-E1-D8								
571	1499	701069538H1	SOYMON034	g1155255	BLASTX	221	1e-23	52
572	1499	701064351H1	SOYMON034	g21953	BLASTX	153	1e-17	50
573	1499	700651844H1	SOYMON003	g1155090	BLASTX	91	1e-10	55
574	1499	701068724H1	SOYMON034	g1155090	BLASTX	125	1e-9	48
575	150	700653669H1	SOYMON003	g21953	BLASTX	219	1e-23	58
576	150	700651748H1	SOYMON003	g21953	BLASTX	193	1e-19	60
577	150	701127306H1	SOYMON037	g21953	BLASTX	188	1e-18	58
578	150	700561901H1	SOYMON002	g21955	BLASTX	156	1e-14	64
579	150	701129795H1	SOYMON037	g21955	BLASTX	143	1e-12	68
580	150	701126390H1	SOYMON037	g21955	BLASTX	143	1e-12	68
581	150	701125867H1	SOYMON037	g21955	BLASTX	134	1e-11	63
582	150	701142724H1	SOYMON038	g21955	BLASTX	135	1e-11	69
583	150	701060152H1	SOYMON033	g1155255	BLASTX	135	1e-11	62
584	150	701141927H1	SOYMON038	g581738	BLASTX	127	1e-10	64
585	150	701125996H1	SOYMON037	g21955	BLASTX	127	1e-10	66
586	150	701061767H1	SOYMON033	g21953	BLASTX	89	1e-9	65
587	21571	701098566H1	SOYMON028	g21955	BLASTX	152	1e-34	59
588	21571	700846795H1	SOYMON021	g21955	BLASTX	158	1e-31	59
589	21571	701037773H1	SOYMON029	g21953	BLASTX	94	1e-22	59
590	22050	701039143H1	SOYMON029	g21953	BLASTX	86	1e-14	63
591	24776	701149235H1	SOYMON031	g21955	BLASTX	181	1e-17	55
592	30906	LIB3028-006-	LIB3028	g2077896	BLASTX	113	1e-24	47
Q1-B1-F2								
593	3094	700564240H1	SOYMON002	g21955	BLASTX	197	1e-20	80
594	3094	700564288H1	SOYMON002	g21955	BLASTX	199	1e-20	79
595	3094	701042714H1	SOYMON029	g21955	BLASTX	179	1e-17	77
596	3094	700565738H1	SOYMON002	g21955	BLASTX	160	1e-16	85
597	32420	LIB3030-008-	LIB3030	g1143863	BLASTN	551	1e-35	61
Q1-B1-H11								
598	32420	700963106H1	SOYMON022	g804656	BLASTX	219	1e-23	50
599	33821	700847344H1	SOYMON021	g21955	BLASTX	163	1e-26	58
600	4085	LIB3053-005-	LIB3053	g40665	BLASTX	195	1e-42	59
Q1-N1-F10								
601	4085	700548207H1	SOYMON002	g1155255	BLASTX	144	1e-24	60
602	587	LIB3039-005-	LIB3039	g1155090	BLASTX	199	1e-38	56
Q1-E1-F2								
603	587	LIB3039-048-	LIB3039	g21955	BLASTX	142	1e-27	65
Q1-E1-A6								
604	587	701145333H1	SOYMON031	g21953	BLASTX	149	1e-27	66
605	587	700653427H1	SOYMON003	g1155090	BLASTX	230	1e-26	56
606	587	700652854H1	SOYMON003	g1155090	BLASTX	188	1e-25	60
607	587	700751375H1	SOYMON014	g21955	BLASTX	161	1e-23	64
608	587	701063494H1	SOYMON033	g1155090	BLASTX	223	1e-23	60
609	587	700955066H1	SOYMON022	g1155090	BLASTX	216	1e-22	59
610	587	700982238H1	SOYMON009	g1155090	BLASTX	185	1e-20	61
611	587	701109014H1	SOYMON036	g1155090	BLASTX	189	1e-20	58
612	587	701145904H1	SOYMON031	g21953	BLASTX	204	1e-20	59
613	587	700982608H1	SOYMON009	g1155090	BLASTX	180	1e-19	59
614	587	700986368H1	SOYMON009	g1155090	BLASTX	184	1e-19	62
615	587	701139123H1	SOYMON038	g21953	BLASTX	155	1e-18	63
616	587	701138844H1	SOYMON038	g1155090	BLASTX	178	1e-17	61
617	587	700791212H1	SOYMON011	g1155090	BLASTX	142	1e-15	59
618	587	700646575H1	SOYMON014	g1155090	BLASTX	150	1e-15	63

619	587	700991712H1	SOYMON011	g1155090	BLASTX	153	1e-15	57
620	587	700904947H1	SOYMON022	g21955	BLASTX	159	1e-15	59
621	587	700730081H1	SOYMON009	g21955	BLASTX	159	1e-15	59
622	587	701060675H1	SOYMON033	g21955	BLASTX	160	1e-15	59
623	587	700983905H1	SOYMON009	g21953	BLASTX	114	1e-14	47
624	587	701135826H1	SOYMON038	g21955	BLASTX	155	1e-14	58
625	587	701142683H1	SOYMON038	g21953	BLASTX	155	1e-14	60
626	587	700656303H1	SOYMON004	g1155255	BLASTX	96	1e-13	41
627	587	701064503H1	SOYMON034	g21953	BLASTX	118	1e-13	62
628	587	700959789H1	SOYMON022	g1155090	BLASTX	145	1e-13	57
629	587	701104579H1	SOYMON036	g21955	BLASTX	146	1e-13	55
630	587	700975311H1	SOYMON009	g21955	BLASTX	148	1e-13	55
631	587	700987858H1	SOYMON009	g1155090	BLASTX	151	1e-13	58
632	587	700787696H2	SOYMON011	g21953	BLASTX	151	1e-13	60
633	587	700755020H1	SOYMON014	g21955	BLASTX	124	1e-12	58
634	587	700961408H1	SOYMON022	g1155090	BLASTX	126	1e-12	47
635	587	700975523H1	SOYMON009	g1155090	BLASTX	128	1e-12	50
636	587	700956261H1	SOYMON022	g1155090	BLASTX	139	1e-12	60
637	587	700986691H1	SOYMON009	g1155090	BLASTX	140	1e-12	60
638	587	700751271H1	SOYMON014	g1155090	BLASTX	141	1e-12	59
639	587	700730156H1	SOYMON009	g1155090	BLASTX	141	1e-12	59
640	587	701141713H1	SOYMON038	g21955	BLASTX	141	1e-12	60
641	587	701157330H1	SOYMON031	g1155090	BLASTX	141	1e-12	59
642	587	700967834H1	SOYMON033	g21955	BLASTX	142	1e-12	55
643	587	701155566H1	SOYMON031	g21955	BLASTX	142	1e-12	61
644	587	700751706H1	SOYMON014	g21955	BLASTX	144	1e-12	61
645	587	701145403H1	SOYMON031	g21953	BLASTX	131	1e-11	63
646	587	700959567H1	SOYMON022	g757740	BLASTX	134	1e-11	60
647	587	701064274H1	SOYMON034	g505279	BLASTX	135	1e-11	41
648	587	701151995H1	SOYMON031	g21953	BLASTX	135	1e-11	63
649	587	701050236H1	SOYMON032	g21955	BLASTX	135	1e-11	62
650	587	701152375H1	SOYMON031	g21955	BLASTX	135	1e-11	67
651	587	701155583H1	SOYMON031	g21953	BLASTX	135	1e-11	63
652	587	701156782H1	SOYMON031	g21953	BLASTX	135	1e-11	63
653	587	701149881H1	SOYMON031	g21955	BLASTX	136	1e-11	65
654	587	701151802H1	SOYMON031	g21953	BLASTX	136	1e-11	63
655	587	701147107H1	SOYMON031	g21953	BLASTX	136	1e-11	63
656	587	701157340H1	SOYMON031	g21955	BLASTX	137	1e-11	60
657	587	701142839H1	SOYMON038	g21953	BLASTX	138	1e-11	61
658	587	701148320H1	SOYMON031	g21955	BLASTX	138	1e-11	60
659	587	701156604H1	SOYMON031	g1155090	BLASTX	86	1e-10	60
660	587	701139062H1	SOYMON038	g21955	BLASTX	125	1e-10	60
661	587	701068825H1	SOYMON034	g21953	BLASTX	125	1e-10	55
662	587	701147382H1	SOYMON031	g21955	BLASTX	127	1e-10	68
663	587	701154153H1	SOYMON031	g21953	BLASTX	127	1e-10	64
664	587	701155501H1	SOYMON031	g21955	BLASTX	129	1e-10	61
665	587	701155731H1	SOYMON031	g21955	BLASTX	129	1e-10	58
666	587	701157725H1	SOYMON031	g21955	BLASTX	130	1e-10	61
667	587	700967321H1	SOYMON031	g21953	BLASTX	120	1e-9	61
668	587	701108022H1	SOYMON036	g21953	BLASTX	121	1e-9	60
669	587	701150196H1	SOYMON031	g21953	BLASTX	122	1e-9	63
670	587	701150439H1	SOYMON031	g21953	BLASTX	122	1e-9	63
671	587	701145973H1	SOYMON031	g1155090	BLASTX	99	1e-8	63
672	587	701130507H1	SOYMON038	g1155090	BLASTX	116	1e-8	50

673	587	701155886H1	SOYMON031	g21953	BLASTX	116	1e-8	62
674	587	700753795H1	SOYMON014	g21953	BLASTX	117	1e-8	62
675	7163	700560905H1	SOYMON001	g1206013	BLASTX	125	1e-16	59
676	7163	700964094H1	SOYMON022	g1155255	BLASTX	106	1e-15	61
677	7535	701065656H1	SOYMON034	g1155255	BLASTX	161	1e-28	59
678	7535	701065608H1	SOYMON034	g1155255	BLASTX	92	1e-20	61
679	7535	701063444H1	SOYMON033	g1155090	BLASTX	195	1e-19	62
680	9186	700995628H1	SOYMON011	g21953	BLASTX	161	1e-18	68
681	921	LIB3051-108-Q1-K1-H11	LIB3051	g804656	BLASTX	217	1e-58	67
682	921	700651438H1	SOYMON003	g804656	BLASTX	454	1e-55	67
683	921	LIB3051-101-Q1-K1-A6	LIB3051	g804656	BLASTX	228	1e-51	62
684	921	LIB3051-091-Q1-K1-G8	LIB3051	g804656	BLASTX	217	1e-50	67
685	921	LIB3051-096-Q1-K1-A7	LIB3051	g804656	BLASTX	269	1e-46	65
686	921	701068457H1	SOYMON034	g804656	BLASTX	230	1e-43	73
687	921	701134773H2	SOYMON038	g804656	BLASTX	332	1e-38	66
688	921	700978751H1	SOYMON009	g804656	BLASTX	211	1e-37	68
689	921	700727744H1	SOYMON009	g804656	BLASTX	305	1e-35	71
690	921	LIB3049-024-Q1-E1-G5	LIB3049	g804656	BLASTX	187	1e-34	61
691	921	700757238H1	SOYMON015	g1143863	BLASTN	465	1e-29	68
692	921	700972951H1	SOYMON005	g804656	BLASTX	168	1e-16	69
693	921	700986340H1	SOYMON009	g804656	BLASTX	172	1e-16	72
694	921	700564114H1	SOYMON002	g804656	BLASTX	172	1e-16	72
695	921	700851301H1	SOYMON023	g804656	BLASTX	172	1e-16	72
696	921	701104026H1	SOYMON036	g804656	BLASTX	172	1e-16	72
697	921	701060132H1	SOYMON033	g804656	BLASTX	172	1e-16	72
698	921	701211625H1	SOYMON035	g804656	BLASTX	172	1e-16	72
699	921	701136164H1	SOYMON038	g804656	BLASTX	173	1e-16	68
700	921	701137640H1	SOYMON038	g804656	BLASTX	160	1e-15	67
701	921	701142240H1	SOYMON038	g804656	BLASTX	157	1e-14	74
702	921	700842016H1	SOYMON020	g804656	BLASTX	136	1e-11	74
703	921	701210015H1	SOYMON035	g804656	BLASTX	136	1e-11	74
704	921	701204245H2	SOYMON035	g804656	BLASTX	125	1e-10	72
705	921	700841069H1	SOYMON020	g804656	BLASTX	119	1e-9	79
706	921	700852034H1	SOYMON023	g804656	BLASTX	119	1e-9	79
707	921	700837045H1	SOYMON020	g804656	BLASTX	120	1e-9	74
708	921	701210726H1	SOYMON035	g804656	BLASTX	120	1e-9	74
709	921	700840844H1	SOYMON020	g804656	BLASTX	122	1e-9	57
710	921	700839037H1	SOYMON020	g804656	BLASTX	115	1e-8	79

SOYBEAN ISOPENTYLTRANSFERASE

Seq No.	Cluster ID	CloneID	Library	NCBI gi	Method	Score	P-value	%Ident
711	-GM17896	LIB3055-003-Q1-N1-B10	LIB3055	g1419759	BLASTX	241	1e-42	40

***Table Headings**

Cluster ID

A cluster ID is arbitrarily assigned to all of those clones which belong to the same cluster at a given stringency and a particular clone will belong to only one cluster at a given stringency. If a cluster contains only a single clone (a “singleton”), then the cluster ID number will be negative, with an absolute value equal to the clone ID number of its single member. The cluster ID entries in the table refer to the cluster with which the particular clone in each row is associated.

Clone ID

The clone ID number refers to the particular clone in the PhytoSeq database. Each clone ID entry in the table refers to the clone whose sequence is used for (1) the sequence comparison whose scores are presented and/or (2) assignment to the particular cluster which is presented. Note that a clone may be included in this table even if its sequence comparison scores fail to meet the minimum standards for similarity. In such a case, the clone is included due solely to its association with a particular cluster for which sequences of one or more other member clones possess the required level of similarity.

Library

The library ID refers to the particular cDNA library from which a given clone is obtained. Each cDNA library is associated with the particular tissue(s), line(s) and developmental stage(s) from which it is isolated.

NCBI gi

Each sequence in the GenBank public database is arbitrarily assigned a unique NCBI gi (National Center for Biotechnology Information GenBank Identifier) number. In this table, the

NCBI gi number which is associated (in the same row) with a given clone refers to the particular GenBank sequence which is used in the sequence comparison. This entry is omitted when a clone is included solely due to its association with a particular cluster.

Method

The entry in the “Method” column of the table refers to the type of BLAST search that is used for the sequence comparison. “CLUSTER” is entered when the sequence comparison scores for a given clone fail to meet the minimum values required for significant similarity. In such cases, the clone is listed in the table solely as a result of its association with a given cluster for which sequences of one or more other member clones possess the required level of similarity.

Score

Each entry in the “Score” column of the table refers to the BLAST score that is generated by sequence comparison of the designated clone with the designated GenBank sequence using the designated BLAST method. This entry is omitted when a clone is included solely due to its association with a particular cluster. If the program used to determine the hit is HMMSW then the score refers to HMMSW score.

P-Value

The entries in the P-Value column refer to the probability that such matches occur by chance.

%Ident

The entries in the “%Ident” column of the table refer to the percentage of identically matched nucleotides (or residues) that exist along the length of that portion of the sequences which is aligned by the BLAST comparison to generate the statistical scores presented. This entry is omitted when a clone is included solely due to its association with a particular cluster.